

**A Preliminary Classification of Plant
Communities in the Central Parkland
Natural Sub-Region of Alberta.**

Prepared for:

Alberta Sustainable Resource Development
Public Lands Division
Resource Data Branch
Edmonton, Alberta

Prepared by:

Matthew Wheatley
and
Jerry Bentz

Geowest Environmental Consultants Ltd.
Suite 203 4208 – 97th Street
Edmonton, Alberta, T5K 2J6

February 2002

TABLE OF CONTENTS

Objectives 1

Study Area Background 1

Methods 2

Results 5

Dwarf Shrubland 5

Evergreen 5

Broad-leaved 5

Evergreen dwarf-shrubland 5

ARCTOSTAPHYLOS UVA-URSI DWARF-SHRUBLAND ALLIANCE 5

Arctostaphylos uva-ursi 5

Scale-leaved 6

Xeric evergreen dwarf-shrubland 6

JUNIPERUS HORIZONTALIS DWARF-SHRUBLAND ALLIANCE 6

Juniperus horizontalis 6

Juniperus horizontalis - Selaginella densa - Calamovilfa longifolia 7

Forest/Woodland 7

Deciduous 7

Broad-leaved 7

Intermittently flooded cold-deciduous forest/woodland 7

ALUNUS TENUIFOLIA FOREST/WOODLAND ALLIANCE 8

Alnus tenuifolia - Salix discolor / Lonicera involucrata - Cornus stolonifera - Ribes hudsonianum / Plagiomnium cuspidatum 8

Cold-deciduous forest/woodland 8

ACER NEGUNDO FOREST/WOODLAND ALLIANCE 8

Acer negundo / Prunus virginiana 8

BETULA NEOALASKANA – POPULUS BALSAMIFERA FOREST/WOODLAND ALLIANCE 9

Betula neoalaskana - Populus balsamifera / Cornus stolonifera / Calamagrostis canadensis 9

POPULUS BALSAMIFERA FOREST/WOODLAND ALLIANCE 10

Populus balsamifera / Salix discolor - (Betula occidentalis) 10

Populus balsamifera / Salix petiolaris 10

Populus balsamifera / Viburnum opulus - Cornus stolonifera / Aralia nudicaulis 11

POPULUS TREMULIODES FOREST/WOODLAND ALLIANCE 11

Populus tremuloides / (sparse understory) 11

Populus tremuloides / Corylus cornuta / Low Shrubs and Herbs 12

Populus tremuloides / Prunus virginiana – (Shrub) / Low Herb 13

Populus tremuloides / Prunus virginiana / Solidago canadensis 14

Populus tremuloides / Rosa acicularis - Rubus idaeus 15

Populus tremuloides / Symphoricarpos occidentalis / Herbs 16

Populus tremuloides / Viola adunca - Orthilia secunda 16

<i>POPULUS TREMULOIDES</i> – <i>POPULUS BALSAMIFERA</i>	
FOREST/WOODLAND ALLIANCE	17
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> - <i>Betula papyrifera</i> /	
<i>Symphoricarpos occidentalis</i>	17
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> / <i>Amelanchier alnifolia</i>	17
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> / <i>Cornus stolonifera</i> - <i>Rosa</i>	
<i>acicularis</i> - <i>Viburnum edule</i> / <i>Aralia nudicaulis</i>	18
Evergreen	18
Needle-leaved	18
Saturated evergreen forest/woodland	18
<i>PICEA GLAUCA</i> SATURATED FOREST/WOODLAND ALLIANCE.....	18
<i>Picea glauca</i> / <i>Cornus stolonifera</i> / <i>Climacium dendroides</i>	18
<i>PICEA MARIANA</i> SATURATED FOREST/WOODLAND ALLIANCE	19
<i>Picea mariana</i> / <i>Ledum groenlandicum</i> / Low Shrubs and Moss.....	19
<i>PINUS BANKSIANA</i> FOREST/WOODLAND ALLIANCE	20
<i>Pinus banksiana</i> / <i>Arctostaphylos uva-ursi</i>	20
Mixed Evergreen-deciduous	20
Broad-leaved	20
Mixed-evergreen-deciduous cold-deciduous forest/woodland	20
<i>PICEA GLAUCA</i> MIXED-EVERGREEN DECIDUOUS	
FOREST/WOODLAND ALLIANCE	20
<i>Picea glauca</i> - <i>Populus tremuloides</i> - <i>Populus balsamifera</i> / (shrubs and	
herbs)	20
<i>POPULUS TREMULOIDES</i> MIXED-EVERGREEN DECIDUOUS	
FOREST/WOODLAND ALLIANCE	21
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> - <i>Picea glauca</i> / <i>Rubus idaeus</i> ...21	
<i>Populus tremuloides</i> (<i>Picea glauca</i>) / <i>Aralia nudicaulis</i> / <i>Cornus canadensis</i>	
.....	22
Herbaceous	22
Annual	22
Narrow-leaved	22
Riparian herbaceous vegetation	22
<i>SENECIO CONGESTUS</i> HERBACEOUS ALLIANCE	22
<i>Senecio congestus</i>	22
Herb	23
Thread-like leaved	23
Intermittently flooded brackish herbaceous vegetation	23
<i>RUPPIA MARITIMA</i> INTERMITTENTLY FLOODED HERBACEOUS	
ALLIANCE	23
<i>Ruppia maritima</i>	23
Toothed, broad leaves	24
Brackish herbaceous vegetation	24
<i>POTENTILLA ANSERINA</i> HERBACEOUS ALLIANCE.....	24
<i>Potentilla anserina</i> - <i>Distichlis stricta</i>	24
Sedges and Rushes	24
Spiked Sedges	24

Marl fen herbaceous vegetation	24
<i>ELEOCHARIS QUINQUEFOLIA</i> HERBACEOUS ALLIANCE.....	24
<i>Eleocharis quinquefolia</i>	24
Temperate or subpolar sedge grassland	25
<i>CAREX SPP. SEDGE</i> GRASSLAND ALLIANCE.....	25
<i>Carex</i> group - <i>Calamovilfa longifolia</i>	25
<i>Carex</i> group - <i>Koeleria macrantha</i>	26
<i>Carex</i> group - <i>Stipa curtisetata</i>	26
Temperate or subpolar intermittently flooded sedges and rushes	27
<i>CAREX ATHERODES</i> INTERMITTENTLY FLOODED HERBACEOUS ALLIANCE.....	27
<i>Carex atherodes</i> - <i>Carex rostrata</i>	27
Temperate or subpolar permanently-flooded sedges and rushes	27
<i>CAREX SPP. PERMANENTLY FLOODED</i> HERBACEOUS ALLIANCE.....	27
<i>Carex atherodes</i> - <i>Scolochloa festucacea</i> - <i>Eleocharis palustris</i> - <i>Typha latifolia</i>	27
<i>Carex pseudo-cyperus</i> - <i>Calla palustris</i>	28
Temperate or subpolar seasonally flooded sedges and rushes	28
<i>CAREX ATHERODES</i> SEASONALLY-FLOODED HERBACEOUS ALLIANCE.....	28
<i>Carex atherodes</i> - <i>Calamagrostis inexpansa</i>	28
<i>Carex atherodes</i> - <i>Scolochloa festucacea</i> - <i>Poa palustris</i> - <i>Cirsium arvense</i>	29
<i>ELEOCHARIS PALUSTRIS</i> SEASONALLY FLOODED HERBACEOUS ALLIANCE.....	29
<i>Eleocharis palustris</i>	29
Temperate or subpolar temporarily flooded sedges and rushes	30
<i>CAREX ATHERODES</i> TEMPORARILY FLOODED HERBACEOUS ALLIANCE.....	30
<i>Carex atherodes</i> / <i>Lemna minor</i>	30
Temperate or subpolar semipermanently flooded sedges and rushes	30
<i>CAREX AQUATILIS</i> SEMIPERMANENTLY FLOODED HERBACEOUS ALLIANCE.....	30
<i>Carex aquatilis</i> - <i>Carex utriculata</i>	30
<i>SCIRPUS PALUDOSIS</i> PERMANENTLY FLOODED HERBACEOUS ALLIANCE.....	31
<i>Scirpus paludosus</i>	31
<i>SCIRPUS SPP. SEMIPERMANENTLY FLOODED</i> HERBACEOUS ALLIANCE.....	32
<i>Scirpus acutus</i>	32
<i>Scirpus paludosus</i> - <i>Puccinellia nuttalliana</i> - <i>Hordeum jubatum</i> - <i>Distichlis stricta</i>	32
<i>Scirpus acutus</i> – Liverworts.....	33
<i>Scirpus pungens</i>	33
<i>Scirpus pungens</i> - <i>Hordeum jubatum</i> - <i>Puccinellia nuttalliana</i>	33
Temperate or subpolar sedges or rushes on sandy substrate	34
<i>CAREX SPP. RAPIDLY DRAINED</i> HERBACEOUS ALLIANCE.....	34

<i>Carex siccata</i> - <i>Calamovilfa longifolia</i>	34
Aquatic Rushes	35
Temperate or subpolar semipermanently flooded aquatic rushes	35
TYPHA LATIFOLIA SEMIPERMANENTLY FLOODED HERBACEOUS	
ALLIANCE	35
<i>Typha latifolia</i>	35
Graminoid	35
Linear, parallel-veined leaves	35
Temperate or subpolar grasslands	35
AGROPYRON TRACHYCAULUM HERBACEOUS ALLIANCE	35
<i>Agropyron trachycaulum</i> - <i>Hordeum jubatum</i> - <i>Elymus canadense</i>	35
AGROSTIS SCABRA HERBACEOUS ALLIANCE	36
<i>Agrostis scabra</i> - <i>Achillea millefolium</i> - <i>Antennaria parvifolia</i>	36
BOUTELOUA GRACILIS HERBACEOUS ALLIANCE	36
<i>Bouteloua gracilis</i> - <i>Artemisia frigida</i>	36
CALAMAGROSTIS SPP. HERBACEOUS ALLIANCE	37
<i>Calamagrostis canadensis</i>	37
<i>Calamagrostis stricta</i> - <i>Juncus balticus</i> - <i>Potentilla anserina</i> - <i>Aster hesperius</i>	
.....	37
DISTICHLIS STRICTA HERBACEOUS ALLIANCE	38
<i>Distichlis stricta</i>	38
<i>Distichlis stricta</i> - <i>Agropyron smithii</i>	39
ELYMUS CANADENSIS HERBACEOUS ALLIANCE	39
<i>Elymus canadensis</i> - <i>Poa pratensis</i> - <i>Glycyrrhiza lepidota</i>	39
KOELERIA MACRANTHA HERBACEOUS ALLIANCE	40
<i>Koeleria macrantha</i> - <i>Agropyron smithii</i>	40
STIPA SPP. HERBACEOUS ALLIANCE	40
<i>Stipa comata</i> - <i>Koeleria macrantha</i> - <i>Agropyron smithii</i>	40
<i>Stipa comata</i> - <i>Koeleria macrantha</i> - <i>Agropyron trachycaulum</i> - <i>Agropyron</i>	
<i>smithii</i> - <i>Carex obtusata</i>	41
<i>Stipa comata</i> - <i>Poa sandbergii</i> - <i>Carex obtusata</i>	41
<i>Stipa comata</i> - <i>Stipa viridula</i> - <i>Festuca hallii</i>	41
<i>Stipa curtisetata</i> - <i>Artemisia frigida</i>	42
<i>Stipa curtisetata</i> - <i>Bouteloua gracilis</i>	42
<i>Stipa viridula</i> - <i>Poa sandbergii</i> - <i>Carex obtusata</i>	43
Temperate or subpolar grasslands on sandy substrate	43
CALAMOVILFA LONGIFOLIA SANDY HERBACEOUS ALLIANCE	43
<i>Calamovilfa longifolia</i> - <i>Artemisia frigida</i> - <i>Koeleria macrantha</i> - <i>Festuca</i>	
<i>saximontana</i> - <i>Festuca hallii</i> / <i>Selaginella densa</i>	43
MUHLENBERGIA ASPERIFOLIA SANDY HERBACEOUS ALLIANCE	44
<i>Muhlenbergia asperifolia</i> - <i>Scirpus nevadensis</i> - <i>Distichlis stricta</i>	44
PUCCINELLIA NUTTALLIANA SANDY HERBACEOUS ALLIANCE	44
<i>Puccinellia nuttalliana</i> - <i>Distichlis stricta</i>	44
SPARTINA GRACILIS HERBACEOUS ALLIANCE	45
<i>Spartina gracilis</i> - <i>Agropyron smithii</i>	45
STIPA CURTISETA SANDY HERBACEOUS ALLIANCE	45

<i>Stipa curtisetata</i> - <i>Koeleria macrantha</i> - <i>Agropyron trachycaulum</i> / <i>Cladina mitis</i>	45
Temperate or subpolar seasonally flooded grasslands	45
HORDEUM JUBATUM SEASONALLY FLOODED HERBACEOUS ALLIANCE	46
<i>Hordeum jubatum</i>	46
PUCCINELLIA NUTTALLIANA HERBACEOUS ALLIANCE	46
<i>Puccinellia nuttalliana</i>	46
PHALARIS ARUNDINACEA HERBACEOUS ALLIANCE	47
<i>Phalaris arundinacea</i> - <i>Carex atherodes</i> - <i>Calamagrostis stricta</i>	47
Temperate or subpolar intermittently flooded grasslands	47
CALAMAGROSTIS INEXPANSA INTERMITTENTLY FLOODED HERBACEOUS ALLIANCE	48
<i>Calamagrostis inexpansa</i> - <i>Juncus balticus</i>	48
<i>Hordeum jubatum</i> - <i>Puccinellia nuttalliana</i>	48
Composites	48
Oblong, spatulate-leaved	48
Temperate or subpolar composite grasslands	49
ANTENNARIA PARVIFOLIA HERBACEOUS ALLIANCE	49
<i>Antennaria parvifolia</i> - <i>Artemisia frigida</i> - <i>Koeleria macrantha</i>	49
Twice-pinnate leaves	49
Temperate or subpolar composite grasslands	49
ARTEMISIA FRIGIDA HERBACEOUS ALLIANCE	49
<i>Artemisia frigida</i> / <i>Koeleria macrantha</i> - <i>Carex obtusata</i>	49
Leaves large, petioled	50
Temperate or subpolar permanently flooded composite wetlands	50
POLYGONUM AMPHIBIUM PERMANENTLY FLOODED HERBACEOUS ALLIANCE	50
<i>Polygonum amphibium</i>	50
Shrubland	50
Deciduous	50
Broad-leaved	50
Temperate or subpolar cold-deciduous shrubland	50
ELAEAGNUS COMMUTATA SHRUBLAND ALLIANCE	50
<i>Elaeagnus commutata</i>	50
<i>Elaeagnus commutata</i> / <i>Symphoricarpos occidentalis</i> - <i>Rosa woodsii</i> / <i>Poa palustris</i>	51
POPULUS TREMULOIDES SHRUBLAND ALLIANCE	52
<i>Populus tremuloides</i> - <i>Amelanchier alnifolia</i> - <i>Prunus virginiana</i> / <i>Rosa spp.</i>	52
PRUNUS VIRGINIANA SHRUBLAND ALLIANCE	52
<i>Prunus virginiana</i> - <i>Amelanchier alnifolia</i> / <i>Agropyron trachycaulum</i> - <i>Poa pratensis</i>	52
ROSA ACCICULARIS SHRUBLAND ALLIANCE	53
<i>Rosa acicularis</i> – (shrub) / <i>Solidago spp.</i>	53
<i>Rosa acicularis</i> - <i>Rubus idaeus</i> / <i>Lathyrus venosus</i>	53

<i>SALIX SPP.</i> SHRUBLAND ALLIANCE	53
<i>Salix petiolaris</i> / (low shrubs and herbs)	53
<i>SYMPHORICARPOS OCCIDENTALIS</i> SHRUBLAND ALLIANCE	54
<i>Symphoricarpos occidentalis</i>	54
<i>Symphoricarpos occidentalis</i> / <i>Antennaria parvifolia</i> - <i>Festuca hallii</i>	55
Temperate or subpolar permanently flooded cold-deciduous shrubland	56
<i>BETULA PUMILA</i> PERMANENTLY FLOODED SHRUBLAND ALLIANCE	56
<i>Betula pumila</i> / <i>Carex aquatilis</i>	56
<i>SALIX PYRIFOLIA</i> PERMANENTLY FLOODED SHRUBLAND ALLIANCE	56
<i>Salix pyrifolia</i> - <i>Betula nealaskana</i> / <i>Ledum groenlandicum</i> - <i>Calamagrostis canadensis</i> / <i>Sphagnum warnstorffii</i>	56
<i>Salix pyrifolia</i> / <i>Carex utriculata</i>	57
Temperate or subpolar semi-permanently flooded cold-deciduous shrubland	57
<i>SALIX PETIOLARIS</i> SEMI-PERMANENTLY FLOODED SHRUBLAND	57
<i>Salix petiolaris</i> - <i>Salix planifolia</i> / <i>Carex utriculata</i>	57
<i>Salix petiolaris</i> / <i>Juncus balticus</i>	57
Temperate or subpolar cold-deciduous shrubland on sandy substrate	58
<i>POPULUS TREMULOIDES</i> SANDY SHRUBLAND ALLIANCE	58
<i>Populus tremuloides</i> / <i>Juniperus horizontalis</i> / <i>Cladina mitis</i>	58
<i>SYMPHORICARPOS OCCIDENTALIS</i> SANDY SHRUBLAND ALLIANCE	58
<i>Symphoricarpos occidentalis</i> - <i>Rosa arkansana</i> / <i>Juniperus horizontalis</i>	58
Sparse	59
Xeric sandy soil	59
Dunes and blowouts	59
Temperate or subpolar sparse graminoids on sand	59
<i>Calamovilfa longifolia</i> - <i>Sporobolus cryptandrus</i> - <i>Koeleria macrantha</i> - <i>Carex obtusata</i>	59
Temperate or subpolar sparse cold-deciduous shrubs on sand	59
<i>Populus tremuloides</i> / <i>Juniperus horizontalis</i>	59
<i>Populus tremuloides</i> - <i>Prunus virginiana</i> / <i>Juniperus horizontalis</i> / <i>Selaginella densa</i>	60
Temperate or subpolar sparse evergreen shrubs on sand	60
<i>Juniperus horizontalis</i> / (<i>Koeleria macrantha</i>) / <i>Cladina mitis</i>	60
Lake and slough edges	60
Saline wetlands	60
Saline wetland sparse herbaceous vegetation	60
<i>Salicornia europaea</i>	60
<i>Scirpus nevadensis</i>	61
<i>Scirpus nevadensis</i> - (<i>Triglochin maritima</i>)	61
<i>Suaeda calceoliformis</i>	62
<i>Triglochin maritima</i>	62
Crosswalk Table – Literature Comparisons.	63
Discussion	77

Personal Communications 78
Literature Cited..... 78

Objectives

The objective of this report was to produce a synopsis of historical data on Central Parkland natural/near natural plant community types, and to create a preliminary provincial classification of these types. In addition, each community was to be assigned a preliminary provincial conservation rank based on available information or expert opinion, and comparisons with similar plant communities found in adjacent provinces and states were to be made. There was no field component in this process, all conclusions were based on a review and synthesis of the best available information.

The Resource Data Division of Alberta Environment contracted Geowest Environmental Consultants Ltd. to compile available information on plant community types within the Central Parkland natural sub-region of Alberta, excluding *Festuca spp.*, *Agropyron spp.*, and *Artemisia cana* communities which are included in other reports (Weerstra and Weerstra 1998, Vujnovic and Bentz 2001, and Weerstra 2001, respectively). This initiative is in support of the Alberta Natural Heritage Information Center (ANHIC). ANHIC collects, evaluates and makes available information on elements of natural biodiversity of Alberta, including flora, fauna and natural plant communities. ANHIC develops tracking lists of elements that are considered of high priority because they are considered rare or special in some way. ANHIC's long-term goal is to develop a list of plant community types which occur throughout the province and to attempt to identify community types that require conservation initiatives.

Study Area Background

Parkland

The Parkland Region extends from northern Minnesota northeastward through southern Manitoba, central Saskatchewan, and Alberta (Bird 1961). It is also found along the Rocky Mountain Foothills and in the Peace River Area of Alberta, and in Glacier County, Montana (Bird 1961, Lynch 1955). Alberta's Parkland Region can be divided into three natural subregions; the Central, the Foothills, and the Peace River subregions, all part of a landscape classification system consisting of six regions and 20 subregions throughout the province (Alberta Environmental Protection 1994b). The Parkland Natural Region forms a transition zone between the drier grasslands of the plains and the mixedwood and coniferous forests of the Boreal Forest and the Rocky Mountains (Alberta Environmental Protection 1997).

Central Parkland

The Central Parkland Natural Subregion occurs between the grasslands of southern Alberta and the boreal forest to the north. Its northern boarder runs from Edmonton to the Saskatchewan border, roughly following the North Saskatchewan River, while its southern border arcs from Calgary northeast, just south of Stettler, to the Saskatchewan border. The region is a mixture of vegetation and landforms transitional to the boreal and grassland regions. Surficial deposits include hummocky and ground moraines, glaciolacustrine deposits, coarse outwash, kame moraines, and dune fields (Alberta Environmental Protection

1997). Associated plant communities in this subregion can include a mixture of grasslands, mixed deciduous and mature aspen forests, saline wetlands, shrublands extending out from forested areas, and sparse communities stabilizing sand dune slopes. Most of the natural elements of the Parkland Region are now limited to marginal lands or ecological reserves (e.g. Rumsey and Wainwright). The majority of the region has been permanently altered from pre-settlement conditions primarily by agriculture, oil and gas development, and cattle farming.

This area is highly productive in agriculture and oil and gas production, two industries that are visibly prominent across the subregion's landscape. The Central Parkland is the most densely populated region of the province and the majority of the Parkland's native vegetation has been lost to human development. Between the mid 1800s and present day it is estimated that 85-95% of the Parkland Region has been lost to cultivation, urbanization, roads and other anthropogenic development (IISD 1994). Despite the historically high human presence the Central Parkland, many native vegetation community types were not adequately documented before they were eliminated or greatly reduced. The area of uncultivated Parkland continues to decline today (Alberta Environmental Protection 1997). Therefore, Alberta Environment has deemed it a priority to compile and organize existing information on native or near-native vegetation communities in the Central Parkland natural subregion.

Methods

Vegetation Community Information Collection

Information provided in this report was generated by a literature review. Discussions with researchers familiar with and experienced in sampling vegetation communities within the Central Parkland natural subregion of Alberta were also incorporated. There was no field component to this study.

Community Classification System

Community classification followed the hierarchical guidelines set forth by Grossman *et al.* (1998). Community types were grouped first by physiognomic levels, and second by floristic levels. Physiognomic groupings consist of five levels: Class, Subclass, Group, Subgroup, and Formation.

Physiognomic Levels:

Class: Community class is based on the vegetation structure of the dominant, uppermost life form and contains six different levels:

1. Forest/Woodland – Trees with crowns overlapping (25-100% cover)
2. Shrubland – Shrubs generally >0.5m tall forming >25% cover
3. Dwarf-Shrubland – Shrubs <0.5m tall forming >25% cover
4. Herbaceous – Graminoids, ferns and forbs dominant
5. Nonvascular – bryophytes, lichens and algae dominant
6. Sparse – Abiotic substrate dominant

Subclass: Community subclass is based upon the growth-form characteristics of the dominant life form, predominantly leaf phenology for classes 1-3 (e.g. evergreen, deciduous, mixed evergreen/deciduous), persistence for class 4 (perennial, annual), and substrate characteristics for class 6 (e.g. rocks, sand, exposed lakeshore). Class 5 is not represented in this report.

Group: Community groups are defined based on leaf characters (broad leaf, needle, scale, etc.). The presence of woody strata separates groups in the Herbaceous and Nonvascular classes, and Sparse vegetation communities are separated on the basis of topographic position (shores, cliffs, etc.).

Subgroup: Community subgroups divide each group into either a Natural/Seminatural or a Cultural subgroup. All plant communities discussed in this report are in the Natural/Seminatural subgroup.

Formation: Community formations represent vegetation types that share a definite physiognomy or structure within broadly defined environmental factors, landscape positions, or hydrological regimes (e.g. Temperate or subpolar cold-deciduous shrubland).

Floristic Levels:

Two floristic levels are used: the Alliance and the Association. Both are based on the dominant species in the community.

Alliance: The alliance is a physiognomically uniform group of plant associations (see below) sharing one or more diagnostic species, which as a rule are found in the uppermost stratum of the vegetation (Grossman *et al.* 1998).

Association: The association is the lowest level of the hierarchy and is defined as a plant community type of definite floristic composition, uniform habitat conditions, and uniform physiognomy (Grossman *et al.* 1998). Nomenclature at this level is based on the diagnostic species. Species occurring in the uppermost strata are listed first (separated by a hyphen if in the same strata, and a slash if in separate strata) followed successively by those occurring in lower levels. Within the same stratum, the order of species names generally reflects decreasing levels of dominance or constancy (Grossman *et al.* 1998).

Estimating Conservation Preliminary Ranks

Each community was given a preliminary provincial conservation rank based on available occurrence or abundance information, or expert opinion. The ranking system used is based on The Nature Conservancy's species ranking system (Grossman *et al.* 1994). The two major criteria used in ranking communities are the total number of occurrences and the total acreage or area of each community. Trends in range expansion and immediate threats to

community persistence are considered in ranking as well. Preliminary Ranks range from S1 to S5, from rare or threatened to widespread, respectively (Table 1).

The information used as rationale behind selected ranks was provided from both the published literature (if available) and professional opinion. The ranks generated using these methods are highly subjective. Most published community-level vegetation descriptions applicable to the Central Parkland Natural Subregion of Alberta are qualitative and without abundance measurements, and professional opinion in the absence of empirical support is tenuous at best. The rankings provided in this report are as such considered preliminary and should be consistently under review as new, quantitative information is acquired.

Table 1. Provincial conservation ranks and definitions.

Preliminary Rank	Criteria
S1	Five or fewer occurrences, very few remaining hectares, especially vulnerable to extirpation.
S2	Six to 20 occurrences, few remaining hectares, vulnerable to extirpation throughout its range.
S3	21 to 100 occurrences; may be rare and local throughout its range or found locally, even abundantly, in a restricted range, or vulnerable to extirpation throughout its range because of some specific factor.
S4	Uncommon, but not rare, although it may be quite rare in parts of its range, especially at the periphery; apparently not vulnerable in most of its range.
S5	Common, widespread and abundant provincially; although it may be quite rare in parts of its range, especially at the periphery; not vulnerable in most of its range.
SU	Not able to rank; status is uncertain.
SH	Historic; presumed eliminated in the province with little or no likelihood it will be rediscovered; there may be potential for restoration.
SX	Believed to be eliminated throughout the province with virtually no likelihood that it will be rediscovered; no restoration potential due to extinction of dominant or characteristic species.
SP	Potentially exists; further documentation required.
S?	Element is not yet ranked.

Where information was available, plant communities from the Central Parkland of Alberta were compared with any similar communities found within the subregion, in other regions of the province, or from other provinces and states.

For plant communities containing one or a combination of the following ranks; S1, S2, S3, or SU, an Element Characterization Abstract was completed in a format provided by ANHIC. The abstract format included scientific and common name, physiognomic class, range and distribution information, environmental description, vegetation description, proposed conservation rank and justification, general comments and relevant bibliographic references.

Results

Overview

A total of 106 plant communities occurring and described within the Central Parkland natural subregion were derived from the literature, unpublished data and expert opinion. The majority of the communities were ranked S5 or SU. Community types included representatives from all classes except Nonvascular. They are presented below by major physiognomic level.

Dwarf Shrubland

Evergreen

Broad-leaved

Evergreen dwarf-shrubland

ARCTOSTAPHYLOS UVA-URSI DWARF-SHRUBLAND ALLIANCE

Arctostaphylos uva-ursi

Common Bearberry

Preliminary Rank: S5

In the Central Parkland, this community type is described from one site in the Dillberry Lake Provincial Park area of Alberta by Meijer and Karpuk (1999). This community occurs on Orthic Regosolic soils where drainage is rapid, moisture conditions are xeric, and nutrients are submesotrophic. It forms communities on sandy, upland plains in parkland regions in areas where very little soil formation is evident and forbs comprise a higher cover than grass. Common associates for this Parkland community include *Juniperus horizontalis*, *Heterotheca villosa*, *Solidago missouriensis*, *Artemisia frigida*, *Artemisia campestris*, *Calamovilfa longifolia* and *Agropyron dasystachyum*. Because this type is only described from one site, an assessment of variability between occurrences in the Central Parkland is not available.

Remphrey *et al.* (1983) note that the occurrence of *A. uva-ursi* is common on sand dune areas of sub-boreal regions of Canada. In the southern boreal forests of Saskatchewan and Manitoba, bearberry is characteristic of dry and very dry forests (Moss 1955, Beckingham *et al.* 1996). This species is

common in Alberta particularly in the foothills and mountain regions and, when present, often forms extensive, low-growing patches (Moss 1983) which can be considered bearberry communities. However, the status of this species as a community and not an associated understory species in many cases is not clear. More work is needed to clarify its distribution as a community type within the Central Parkland.

Elsewhere, Bearberry is a dominant understory species that grows best in high light situations where it forms a compact and intricate mat (Natureserve 2001). Bearberry forms low-growing communities in open pine forests under *Pinus banksiana* and *Pinus contorta* in boreal regions (Moss 1955, Beckingham and Archibald 1996) where it is often an extensive understory associate of conifer forest communities (but is otherwise very similar to the Central Parkland *A. uva-ursi* type). Douglas and Bliss (1977) describe an alpine *A. uva-ursi* dwarf-shrubland community in the western and eastern Cascade Mountains of Washington State as occurring in coniferous forest and alpine meadows (ranked S3 by The Nature Conservancy). This is an alpine community type that does not occur on a sand substrate, characteristics that differentiate this community from Alberta Central Parkland types.

(Dwarf shrubland)

(Evergreen)

Scale-leaved

Xeric evergreen dwarf-shrubland

JUNIPERUS HORIZONTALIS DWARF-SHRUBLAND ALLIANCE

Juniperus horizontalis

Creeping Juniper

Preliminary Rank: SU

In the Central Parkland, this community has been described from 11 sites in the Dillberry Lake Provincial Park area of Alberta by Meijer and Karpuk (1999). In this subregion it occurs on steep, north-east and south-facing slopes, on well drained, glaciofluvial or eolian sites with Orthic Dark Brown or Orthic Regosolic soils with xeric moisture regimes and submesotrophic nutrient conditions. A sparse association of forbs and grasses can include *Heterotheca villosa*, *Thermopsis rhombifolia*, *Comandra umbellata*, *Stipa curtisetata* (and *comata*) and *Agropyron trachycaulum*. Under steep, dry, and nutrient poor conditions this community grows low and mat-like as it stabilizes exposed mineral soils. An assessment of variability between occurrences for this community in the Central Parkland is not available in the literature.

Juniperus horizontalis is documented from sandy and rocky areas in the foothills and mountains of Alberta (Moss 1983, Johnson *et al.* 1995) but its

occurrence as a community type in these areas is not clear. Associated species, or lack thereof, differentiate this community from the *Juniperus horizontalis* - *Selaginella densa* - *Calamovilfa longifolia* described by Pearson Timberline Consultants (1993). Creeping juniper is an indicator in a number of grassland or drier coniferous forest habitat types of the northern Great Plains where it occurs most commonly as a co-dominant (Johnston 1987), but is not documented as a community type. *Juniperus horizontalis* is described as a dominant or an associate of several plant communities in the Central Parkland, but more work is needed to determine its abundance as a community type in this subregion.

Juniperus horizontalis - *Selaginella densa* - *Calamovilfa longifolia*
Creeping juniper – Prairie selaginella – Sand grass
Preliminary Rank: S4

In the Central Parkland this community type is described from one area in the Wainwright Dunes Ecological Reserve (Pearson Timberline Consultants 1993). This community inhabits dry outwash plains, dune complexes, and kame moraines on grassland areas in the Central Parkland (Pearson Timberline Consultants 1993). It is associated with variable aspect and slope, on rapidly drained areas with xeric to subxeric moisture regimes, and Orthic Regosolic soils. Other shrubs present at lower levels can be *Rosa spp.* and *Prunus virginiana*. *Artemisia frigida* and *Festuca hallii* can also be present. This community is area-restricted in the Central Parkland to unbroken, sandy areas but it is not considered rare (A.H. Weerstra pers comm.). An assessment of variability between occurrences for this community in the Central Parkland is not available in the literature.

A similar community type, *Juniperus horizontalis* / *Koeleria macrantha* - *Artemisia frigida* – *Selaginella densa*, with similar dominants, similar associated species and the same site conditions is described by Thorpe and Goodwin (1993) from the Manito Sand Hills in Saskatchewan. *Festuca hallii* is a common associate of this community type. The main difference of this community to the one described above is the presence of *Koeleria macrantha* as a dominant. There is no available assessment of the constancy of this species as a dominant for this community type, a characteristic that could differentiate these as different types. The distribution of this Saskatchewan type is not clear.

Forest/Woodland

Deciduous

Broad-leaved

Intermittently flooded cold-deciduous forest/woodland

ALUNUS TENUIFOLIA FOREST/WOODLAND ALLIANCE

Alnus tenuifolia - *Salix discolor* / *Lonicera involucrata* - *Cornus stolonifera* - *Ribes hudsonianum* / *Plagiomnium cuspidatum*

River alder – Pussy willow / Bracted honeysuckle – Red osier dogwood – Wild black currant / Woodsy mniium

Preliminary Rank: S5

Griffiths *et al.* (1997) describe this community from one area in the northern Central Parkland of Alberta. It is described as an alder swamp dominated by river alder of tree size, under very wet, intermittently flooded conditions, especially on bog margins and along stream channels (Griffiths *et al.* 1997). Nutrient conditions and soil types for this community are not discussed in the literature. Common associates include *Salix discolor* (also of tree size), *Lonicera involucrata*, *Cornus stolonifera*, *Ribes hudsonianum*, *Caltha palustris*, *Viola palustris* and *Calamagrostis canadensis*. Moss coverage (*Plagiomnium cuspidatum*) is extensive. An assessment of variability between occurrences for this community in the Central Parkland is not available in the literature.

Beckingham and Archibald (1996) describe a similar community from the Boreal Region of Alberta as occurring in lower slope, moist areas (the d1.2 horsetail Pb-Bw community type). However, it includes *Populus balsamifera* as a co-dominant species, lacks *Lonicera involucrata* and the moss component is primarily *Brachythecium spp.* We have observed this community in riparian areas of the northern Central Parkland and the Boreal Region of Alberta. We do not consider it rare, even though a formal assessment of its provincial abundance is not available.

Cold-deciduous forest/woodland

ACER NEGUNDO FOREST/WOODLAND ALLIANCE

Acer negundo / *Prunus virginiana*

Manitoba Maple / Choke cherry Forest

Preliminary Rank: S1S2

This community is not formally described in the literature for the Central Parkland of Alberta. However, it is known to occur in coulees and on protected slopes, primarily in the Grassland Natural Region, and extending into the Central Parkland Subregion (Allen 2001). A tall shrub layer of *Prunus virginiana* is characteristic of undisturbed stands, but can be replaced by a low shrub layer of *Symphoricarpos occidentalis* with disturbance such as cattle grazing. Soil type, nutrient conditions and moisture conditions are not described for this community. Associated species of undisturbed stands can

include *Carex sprengelii* and *Glyceria striata*, which are replaced by weedy species under disturbance. Because this type has not been formally described, an assessment of variability between occurrences in the Central Parkland is not available.

This community is included in the ANHIC plant community tracking list (Allen 2001). Manitoba Maple is widely planted but uncommon as a native community in Alberta. It is possibly native along streams in south-eastern Alberta (Moss 1983), but restricted to river floodplains in this area. More work is needed to document the locations and extent of this community in the province.

Wallis (1977) describes a riparian Manitoba Maple community in the Dry Mixedgrass Natural Subregion of Alberta known particularly from the South Saskatchewan and Red Deer River valleys. Manitoba Maple is a component of various deciduous forest plant communities in the Great Plains of North America (Hansen and Hoffman 1988), and in Arizona and New Mexico. Manitoba Maple is the overstory dominant in several high elevation riparian forests. This species is widespread in riparian and palustrine communities throughout most of the contiguous United States (Little 1979). Natureserve (2001) provides information only on a riparian floodplain *Acer negundo* / *Prunus virginiana* community in the western Great Plains of the United States. This community has similar associated species in addition to *Prunus virginiana* (*Rhus trilobata*, *Symphoricarpos occidentalis*, *Ribes aureum*) but would differ from Central Parkland types in substrate and topographic position (floodplain versus coulee bottoms). Throughout much of the range of *Acer negundo* there are no described Manitoba Maple communities.

BETULA NEOALASKANA – POPULUS BALSAMIFERA FOREST/WOODLAND ALLIANCE

Betula neoalaskana - *Populus balsamifera* / *Cornus stolonifera* / *Calamagrostis canadensis*

Alaskan Birch – Balsam Poplar / Red osier dogwood / Marsh reed grass

Preliminary Rank: SU

In the Central Parkland, Griffiths and Griffiths (1987a) describe this community from one site near the Sylvan Lake area of Alberta. This community occurs on lower ground where the *Populus tremuloides* community gives way to a moister forest and *Betula neoalaskana* and *Populus balsamifera* become dominant (Griffiths and Griffiths 1987a). Soils, hydrological conditions and nutrient regimes are not discussed. The understory of this forest type is heterogeneous. Common associates include *Cornus stolonifera*, *Rubus idaeus*, *Ribes oxyacanthoides*, *Calamagrostis canadensis*, *Heracleum lanatum* and *Aralia nudicaulis*. Variation in the understory can occur along streams and seepage channels where *Alnus*

tenuifolia, *Caltha palustris*, and *Carex bebbii*, among others can also occur. Because this type is only described from one site, a quantitative assessment of variability between occurrences is not available. Little is documented on Alaskan Birch communities in Alberta, so the distribution and preliminary rank of this community is unknown.

According to Griffiths and Griffiths (1987a), this community is comparable to the "balsam poplar consociation" described by Moss (1932) who did not discriminate based the presence of *Betula*. It is also similar to the *Betulo-Populetum balsamiferae* association described from similar, mesic site conditions by Looman (1987b). However, Looman's (1987b) association includes trace amounts of *Rhus radicans* and *Bromus ciliatus*. He formed alliances based on the presence of *Betula*, but makes no reference to association abundance.

POPULUS BALSAMIFERA FOREST/WOODLAND ALLIANCE

Populus balsamifera / *Salix discolor* - (*Betula occidentalis*)

Balsam Poplar / Pussy willow – (Water birch)

Preliminary Rank: SU

In the Central Parkland, Meijer and Karpuk (1999) describe this community type from two sites in the Dillberry Lake Provincial Park area. This "balsam woodland" community occurs on lacustrine deposits adjacent to wetlands (lake perimeters) on wet and nutrient rich substrates with a continuous moisture source (Meijer and Karpuk 1999). Soils for this community range from Rego Gleysols to Orthic and Gleyed Regosols. Common associates include *Hordeum jubatum*, *Salix spp.*, *Juncus balticus*, *Equisetum spp.* and *Betula occidentalis*. An assessment of variability between occurrences for this community in the Central Parkland is not available from the literature. The distribution and abundance of this community is not clear, and more work is needed to determine its abundance in the province.

This association is similar to the "Populo-Betuletum occidentalis" association as described by Looman (1987a). Similar associates are described for this type but can also include *Populus angustifolia* at low levels. Looman (1987a) suggests that stands of this association seem to be restricted to southern Alberta, where they occur on sandy soils along rivers, and in some low-lying sand dune areas.

Populus balsamifera / *Salix petiolaris*

Balsam poplar / Basket willow

Preliminary Rank: S5

In the Central Parkland, Griffiths *et al.* (1996) describe this community from seven sites on higher ground near Lister Lake, Alberta. These stands are

dominated by balsam poplar of approximately 75 years of age, with a shrub layer of *Salix petiolaris* persisting from an earlier stage of succession. It occupies areas where the water table is at a depth of roughly 1.5 m and soils are Gleyed Eutric Brunisols. This community can contain significant coverages of *Juncus balticus*. Griffiths *et al.* (1996) interpreted this community type as transitional between upland poplar forest and willow shrubbery. Although this type is described from more than one site in the Lister Lake area, an assessment of variability between occurrences is not available. The distribution of this community within the Central Parkland is not well documented, however, this community is considered a common type around lakes and sloughs in the Boreal Forest, Foothills and Parkland Regions of Alberta (G. Griffiths pers comm.).

There are a number of *Populus balsamifera* – *Salix* communities described in Beckingham and Archibald (1996) and Beckingham *et al.* (1996a) but willows are not identified to species making comparisons to these community types tenuous.

Populus balsamifera / *Viburnum opulus* - *Cornus stolonifera* / *Aralia nudicaulis*
Balsam poplar / High-bush cranberry / Red osier dogwood / Wild sarsaparilla
Preliminary Rank: SU

In the Central Parkland, Fehr (1984) describes this community from two sites in the Wainwright and Ribstone areas of Alberta. This community has a consistently well developed *Populus balsamifera* layer. Humus layers are also well developed and soils are Dark Brown or Black Chernozems, which usually show some signs of gleying because of a fluctuating water table (Fehr 1984). Shrub cover is roughly 60% and can include *Cornus stolonifera*, *Populus tremuloides*, *Rosa acicularis* and *Symphoricarpos occidentalis*. Herb-dwarf shrub coverage averages 70% and consists of mesophilic or hydrophilic, shade-tolerant species such as *Aralia nudicaulis*, *Thalictrum venulosum*, *Rubus pubescens* and *Aster ciliolatus*. Balsam Poplar stands are usually located in more mesic locations than Aspen stands, and are commonly located at the bottom of slopes or in depressions surrounding the more hydrophilic willow communities in areas that are moderately well drained (Fehr 1984). An assessment of variability between occurrences for this community is not available from the literature.

Viburnum opulus is not common in Alberta (Moss 1983) and this community type is not well documented in the literature, so more work is required to determine the distribution and abundance of this type.

POPULUS TREMULIODES FOREST/WOODLAND ALLIANCE

Populus tremuloides / (sparse understory)
Trembling aspen Forest

Preliminary Rank: S5

In the Central Parkland, this community is described from one site near the Rumsey Ecological Reserve (Fehr 1982) and from one site near Rumsey, Alberta (Wroe 1971). Looman (1987a) also reports this association from the Grassland and Parkland regions of Alberta, but does not discuss occurrence frequency. Little or no understory growth is typical of these young aspen stands (< 30 years old) where grassland understory species are shaded out, plant debris accumulates, and the understory becomes sparse and species-poor (Looman 1987a). This community can be found in depressions and on gradually-moderate to steep slopes of northerly, sometimes southerly facing aspects (Fehr 1982). It occurs on sites that are submesic to subhydric with moderately to well drained Black Chernozems and Humic Gleysols. Shrub species can be present but are represented by only a few individuals (e.g. *Symphoricarpos occidentalis*, *Rosa acicularis*, *Prunus virginiana*, and *Amelanchier alnifolia*) (Looman 1987a). This is considered a stable, slow-changing community (Fehr 1982) forming incomplete rings around willow-edged sloughs (Wroe 1971) and ranging in size from 0.02-0.05 ha (Looman 1987a, Wroe 1971). This community is differentiated from other aspen communities by its lack of a significant understory component.

Aspen is found province-wide (Moss 1983) and occurrences with a sparse understory are reported from the Central Parkland and Dry Mixedwood Subregions of Alberta (Looman 1987a). Similar stands further north in Alberta tend to have minor *Picea spp.* components of < 2%, and considerably higher herb diversity (Beckingham and Archibald 1996).

Populus tremuloides / *Corylus cornuta* / Low Shrubs and Herbs
Trembling aspen / Beaked Hazelnut / Low Shrubs and Herbs
Preliminary Rank: S5

This community is described as common from several locations in the Central Parkland (Bork 1993, Coupland 1950, Griffiths and Griffiths 1987b, Griffiths *et al.* 1997, Looman 1987a, Moss 1932, Polster and Watson 1979, Timoney and Robinson 1998). It is found on upland, steep riparian, and ravine areas and on mesic, well drained sites with Orthic Gray Luvisolic soils (Timoney and Robinson 1998). It is typically found on north or south-facing slopes and is dominated by a mature aspen canopy with a shrub layer consisting primarily of *Corylus cornuta* at varying heights depending on ungulate browsing pressure (Bork 1993). Given the dense overstory cover, the herb layer tends to be sparse. Associated shrubs and herbs include *Rosa acicularis*, *Amelanchier alnifolia*, *Viburnum edule*, *Rubus idaeus*, *Rubus pubescens*, *Cornus stolonifera*, *Symphoricarpos albus*, *Aralia nudicaulis*, *Cornus canadensis*, *Pyrola asarifolia* and *Galium triflorum*. An assessment of variability between occurrences for this community is not available from the literature.

In the Central Parkland, Griffiths *et al.* (1997) describe this type as a *Populus tremuloides* / *Corylus cornuta* community forming the dominant upland vegetation blanketing most of the higher ground throughout the Blackfoot Recreation Area 26 km east of Edmonton. Timoney and Robinson (1998) describe the same community from this area. Bork (1993) describes a *Populus tremuloides* / *Corylus cornuta* / *Aster conspicuus* - *Disporum trachycarpum* - *Epilobium angustifolium* community and a *Populus tremuloides* / *Corylus cornuta* / *Oryzopsis asperifolia* community from two areas on north facing slopes in Elk Island National Park. From one area on the western edge of the Central Parkland, Griffiths and Griffiths (1987b) describe a *Populus tremuloides* / *Rosa acicularis* - *Corylus cornuta* community. These communities all occur with similar topographic positions and substrate. Canopy and shrub layers are very similar and only show slight variations in herb dominance. They are therefore all considered as the same community in this report. These aspen forests are considered common throughout the Boreal and Parkland Regions of Alberta (Coupland 1950, Griffiths *et al.* 1997, Moss 1932) and Looman (1987a) suggests this association can be found from northern Alberta to Montana and Manitoba.

This community is similar to the *Corylus* - *Rosa* type of Polster and Watson (1979) and the *Populus tremuloides* / *Corylus cornuta* / *Aralia nudicaulis* type of Hardy and Associates (1986), both also from Elk Island National Park. Bork (1993) suggests this community is similar to (unspecified) mature aspen forests found in the Central and Dry Mixedwood Subregions of Alberta. Similarities can also be noted in the "low-bush cranberry Aw" (community d1.3) described by Beckingham and Archibald (1996) from the Boreal Region of Alberta. Moss (1932) called these community types the "aspen consociation", and Looman (1987a) called them the "Alliance *Populion tremuloides* in the association *Corylo-Populetum* (forests dominated with aspen having *Corylus* prominent in the shrub layer)."

Populus tremuloides / *Prunus virginiana* – (Shrub) / Low Herb

Trembling aspen / Choke cherry (shrub) / Low herb

Preliminary Rank: S5

Variations of this community are described from several locations in the Central Parkland by Fehr (1984), Meijer and Karpuk (1999) and Pearson Timberline (1993). *Populus tremuloides* cover averages 41%, and average tree height is 6m (Fehr 1984). Cover of the low shrub layer averages 43%, average cover of the herb-dwarf shrub layer is 58% (Fehr 1984) and bryoid cover is generally low. This community is found on upland eolian and glaciofluvial parent materials on exposed upper slopes where there is no moisture accumulation (Meijer and Karpuk 1999). It is found on dry, sandy outwash plains, dune complexes, interdune depressions, and kame moraines (Pearson Timberline 1993) on the upper portion of dry, north-facing dune

slopes where moisture conditions are subxeric (Fehr 1984). Soils within this community are poorly developed and can be Ortho Regosols, Rego Dark Brown or Ortho Dark Brown Chernozems (Meijer and Karpuk 1999). Drainage is rapid, and nutrient conditions tend to be submesotrophic. Slope can range from 0-70% (Pearson Timberline 1993). Other common species can include *Populus tremuloides*, *Amelanchier alnifolia*, *Symphoricarpos spp.*, *Rosa spp.*, *Juniperus horizontalis*, *Apocynum androsaemifolium*, *Agropyron spp.*, *Thalictrum venulosum*, *Campanula rotundifolia*, *Lathyrus ochroleucus*, *Schizachne purpurascens*, *Carex foenea*, *Thermopsis rhombifolia*, *Galium boreale* and *Smilacina stellata* (Fehr 1984, Meijer and Karpuk 1999, Pearson Timberline 1993). The herb-dwarf shrub layer may contain *Symphoricarpos occidentalis* instead of *S. albus* (Fehr 1984, H. Loonen pers comm.).

Miejer and Karpuk (1999) describe a *Populus tremuloides* / *Populus tremuloides* - *Prunus virginiana* - *Amelanchier alnifolia* / *Juniperus horizontalis* - *Arctostaphylos uva-ursi* community from three areas in Dillberry Lake Provincial Park. Pearson Timberline (1993) describe a *Populus tremuloides* / *Prunus virginiana* - *Amelanchier alnifolia* - *Rosa acicularis* / *Galium boreale* forest type (sometimes lacking one of the shrub components; H. Loonen, pers comm.) from one area in the Wainwright Dunes Ecological Reserve. Fehr (1984) describes a *Populus tremuloides* / *Prunus virginiana* - *Rosa acicularis* community type with a less-developed low shrub layer (10-40% cover), and a common (H. Loonen, pers comm.) but poorly described *Populus tremuloides* / *Rosa acicularis* - *Prunus virginiana* / *Carex siccata* community type from the Wainwright area of Alberta. Fehr (1984) also describes a *Populus tremuloides* / *Prunus virginiana* / *Corylus cornuta* - *Rosa acicularis* / *Symphoricarpos albus* community type from near Wainwright. And finally, Meijer and Karpuk (1999) describe a *Populus tremuloides* / *Amelanchier alnifolia* - *Prunus virginiana* / *Symphoricarpos occidentalis* community as the "buckbrush associated tall shrub community" noted as common in Dillberry Lake Provincial Park. These communities all have consistent dominants, similar associated species and identical site conditions and are as such all considered the same community type in this report.

Beckingham and Archibald (1996) describe a "low-bush cranberry Aw" community type from the Boreal Region of Alberta with a high component of saskatoon and prickly rose. It has similar dominants to the communities described above, but has moister site conditions and non-sandy substrate. Thorpe and Goodwin (1993) describe a *Populus tremuloides* / *Prunus virginiana* - *Amelanchier* / *Galium boreale* community with similar associates and site conditions from the Manito Sand Hills in Saskatchewan.

Populus tremuloides / *Prunus virginiana* / *Solidago canadensis*
Trembling aspen / Choke cherry / Canada goldenrod
Preliminary Rank: S5

This community is described in the Central Parkland from one site near Elk Island National Park (Bork 1993). This community consists of young aspen forests with less developed canopies resulting in a tall forb layer of *Solidago canadensis*, *Aster conspicuous* and *Lathyrus venosus*. *Prunus virginiana* is a key indicator species for this community type. It can be found in lightly browsed areas with variable topography (mainly crests and south-facing slopes) and variable soil conditions (Bork 1993). Further detail on site conditions is not available. The authors have observed this community type in the Lower Foothills, Central and Dry Mixedwood Natural Subregions of Alberta and, although not considered rare, a formal assessment of provincial abundance is not available.

Beckingham *et al.* (1996a) describe a similar community type ("hairy wild rye Aw" c3.3) with variation in shrub composition (*Rosa acicularis* and *Amelanchier alnifolia* present) and herb composition (*Aster conspicuous* and *Lathyrus spp.* absent) from west-central Alberta. Species composition is similar to the *Populus tremuloides* / *Prunus virginiana* – (Shrub) / Low Herb community but site conditions are quite different; soils are not sandy and sites are not well drained which would classify this type as separate based on substrate and hydrological regime (see Grossman *et al.* 1998).

Populus tremuloides / *Rosa acicularis* - *Rubus idaeus*
Trembling aspen / Prickly rose – Wild red raspberry
Preliminary Rank: S5

In the Central Parkland, Meijer and Karpuk (1999) describe this community from two sites in Dillberry Lake Provincial Park. It occurs on north or south-west facing slopes, on Orthic Dark Brown Chernozemic soils in areas with rapid drainage, submesic to xeric moisture regimes, and submesotrophic nutrient conditions (Meijer and Karpuk 1999). Associated shrubs are a minor part of this community but if present can include *Salix bebbiana*, *Rosa woodsii* and *Symphoricarpos occidentalis*. *Lathyrus ochroleucus*, *Disporum trachycarpum* and *Poa spp* can also be present. The authors have encountered this community in the Dry and Central Mixedwood subregions of Alberta. Although not considered rare, a formal assessment of provincial abundance is not available. An assessment of variability between occurrences for this community in the Central Parkland is also not available from the literature.

Beckingham and Archibald (1996) describe two similar communities with common canopy and understory dominants from the Boreal Forest Region of Alberta. However, in both cases there is a conifer tree component (either white spruce or lodgepole pine) and variation in the abundance of *Rosa acicularis* and *Rubus idaeus* (sometimes not present together).

Populus tremuloides / *Symphoricarpos occidentalis* / Herbs
Trembling aspen / Buckbrush / Herbs
Preliminary Rank: S5

In the Central Parkland, this community is described from three sites; two near Dry Island Buffalo Jump Provincial Park (Smith and Kondla 1972, Wroe 1971) and one site near Kinsella, Alberta (Hilton 1970). Wroe (1971) measured tree cover at approximately 90% and a shrub cover of *S. occidentalis* averaging 69%. Smith and Kondla (1972) describe the herb layer as well-developed. It is located on north-facing slopes around depressional areas (Wroe 1971) on "knob and kettle" topography where the water table is relatively high (Smith and Kondla 1972). Soils are Carbonated Regosol Humic Gleysols (near depressions), and Black Chernozems (mature stands) (Wroe 1971). Associated species include *Rosa woodsii*, *Amelanchier alnifolia*, *Aster* spp., *Actaea rubra*, *Aralia nudicaulis*, *Maianthemum canadense*, *Smilacina stellata*, *Galium boreale*, *Calamagrostis inexpansa*, *Vicia americana*, *Fragaria virginiana*, *Epilobium angustifolium*, *Lathyrus* spp., *Carex praticola*, *Viola canadensis* and *Carex lasiocarpa*. An assessment of variability between occurrences for this community is not available from the literature.

Wroe (1971) describes this community as a *Populus tremuloides* / *Symphoricarpos occidentalis* stand from one site near Dry Island Buffalo Jump. Similarly, Smith and Kondla (1972) describe a *Populus tremuloides* / *Symphoricarpos occidentalis* / *Galium boreale* community from one site near the same area having similar site conditions and associated species. Lastly, a *Populus tremuloides* / *Symphoricarpos occidentalis* / *Rubus pubescens* community is briefly described as the "Large Poplar" forest community by Hilton (1970) from near Kinsella, Alberta with identical site conditions and associated species. As such, these are all considered the same community type in this report. This community appears similar to the Poplar Association described by Moss (1932) in central Alberta. We have also observed these forest types in the Central and Dry Mixedwood Natural Subregions of Alberta. Although we do not consider them rare, there is no reliable assessment of their overall abundance.

Populus tremuloides / *Viola adunca* - *Orthilia secunda*
Trembling aspen / Early blue violet – One-sided wintergreen
Preliminary Rank: SU

In the Central Parkland, this community is described by Anderson (1972) from one site near Kinsella, Alberta. He called it the "large poplar community", where the average age of aspen trees is roughly 30 years with a herb community dominated by shade tolerant species. Soil, moisture and nutrient conditions are not discussed and the community description is brief. This community occurs at a lower elevation than the poorly described "small

poplar community” mentioned by Hilton (1970) which contains *Populus tremuloides* but has an understory (structure and dominance not clear) of *Symphoricarpos occidentalis*, *Rosa woodsii*, *Rosa acicularis* and *Rubus strigosus*. There are no other records of this community type in the literature, and no mention of community abundance. Because this type is only described from one site, an assessment of variability between occurrences in the Central Parkland is not available.

POPULUS TREMULOIDES – POPULUS BALSAMIFERA FOREST/WOODLAND ALLIANCE

Populus tremuloides - *Populus balsamifera* - *Betula papyrifera* / *Symphoricarpos occidentalis*

Trembling aspen – Balsam Poplar – Paper birch / Buckbrush

Preliminary Rank: SU

This community is described briefly by Smith and Kondla (1972) from one site near Dry Island Buffalo Jump. The dominant species are mentioned but there is no structural description of the strata within this community type. It is frequent along streams and riverbanks where the substrate is usually sand with little or no soil development. Moisture, nutrient and other site conditions are not discussed. Understory shrubs appear to be poorly developed but can also include *Amelanchier alnifolia*, *Prunus spp.* and *Lonicera involucrata* at trace amounts. Because this type is only described from one site, an assessment of variability between occurrences in the Central Parkland is not available. We have observed communities with similar species composition but different site conditions in the northern Central Parkland, Dry Mixedwood and Central Mixedwood subregions of Alberta, but more work is needed to determine the provincial abundance of this community occurring on a sandy substrate.

Populus tremuloides - *Populus balsamifera* / *Amelanchier alnifolia*

Trembling aspen – Balsam Poplar / Saskatoon

Preliminary Rank: S5

In the Central Parkland, this community is described from one site near Sylvan Lake, Alberta by Griffiths and Griffiths (1987a). This community is an upland forest with dense shrubs, and a highly diverse herb layer without dominants. Moisture, nutrient and other site conditions are not discussed. Associated shrubs include *Viburnum edule* and *Prunus virginiana*. The most prominent herbs include *Maianthemum canadense*, *Viola canadensis* and *Aralia nudicaulis*. Because this type is only described from one site, an assessment of variability between occurrences in the Central Parkland is not available.

This association can be found along the southern fringes of the boreal forest (Beckingham and Archibald 1996), and occurs in many parks and natural

areas in the Edmonton region (Griffiths and Griffiths 1987a). Though not considered rare, the provincial abundance of this community type requires clarification.

Populus tremuloides - *Populus balsamifera* / *Cornus stolonifera* - *Rosa acicularis* - *Viburnum edule* / *Aralia nudicaulis*

Trembling aspen – Balsam poplar / Red osier dogwood – Prickly rose – Lowbush cranberry / Wild sarsaparilla

Preliminary Rank: S5

In the Central Parkland, Pearson Timberline (1993) describes this community from one location in the Wainwright Dunes Ecological Reserve. This forest type is found in wet, eolian depressions with Gleyed, Dark Brown and Dark Grey Chernozemic soils. It occurs in hygric to subhygric areas with imperfect drainage, variable aspect, and slopes ranging from 0-5%. Associated species besides dominants are not discussed. It is also found around lakes and wet areas in the northern Central Parkland, Dry Mixedwood and Central Mixedwood Subregions of Alberta (personal observation). Beckingham and Archibald (1996) describe a similar community (the horsetail Pb-Aw fl.1) which only differs in that it contains components of *Salix spp.*, *Alnus spp.* and *Equisetum spp.* all at covers of <10%. Because this type is only described from one site, an assessment of variability between occurrences is not available.

(Forest/Woodland)

Evergreen

Needle-leaved

Saturated evergreen forest/woodland

PICEA GLAUCA SATURATED FOREST/WOODLAND ALLIANCE

Picea glauca / *Cornus stolonifera* / *Climacium dendroides*

White spruce / Red osier dogwood / Common tree moss

Preliminary Rank: S5

Griffiths *et al.* (1997) describe this community from one site in the Blackfoot Recreation Area of Alberta. It can be found on lower slope positions in depressional sites. Site conditions and associated species are not discussed. This community is classified as a white spruce swamp by Griffiths *et al.* (1997) and can contain *Moneses uniflora* considered locally rare. We have observed this forest type in the Upper Foothills, Dry Mixedwood and Central Mixedwood Natural Subregions of Alberta and consider it a common spruce habitat, however, a formal provincial abundance assessment of this community type is not available. Because this type is only described from one site, an assessment of variability between occurrences is not available.

Similar communities are described by Beckingham and Archibald (1996) from the Boreal Region of Alberta (the dogwood Sw e3.1 community). These boreal communities can contain aspen and balsam poplar as minor canopy components, and have *Rosa acicularis*, *Viburnum edule*, *Linnaea borealis* and *Rubus idaeus* as understory associates.

PICEA MARIANA SATURATED FOREST/WOODLAND ALLIANCE

Picea mariana / *Ledum groenlandicum* / Low Shrubs and Moss

Black spruce / Labrador tea / Low Shrubs and Moss

Preliminary Rank: S5

This community is considered a common black spruce bog and is described from two sites in the Central Parkland (Griffiths and Griffiths 1987b, Griffiths *et al.* 1997). This type typically consists of an open spruce canopy with *Ledum groenlandicum* less than 50cm in height on an extensively moss-covered substrate. It can be found in wetlands or perched depressions on higher ground. Site conditions are not discussed for these parkland communities but are commonly subhydric to hydric. The shrub layer is dominated by *Ledum groenlandicum* but can also include *Vaccinium vitis-idaea*. Herb coverage is often sparse but can include *Rubus chamaemorus*. Moss and lichen cover is extensive and includes *Pleurozium schreberi*, *Cladina rangiferina*, *Hylocomium splendens*, *Ptilium crista-castrensis* and *Sphagnum spp.*

Two occurrences of this community type are described from the Central Parkland with identical site conditions and associated species. Griffiths and Griffiths (1987b) describe a *Picea mariana* / *Ledum groenlandicum* - *Vaccinium vitis-idaea* black spruce bog community from the Bilby Natural Area near Onoway, Alberta. Griffiths *et al.* (1997) describe a *Picea mariana* / *Ledum groenlandicum* / *Pleurozium schreberi* community from the Blackfoot Recreation Area. Given the similarities in topographic position and associated species, these are considered the same community type in this report. An assessment of species variability or constancy between occurrences for this community in the Central Parkland is not available from the literature.

This forest community can be considered common in the northern Central Parkland (Griffiths and Griffiths 1987b, Griffiths *et al.* 1997), and is described in the boreal regions of Saskatchewan (Acton *et al.* 1998) and the conifer forests of Michigan (Marshner 1974). This forest type can be observed throughout the Boreal and Foothills Regions of Alberta (e.g. the “treed bog” communities of Beckingham and Archibald 1996 and Beckingham *et al.* 1996).

PINUS BANKSIANA FOREST/WOODLAND ALLIANCE

Pinus banksiana / *Arctostaphylos uva-ursi*

Jack pine / Bearberry

Preliminary Rank: S5

This typical jack pine community is described from one location in the northern Central Parkland by Griffiths and Griffiths (1987b). Topographic and site conditions are not discussed but this community typically occurs on upland, sandy areas with rapid drainage. Associated species for this Central Parkland type include *Vaccinium myrtilloides*, *Aster laevis*, *Oxytropis deflexa* and *Solidago spathulata*. Although not formally assessed for abundance, this community is not considered rare and can be observed frequently from the north Central Parkland well into the Dry and Central Mixedwood Regions of Alberta.

Beckingham and Archibald (1996) describe several *Pinus banksiana* / *Arctostaphylos uva-ursi* ("bearberry Pj") communities from the Boreal Region of Alberta with similar associates consisting of *Vaccinium myrtilloides* (and *vitis-idaea*), *Linnaea borealis* and *Picea mariana*. They also describe some jack pine stands in this region with minor *Picea glauca* canopy components.

(Forest/Woodland)

**Mixed Evergreen-deciduous
Broad-leaved**

Mixed-evergreen-deciduous cold-deciduous forest/woodland

PICEA GLAUCA MIXED-EVERGREEN DECIDUOUS FOREST/WOODLAND ALLIANCE

Picea glauca - *Populus tremuloides* - *Populus balsamifera* / (shrubs and herbs)

White spruce – Trembling aspen – Balsam poplar

Preliminary Rank: S5

In the Central Parkland, this community is described from one site in Dry Island Buffalo Jump Provincial Park (Smith and Kondla 1972). A detailed description of community structure is not provided, but this type can be found on hillsides and along river valleys. Site conditions and understory composition are not described but the community is included in this report because of the spruce-dominated mixed-deciduous canopy. Total canopy coverage is roughly 12% aspen, 12% balsam poplar and 25% white spruce. This is a well drained, spruce-dominated community; two factors that differentiate it from others within this formation subgroup. This community represents a successional stage towards pure white spruce communities in the

Parkland and Boreal Regions of Alberta. Forests with this cover type are not uncommon in Alberta, but clarification of understory species for this cover type in the Central Parkland is required.

Beckingham and Archibald (1996) describe three similar community types from the Boreal Region of Alberta (the dogwood Sw type e3.4, the dogwood Pb-Sw type e2.3 and the horsetail Pb-Sw type f2.1). These three communities have are spruce-dominated with aspen and balsam poplar canopy components at near equal proportions to the community described above. Understory associates of these boreal types include *Rosa acicularis*, *Linnaea borealis*, *Viburnum edule* and *Aralia nudicaulis*, but it is unclear whether these are included in the Parkland type. Similar spruce-dominated communities, as described by Beckingham *et al.* (1996), can be found in the Foothills Region of Alberta (the low-bush cranberry Sw type e4.2 and the horsetail Pb Sw type i2.1) but tend to have a *Betula papyrifera* or an *Abies spp.* canopy component at trace amounts.

POPULUS TREMULOIDES MIXED-EVERGREEN DECIDUOUS FOREST/WOODLAND ALLIANCE

Populus tremuloides - *Populus balsamifera* - *Picea glauca* / *Rubus idaeus*
Trembling Aspen – Balsam Poplar – White spruce / Wild red raspberry
Preliminary Rank: S5

In the Central Parkland, this community is described from five sites in the Blackfoot Provincial Recreation Area by Timoney and Robinson (1987). Canopy height of this type ranges from 21-28 m in height and tend to have high structural and species diversity, but an assessment of species variability between sites is not provided. This mature and old-growth forest occurs on well to imperfectly drained sites with Orthic Gray Luvisols, or Humic and Luvic Gleysolic soils where conditions are mesic-sub-hygic to hygic. *Cornus stolonifera* is a common understory associate. This community is differentiated from the *Picea glauca* - *Populus tremuloides* - *Populus balsamifera* / (*shrubs and herbs*) described above by its lower spruce canopy component and significantly moister site conditions. A formal assessment of abundance for this community is not available, but we have observed this community in the Boreal and Foothills Regions of Alberta and do not consider it rare.

This association is similar to the “aspen-balsam poplar / rose-raspberry / pink wintergreen”, and “aspen / red-osier dogwood” associations of northern Alberta described by Beckingham (1993). Beckingham and Archibald (1996) describe a similar community type (the dogwood Pb-Aw types e1.1, e1.2 and e1.3) from the Boreal Region of Alberta. Shrub associates of this boreal community can include *Rosa acicularis*, *Cornus stolonifera* and *Viburnum edule*.

Populus tremuloides (Picea glauca) / Aralia nudicaulis / Cornus canadensis
Trembling aspen (White spruce) / Wild sarsaparilla / Bunchberry
Preliminary Rank: S5

In the Central Parkland, this community is described from two sites, both from the Blackfoot Recreation Area (Timoney and Robinson 1998, Griffiths *et al.* 1997). Canopy cover can vary from aspen-dominated to a mix of aspen and white spruce, and some stands can have a significant cover of *Populus balsamifera* and *Betula spp.* This community occurs on mesic, well drained, north-facing slopes, and soils are commonly Orthic Gray Luvisols. Common understory associates can include *Rubus pubescens*, *Rubus idaeus*, *Corylus cornuta*, *Viburnum edule*, *Amelanchier alnifolia*, *Cornus stolonifera* and *Symphoricarpos albus*. The lack of *Populus balsamifera* as a canopy dominant differentiates this community from the *Picea glauca - Populus tremuloides - Populus balsamifera* in this formation. An assessment of variability between occurrences for this community is not available from the literature.

Timoney and Robinson (1998) describe this type as a *Populus tremuloides (Picea glauca) / Aralia nudicaulis / Cornus canadensis* community from south of Elk Island National Park, and Griffiths *et al.* (1997) describe it as a *Picea glauca - Populus tremuloides / low shrub* community from the same area. Both communities have identical site conditions and no variation in associated species, and as such are considered the same community type in this report.

Beckingham and Archibald (1996) describe a similar community type (the low-bush cranberry type d1.2) from the Boreal Region of Alberta, the only difference being a *Rosa acicularis* understory component (<20%).

Herbaceous

Annual

Narrow-leaved

Riparian herbaceous vegetation

SENECIO CONGESTUS HERBACEOUS ALLIANCE

Senecio congestus
Marsh ragwort
Preliminary Rank: S5

In the Central Parkland, this community is described from one site on the foreshore of Beaverhill Lake by Griffiths *et al.* (1996). It is described as the Marsh Ragwort (Riparian) Zone by and can be found occupying land exposed by receding water levels. They describe three subzones of this community; an

inner zone where the ragworts have died after flowering, a zone where the ragwort is in flower, and a zone where the ragwort is colonizing mud exposed during the current season. Soil conditions are not discussed. Associated plant species include *Rumex maritimus*, *Aster brachyactis*, *Ranunculus cymbalaria*, *Chenopodium salinum* and *Hordeum jubatum*. This community can disappear when water levels rise and the area it occupies is flooded. The species *Senecio congestus* is common in the northern Central Parkland Subregion and Boreal Region of Alberta (Moss 1983) and occurrence as a community of this species are not considered rare, although no formal assessments of community abundance are available. Because this type is only described from one site, an assessment of species variability between occurrences is not available. Natureserve (2001) describes a *Senecio congestus* Temporarily Flooded Herbaceous Alliance, but there are currently no details on any associations within that alliance.

(Herbaceous)

Herb

Thread-like leaved

Intermittently flooded brackish herbaceous vegetation

RUPPIA MARITIMA INTERMITTENTLY FLOODED HERBACEOUS ALLIANCE

Ruppia maritima

Widgeon grass

Preliminary Rank: S2

This community is described as a saline open water community occurring generally within the Grassland and Parkland Regions of Alberta by Wallis (1990). It occurs on unvegetated shores, where rooted vegetation is absent, and soil salinity is between 3 and 50 parts per million. It is restricted to saline, seasonally or permanently flooded marshes, channels, ponds, rivers or coastal wetlands or within inland, low-lying basins of high evaporation and low, fresh water inputs. Because this community has very specific water and soil chemistry requirements, its habitat is sensitive to anthropogenic influences. This plant association has been characterized as one of three major foods for waterfowl in the Missouri Coteau (Heidel *et al.* 2000, Metcalf 1931). An assessment of variability between occurrences for this community is not available from the literature.

Wallis (1990) mentions this community for Alberta in the Grassland-Parkland Region of eastern Alberta where it is apparently very localized (i.e. restricted to few water bodies). It may be very extensive in lakes where it occurs and might be more common in Alberta than previous surveys indicate, however, droughts may have severely impacted its habitat (C. Wallis, pers comm.).

This community type occurs in the glaciated plains region of Alberta, Saskatchewan, Montana and North Dakota where it is typically in “closed-basin” watersheds, and often part of glacial outwash channels (Heidel *et al.* 2000). It is briefly described from California and the Lahontan Basin in Nevada (Natureserve 2001) but details on community persistence or variability are not available. Little is known of its current or past distribution other than along the California coast, where it has been much reduced in area by drainage and cultivation.

(Herbaceous)

(Herb)

Toothed, broad leaves

Brackish herbaceous vegetation

POTENTILLA ANSERINA HERBACEOUS ALLIANCE

Potentilla anserina - *Distichlis stricta*

Silverweed – Salt grass

Preliminary Rank: SU

In the Central Parkland, Fehr (1984) describes this community from one area near Wainwright, Alberta. He recorded *Potentilla anserina* at 30% cover and *Distichlis stricta* at 15% cover. Associated species were all at covers of 5% or less. Salt-tolerant species found as associates in this community include *Glaux maritima*, *Hordeum jubatum*, *Puccinellia nuttalliana*, *Salicornia rubra* and *Suaeda calceoliformis*. This type is found in depressions where groundwater discharge makes the soil saline. Other site characteristics are not discussed. Records of this community type were not found elsewhere in the literature so a formal assessment of community variability between occurrences and provincial abundance is not available.

(Herbaceous)

Sedges and Rushes

Spiked Sedges

Marl fen herbaceous vegetation

ELEOCHARIS QUINQUEFOLIA HERBACEOUS ALLIANCE

Eleocharis quinquefolia

Few-flowered spike rush

Preliminary Rank: S3

In the Central Parkland, this community is described from one site in the Bilby Natural Area near Onoway by Griffiths and Griffiths (1987b). It is

described as a Marl Fen or a bog forest associated with seepage of mineral-rich groundwater. Soil and site conditions of this community are not discussed, and a detailed description of the community strata and an assessment of the variability between occurrences of this type in the Central Parkland are not available. Marl fens are considered by naturalists to be important for plant conservation because they tend to contain uncommon, showy plants, such as *Tofieldia glutinosa*, *Parnassia palustris*, *Senecio pauperculus*, *Pedicularis groenlandica* and *Orchis rotundifolia*.

Documentation on the frequency of this species forming communities in Alberta is not available. The species *Eleocharis quinquefolia* is considered to be potentially rare and restricted in range, and is ranked S3 for the province (L. Allen, pers comm.). As a community in the Central Parkland, occurrences are probably not common and, consistent with the species rank, the community type is ranked S3 as well.

In central and northwestern North Dakota, *Eleocharis quinquefolia* communities are found on slopes bordering wetlands and along river valleys (Heidel *et al.* 2000). This plant association is documented from North and South Dakota, and northeastern Montana (Heidel *et al.* 2000) and is similar in site conditions and associated species to the Central Parkland occurrence described above. An upper-subalpine and lower alpine *Eleocharis quinquefolia* wetland community is described from Colorado (Natureserve 2001). Stands are dominated by *Eleocharis quinquefolia* and are easily recognized by their homogeneity (usually little more than *Eleocharis quinquefolia* and *Carex aquatilis*), and the sparse nature of the vegetation growth (these types are ranked G4 by The Nature Conservancy). Because of elevation and hydrological differences (i.e. seasonally flooded), this alpine community could not be considered the same as the Central Parkland type discussed above.

Temperate or subpolar sedge grassland

CAREX SPP. SEDGE GRASSLAND ALLIANCE

Carex group - *Calamovilfa longifolia*

Carex group – Sand grass

Preliminary Rank: S5

In the Central Parkland, this community is described from 15 sites near the Wainwright - Metiskow area of Alberta by H. Loonen (unpublished data). The *Carex* group accounts for approximately 30% cover (range 12-50%) with *Calamovilfa longifolia* at approximately 15% cover (range 7-30%). The *Carex* group consists of *Carex obtusata*, *C. filifolia*, *C. pennsylvanica* and *C. stenophylla* of roughly equal cover. This type is a component in a mosaic with (unspecified) shrub and tree communities. This community occurs on

submesic, rapidly drained areas of various aspects on hummocky, undulating terrain. Soils are Orthic Dark Brown Chernozems. Associated species (both less than 8% cover) include *Stipa curtisetata* and *Koeleria macrantha*. This community frequently makes up 20-70% of the landscape. It is considered common in the Central Parkland Subregion and the Grassland Region of Alberta (H. Loonen pers comm.) but is not documented elsewhere in the literature.

Carex group - *Koeleria macrantha*

Carex group – June grass

Preliminary Rank: S5

In the Central Parkland, this community is described from 40 sites near Wainwright, Alberta by H. Loonen (unpublished data). It is differentiated from the previous community by the dominance of *Koeleria macrantha*. The *Carex* group accounts for approximately 30% cover (range 9-72%) with *K. macrantha* accounting for approximately 8% (range 1-20%). The *Carex* group consists of *Carex obtusata*, *C. filifolia*, *C. pensylvanica* and *C. stenophylla* of roughly equal cover. This type occurs on submesic, rapidly drained sites with Orthic Dark Brown Chernozem soils and hummocky, undulating terrain. Nutrient conditions for this community are described as submesotrophic and it occurs on various aspects. Associated species (less than 5% cover) include *Bouteloua gracilis*, *Calamovilfa longifolia*, *Stipa curtisetata*, *Cerastium arvense* and *Juniperus horizontalis*. In these areas the grassland community makes up approximately 20-70% of the landscape. This community relies on the presence of grazing to exist. It is considered common throughout Central Parkland Natural Subregion and the Grassland Region of Alberta (H. Loonen pers comm.) but is not documented elsewhere in the literature.

Carex group - *Stipa curtisetata*

Carex group - Barkworth

Preliminary Rank: S5

In the Central Parkland, this grassland community type is described from 27 sites near the Ribstone and Czar areas of Alberta by H. Loonen (unpublished data). This community is differentiated from the previous two communities by the dominance of *Stipa curtisetata*. The *Carex* group accounts for approximately 25-50% cover in ungrazed areas, with *Stipa curtisetata* accounting for approximately 5-25% cover. The *Carex* group consists of *Carex obtusata*, *C. filifolia*, *C. pensylvanica*, and *C. stenophylla* of roughly equal cover. Associated species (less than 16%) include *Festuca hallii* and *Calamovilfa longifolia*. This type can be found on rapidly drained, mesic, gentle slopes with various aspects and submesotrophic conditions. Soils are Ortho Dark Brown Chernozems. Where this community is found the grassland component can cover 20-70% of the landscape. This community is considered common throughout Central Parkland Natural Subregion and the

Grassland Region of Alberta (H. Loonen pers comm.) but it is not documented elsewhere in the literature.

Temperate or subpolar intermittently flooded sedges and rushes

CAREX ATHERODES INTERMITTENTLY FLOODED HERBACEOUS ALLIANCE

Carex atherodes - *Carex rostrata*
Awned sedge – Beaked sedge
Preliminary Rank: S2

In the Central Parkland, this community is described from one location near Lister Lake, Alberta by Griffiths *et al.* (1996). *Carex atherodes* and *Carex rostrata* show varying degrees of dominance (not specified) amongst a diversity of associated species at unspecified coverages. Associated species include *Drepanocladus aduncus*, *Calamagrostis stricta*, *Sium suave*, *Epilobium palustre*, *Galium trifidum* and *Rumex occidentalis*. This community can be found in areas where the water table is near ground level. Soil and site conditions are not discussed. This type is characteristic of eutrophic sites subject to flooding. *Carex rostrata* is considered a rare species in Alberta and is ranked S2 for the province (ANHIC 2001); the rank of this community type reflects this. This community type is not documented elsewhere in the literature. Natureserve (2001) notes *Carex atherodes* occurring with several associated species including *Polygonum amphibium*, *Calamagrostis canadensis*, *Carex lasiocarpa*, *Spartina pectinata*, and *Carex aquatilis* but describe no such community comparable to this Central Parkland type.

Temperate or subpolar permanently-flooded sedges and rushes

CAREX SPP. PERMANENTLY FLOODED HERBACEOUS ALLIANCE

Carex atherodes - *Scolochloa festucacea* - *Eleocharis palustris* - *Typha latifolia*
Awned sedge – Spangeltop – Common spike rush – Common cattail
Preliminary Rank: S5

In the Central Parkland, this community is described from one site near Rumsey, Alberta by (Fehr 1982). It occupies the subhydric and hydric margins of relatively deep freshwater sloughs. Drainage is very poor with areas of open water being common. Cover of the herb-dwarf shrub layer is roughly 90%. Other characteristic species can include *Alopecurus aequalis*, *Rumex maritimus*, *Glyceria grandis*, and *Beckmannia syzigachne*. This community is differentiated from the *Carex atherodes* Intermittently flooded Herbaceous Alliance by hydrological regime (i.e. this community is permanently flooded). The perimeters of many sloughs in the Central

Parkland are vegetated with this community type (Fehr 1982). An assessment of variability between occurrences for this community is not available from the literature.

Natureserve (2001) notes a *Scolochloa festucacea* Herbaceous Vegetation association where *Scolochloa festucacea* occurs with low covers of *Carex atherodes*. This marsh type is found in the northeastern Great Plains and northern tallgrass prairie region of the United States and Canada but cannot be considered similar as it lacks *Eleocharis palustris* and *Typha latifolia* components and is seasonally, not permanently, flooded.

Carex pseudo-cyperus - *Calla palustris*
Cyprus sedge - Water arum
Preliminary Rank: S1S2

In the Central Parkland, Moss (1928) only found this association in two areas; forming a narrow fringe around a water body backed by muskeg of young bog forest, and as wetland adjoining two lakes. According to Allen (2001), this community can form islets in ponds where the water is shallow enough, and it is often associated with *Carex diandra*. Specific site and soil conditions are not documented, nor are assessments of variability between occurrences. Well-developed occurrences of this community are known only from three sites near Elk Island National Park (G. Griffiths pers comm.), therefore, the community has been assigned a rank of S1S2. This community type is not well described in the literature and is on the Alberta Plant Community Tracking List (Allen 2001).

Temperate or subpolar seasonally flooded sedges and rushes

CAREX ATHERODES SEASONALLY-FLOODED HERBACEOUS ALLIANCE

Carex atherodes - *Calamagrostis inexpansa*
Awned sedge - Northern reed grass
Preliminary Rank: S4

In the Central Parkland, this community is described by Wroe (1971) from one area near Dry Island Buffalo Jump Provincial Park. He noted that the *Carex spp.* cover is roughly 90% and *Calamagrostis inexpansa* cover is approximately 61%. *Beckmannia syzigachne* cover is approximately 10%. Coverage of plants reflect the depth of water and the length of flooding, in deeper water *Glyceria grandis*, *Typha latifolia*, and *Sium suave* can be found. This community type is associated with rolling topography; it occurs in depressions that hold water during part of the growing season. Site and soil conditions are not discussed and an assessment of species variability between occurrences is not available. Associated species include *Carex rostrata* and *Carex athrostachya*. This community is differentiated from other *Carex*

atherodes Herbaceous Alliances by hydrological regime (i.e. it is not permanently flooded). This is considered a common community type, but it is restricted in the area that it can occupy (A.H. Weerstra pers comm.) and is hence ranked S4. It is not well documented in the literature.

Carex atherodes - *Scolochloa festucacea* - *Poa palustris* - *Cirsium arvense*
Awned sedge – Spangletop – Fowl bluegrass – Canada thistle
Preliminary Rank: SU

In the Central Parkland, this community is described from one area near Wainwright, Alberta by Fehr (1982). Coverages of the dominant species were estimated as follows: *Carex atherodes* coverage is 35%, *Scolochloa festucacea* is 30%, *Poa palustris* is 20% and *Cirsium arvense* is 19%. Fehr (1982) estimated the herb-dwarf shrub layer at between 85 and 95 % cover. Associated species include *Potentilla anserina*, *Mentha arvensis*, and *Hordeum jubatum* all at coverages of less than 8%. This community type is found in hygric and subhydryc depressions. It occurs in very poorly to imperfectly drained areas with Rego Humic Gleysolic soils, typically in sloughs that dry up in summer. This community type is prevalent in the kettle bottoms and valleys of the Central Parkland where there is little open standing water. The provincial abundance of this community type is not clear and it is not well documented in the literature. An assessment of variability between occurrences for this community is not available from the literature.

Natureserve (2001) notes a *Scolochloa festucacea* Herbaceous Vegetation association where *Scolochloa festucacea* occurs with low covers of *Carex atherodes*. This marsh type is found in the northeastern Great Plains and northern tallgrass prairie region of the United States and Canada and is seasonally flooded. It lacks the *Poa palustris* and *Cirsium arvense* components of the Central Parkland community, but these are introduced species; pre-introduction of these non-natives, this Parkland community might have been very similar to the community described by Natureserve (2001).

ELEOCHARIS PALUSTRIS SEASONALLY FLOODED HERBACEOUS ALLIANCE

Eleocharis palustris
Common spike rush
Preliminary Rank: S5

Wallis (1990) describes this community as occurring in moderately saline emergent marshes of the Central Parkland Natural Subregion of Alberta. The structure of this community is not provided in the literature. Associated species for Alberta types are not documented, nor is an assessment of species variability between occurrences. The common spike-rush plant community occurs in slightly to moderately brackish wetlands, including prairie potholes

and riparian settings that are seasonally flooded (Heidel *et al.* 2000). The species *Eleocharis palustris* is widely distributed throughout Alberta (Moss 1983) but its occurrence as a community type in the province is not well documented.

This community type is well documented from the central and northern Great Plains, possibly extending into the Southwest and the Pacific Northwest states, and is a major type at low elevations throughout Montana (Heidel 2000, Hansen *et al.* 1995). This plant association is characterized by short-statured, fine-textured vegetation in localized areas where mineral-rich ground water flow emerges at the ground's surface. Associated species for Montana communities include *Eleocharis acicularis*, *Rumex crispus* and *Hordeum jubatum*, but an assessment of community variability between occurrences is not available.

Temperate or subpolar temporarily flooded sedges and rushes

CAREX ATHERODES TEMPORARILY FLOODED HERBACEOUS ALLIANCE

Carex atherodes / *Lemna minor*

Awed sedge – Common duckweed

Preliminary Rank: S5

In the Central Parkland, this community type is described by Griffiths and Griffiths (1987b) from one site near Onoway, Alberta. This type is described as “beds of awed sedge occurring in morainal depressions, either in moist soil as monocultures of *Carex atherodes* or in shallow water in association with *Lemna minor*” (Griffiths and Griffiths 1987b). Site conditions, soil type and associated species are not discussed. This community is considered common on intermittently flooded ground in Alberta (G. Griffiths, pers comm.) but is not well documented in the literature. An assessment of structural variability between occurrences is not available.

A similar community type is described by Schneider *et al.* (1997). This awed sedge wet meadow occurs in the northern tallgrass prairie region of the United States and Canada. Stands occur on lowland sites, typically in depressions, basins or along streams and rivers, having fresh or moderately saline standing water for several weeks each year. Associated species and site conditions are not discussed for this prairie type.

Temperate or subpolar semipermanently flooded sedges and rushes

CAREX AQUATILIS SEMIPERMANENTLY FLOODED HERBACEOUS ALLIANCE

Carex aquatilis - *Carex utriculata*

Water sedge – Greater beaked sedge

Preliminary Rank: S4

In the Central Parkland, this community is described by Griffiths *et al.* (1997) from one site in the Blackfoot Recreation Area in Alberta. This association is recognized by the presence of both *Carex aquatilis* and *Carex utriculata* in roughly equal proportions, but associated species and soil conditions are not discussed. This community type is a common fen that occurs in wet, morainal depressions and along the shores of lakes and ponds representing the interface between water sedge fen and aquatic, open water vegetation. This is considered a common fen type throughout Alberta (Griffiths *et al.* 1997) but is not well documented provincially in the literature. An assessment of structural variability between occurrences is not available.

This community has similar dominants and moisture regimes (i.e. poorly drained, moist areas) to the *Carex aquatilis* type mentioned from the USA Great Plains by Schneider *et al.* (1997) and to the *Carex rostrata* – *Carex aquatilis* type described from the Manito Sandhills in Saskatchewan by Thorpe and Goodwin (1993). Associated species and coverages are not provided for comparison for the community mentioned by Schneider *et al.* (1997). However, associated species in Saskatchewan types include *Typha latifolia*, *Calamagrostis inexpansa* and *Mentha arvensis* at unspecified coverages. In Colorado, the *Carex aquatilis* - *Carex utriculata* community type is considered common and generally occurs in small to moderate-sized patches in very shallow, slow-moving to still water, or on saturated soils near low-order streams, lakes, and backwater areas of larger rivers (Bourgeron and Engelking 1994). This type is ranked G4 by The Nature Conservancy (Natureserve 2001).

SCIRPUS PALUDOSIS PERMANENTLY FLOODED HERBACEOUS ALLIANCE

Scirpus paludosus

Prairie bulrush

Preliminary Rank: S4

In the Central Parkland, this community is described by Fehr (1982) from one area near Wainwright, Alberta and from the Parkland and Grassland Regions of Alberta in saline marshes by Wallis (1990). It occurs as a wide band around saline sloughs. It is described by Wallis (1990) as a saline, emergent marsh dominated by prairie bulrush with water contributing roughly 20% cover (range 5-35%). Associated species and soil types are not discussed. This community is considered common in the parkland region (C. Wallis pers comm.) as is the species *Scirpus paludosus*, but a formal assessment of its occurrence as a community is not available. Saline marshes are not uncommon in the Parkland or Grassland Regions, but would be the limiting factor for area extent of this community type. Natureserve (2001) describes the species *Scirpus paludosus* as SR for Alberta (SR = reported from) but lists

no communities dominated by this species. This community was not found elsewhere in the literature.

SCIRPUS SPP. SEMIPERMANENTLY FLOODED HERBACEOUS ALLIANCE

Scirpus acutus

Great bulrush

Preliminary Rank: S5

In the Central Parkland, this community is described from two sites in Dillberry Provincial Park (Meijer and Karpuk 1999), one site in the Blackfoot Provincial Recreation Area (Griffiths *et al.* 1997), and generally from emergent and saline marshes in the Grassland and Parkland Regions of Alberta (Wallis 1990). This community occurs along lakeshores in poorly drained areas with Typic and Terric Humisolic soils. Moisture conditions tend to be subhydric with a hypereutrophic or eutrophic nutrient regime (Meijer and Karpuk 1999). Associated species include *Cirsium arvense*, *Rumex occidentalis* and *Solidago canadensis*.

Griffiths *et al.* (1997) describe a *Scirpus acutus* - *Scirpus validus* community occurring in narrow bands along the shores of lakes or ponds in the Blackfoot Recreation Area. Wallis (1990) describes this same community as occurring in moderately saline emergent marshes in the Parkland and Grassland regions of Alberta. Published descriptions of this community do not allow for structural comparisons or comparisons of species coverages between occurrences.

Elsewhere, this bulrush community is noted as among the most common plant associations of semi permanently flooded wetlands in parts of northeastern Montana (Thompson 1994) and at low elevations throughout Montana (Hansen *et al.* 1995, Heidel *et al.* 2000). *Scirpus acutus* is considered common in Alberta and the USA both as a plant species and a dominant community type (Moss 1983, Heidel *et al.* 2000, Hansen *et al.* 1995, Thompson 1994), but documented occurrences are not of sufficient detail to afford comparisons of coverage values and structural diversity.

Scirpus paludosus - *Puccinellia nuttalliana* - *Hordeum jubatum* - *Distichlis stricta*

Prairie bulrush – Nuttall's salt meadow grass – Foxtail barley – Salt grass

Preliminary Rank: SU

In the Central Parkland, this community type is described by Fehr (1982) from three sites near Rumsey, Alberta. This community type occupies subhydric saline depressions where water contributes on average 20% cover (Fehr 1982). It occurs in wide bands around saline sloughs where each of the dominant species occupy a particular location along a moisture gradient. From wettest to driest in this transitional zone the order of species occupancy

is *Scirpus paludosus*, *Puccinellia nuttalliana*, *Hordeum jubatum*, and *Distichlis stricta*. Associated species and soil types are not discussed. Dodd and Coupland (1966) note that because each species in this community can dominate a transitional zone of moisture and salinity, each species could also be considered as its own community type. All dominant species are widespread in Alberta (Moss 1983) but the occurrence of this community type is not well documented provincially or elsewhere. As such, an assessment of structural variability between occurrences is not available.

Scirpus acutus – Liverworts

Great bulrush - Liverworts

Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from one site in Dillberry Lake Provincial Park. It occurs in a poorly drained lacustrine deposit with Rego Gleysolic soils, a subhydric moisture regime and hypereutrophic nutrient conditions. The species composition of the liverwort group was not defined. Associated species include *Solidago canadensis*, *Rumex occidentalis*, *Cirsium arvense* and *Scirpus pungens*. This community could be similar to the *Scirpus acutus* community described above, but further work is required to determine associated species of that community before appropriate comparisons can be made with this type. This community is not well documented in the literature so the abundance of this type is not clear and an assessment of structural variability between occurrences is not available.

Scirpus pungens

Three-square rush

Preliminary Rank: S4

This community is noted by Wallis (1990) as occurring generally in emergent and saline marshes of the Parkland and Grassland Regions of Alberta. It is described as a saline emergent marsh dominated by three-square rush (Wallis 1990). It occurs in depressional areas and around the perimeters of saline marshes in Alberta. Soil and nutrient conditions are not discussed. Little information is available for this community, but it is possible that it is similar to, or part of, the *Scirpus pungens* - *Hordeum jubatum* - *Puccinellia nuttalliana* community described below. Associated species and specific site conditions are not clear from the literature. The species *Scirpus pungens* is well documented across Alberta and Canada (Moss 1983, C. Wallis, pers comm.) but more work is required to assess the frequency of its occurrence as a community. As such, an assessment of structural variability between occurrences is not available.

Scirpus pungens - *Hordeum jubatum* - *Puccinellia nuttalliana*

Three square rush – Foxtail barley – Nuttall's salt meadow grass

Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from three sites in Dillberry Lake Provincial Park. It is described as a shallow marsh with the three dominant species occurring at unspecified coverages. It occurs in poorly drained, saline, shallow marshes. This type is characterized by very rich (hypereutrophic) nutrient conditions, and Rego Gleysolic or Gleyed Rego gleysolic soils. Associated species include *Ranunculus cymbalaria*, *Suaeda calceoliformis* and *Puccinellia nuttalliana*. The three component species of this community are common in Alberta (Moss 1983) but the occurrence of these species together as a community type is not well documented. This community is not described elsewhere in the literature and, therefore, an assessment of structural variability between occurrences is not possible.

Temperate or subpolar sedges or rushes on sandy substrate

CAREX SPP. RAPIDLY DRAINED HERBACEOUS ALLIANCE

Carex siccata - *Calamovilfa longifolia*

Hay sedge – Sand grass

Preliminary Rank: S3

In the Central Parkland, this community is described from one site by Fehr (1984) near the Wainwright area of Alberta. This community occurs in active blowouts on south or south-west-facing slopes on undulating areas where there is little relief. Soils are Orthic Regosols. It occurs on xeric to very xeric sites with rapid drainage. These sites are characterized as sandy, possessing little substrate. Associated species are not discussed. This community is an important colonizer of bare sand areas in the Central Parkland of Alberta. Because this community type is restricted to active blowouts, it has a limited range (Fehr 1984). This community type is not well documented in Alberta and an assessment of structural or species variability between occurrences is not possible.

Natureserve (2001) describes a *Carex siccata* Herbaceous Alliance, however, this alliance occurs in dry to moderately moist, open or thinly wooded, grassy slopes in the Rocky Mountains. Elevations range from 2400 m in Arizona to 3900 m in Colorado. These types are typically found near treeline on south-oriented, stabilized, fine-talus slopes, and are as such not comparable to Central Parkland types that differ in substrate, local disturbance (i.e. active blowouts, blowing sand, etc.) and elevation.

(Herbaceous)

(Sedges and Rushes)

Aquatic Rushes

Temperate or subpolar semipermanently flooded aquatic rushes

TYPHA LATIFOLIA SEMIPERMANENTLY FLOODED HERBACEOUS ALLIANCE

Typha latifolia

Common cattail

Preliminary Rank: S5

In the Central Parkland, this community type is described from two sites in Dillberry Lake Provincial Park by Meijer and Karpuk (1999), from one site near Lister Lake by Griffiths *et al.* (1996) and from one site in the Blackfoot Provincial Recreation Area by Griffiths *et al.* (1997). This community is dominated by *Typha latifolia* and can have significant coverages of floating duckweed (*Lemna trisulca*) indicative of areas containing standing, eutrophic water (Griffiths *et al.* 1995). This marsh community is found in level and depressional areas along the shorelines of water bodies, and is extensive in ditches and in riparian zones. It is found in areas where the water level is above the rooting zone for at least part of the growing season, slope and aspect are level, and nutrient levels are high. In the Central Parkland, this community occurs in very poorly drained areas with Rego Gleysol or Rego Humic Gleysolic soils. Formal assessments of structure and variability between occurrences are not available for this community.

We have observed this community around ponds, lakes, sloughs and flooded areas throughout the province. Beckingham and Archibald (1996) describe this community type from the Boreal Region of Alberta (cattail marsh type h1.1), and Beckingham *et al.* (1996) describe it from the Foothills Region of Alberta (cattail marsh type n1.1). It is also mentioned from throughout the Northern Great Plains of Canada and the USA by Heidel *et al.* (2000) in semi-permanent wetlands, ponds and drainages where, under more alkaline conditions, *Typha angustifolia* can begin to dominate.

(Herbaceous)

Graminiod

Linear, parallel-veined leaves

Temperate or subpolar grasslands

AGROPYRON TRACHYCAULUM HERBACEOUS ALLIANCE

Agropyron trachycaulum - *Hordeum jubatum* - *Elymus canadense*

Slender wheat grass – Foxtail barley – Canada wild rye

Preliminary Rank: SU

In the Central Parkland, this community is described by Bradley and Bradley (1977) from one site on the Neutral Hills near Provost, Alberta. It occurs on south-facing grassland slopes (10% grade) with little or no shrub cover. We have observed this community occurring on upper-slope positions between thickets of silverberry growing in lower-slope positions, and young, stunted aspen growing on the tops and north face of the Neutral Hills. This community is generally low-growing due to browsing pressure of deer and elk in the area, and patches of exposed soil accounting for <5% cover are also characteristic of this community. Associated species are not discussed but *Artemisia longifolia* can be a common. This community is made up of species that are widespread in the Central Parkland (Moss 1983), however, a formal assessment of abundance or structural variability between occurrences is not available for this community. The *Agropyron* communities of Alberta are discussed in detail by Vujnovic and Bentz (2001), but this community type is not mentioned therein.

AGROSTIS SCABRA HERBACEOUS ALLIANCE

Agrostis scabra - *Achillea millefolium* - *Antennaria parvifolia*
Hair grass – Common yarrow – Small-leaved everlasting
Preliminary Rank: SU

In the Central Parkland, this community is described by Fehr (1982) from two sites near Rumsey, Alberta. *Agrostis scabra* is the dominant species with an average cover of 23%. The two subdominant species *Achillea millefolium* and *Antennaria parvifolia* both have covers of 15%. This community occurs in shallow, subxeric to submesic depressions on moraine plateau hilltops. Soils are moderately-well to imperfectly drained Dark Brown Chernozems with variable moisture regimes. The moisture regime between stands can vary considerably (but not specified exactly how). Common associates are *Bouteloua gracilis* (2.5% cover), *Koeleria cristata* (2.5% cover) and *Galium boreale* (2.5% cover). All dominant species that compose this community are widespread, but the occurrence of these species together as a community type is not well documented. The overall abundance of this community is unknown and a range of cover values between occurrences (albeit n=2) is not available.

BOUTELOUA GRACILIS HERBACEOUS ALLIANCE

Bouteloua gracilis - *Artemisia frigida*
Blue grama – Pasture sagewort
Preliminary Rank: S4S5

In the Central Parkland, this community is described by Fehr (1982) from six sites near Rumsey, Alberta. It is described as occupying xeric and subxeric sites on the upper and middle sections of well drained, south-facing slopes of

20-35 degrees. Soils are Dark Brown Chernozems. Coverage of the dominant species, *Bouteloua gracilis*, is 35% and *Artemisia frigida* has a coverage of 17%. Associated species at coverages <12% include *Koeleria cristata*, *Stipa curtiseta*, *Antennaria nitida*, *Heterotheca villosa*, *Comandra pallida*, *Haplopappus spinulosus*, *Lygodesmia juncea* and *Thermopsis rhombifolia*. Although *Bouteloua gracilis* is a common grass species, *Bouteloua* – *Artemisia* communities are not represented elsewhere in the literature. A.H. Weerstra (pers comm.) sampled several *Bouteloua* community types throughout the Central Parkland of Alberta and considers this a common community type in this subregion. This community is composed of provincially common species (Moss 1983) but their occurrence together as a community type has not been formally assessed, and a comparison of structural variability between occurrences is not possible.

CALAMAGROSTIS SPP. HERBACEOUS ALLIANCE

Calamagrostis canadensis

Marsh reed grass

Preliminary Rank: S5

In the Central Parkland, this community type is documented from the Blackfoot Recreation Area (Timoney and Robinson 1997) and from near Onoway, Alberta (Griffiths and Griffiths 1987b). This community is typically a patchy mosaic of *Calamagrostis canadensis*. It occurs on imperfectly to poorly drained sites bordering water bodies or in seasonally-inundated depressions. Plants commonly associated with marsh reed grass include *Poa pratensis*, *Epilobium angustifolium*, *Carex spp.*, *Salix spp.*, *Pinus banksiana*, *Picea engelmannii*, and *Populus tremuloides*. Although only formally documented in a few locations from the Central Parkland, we have observed this community type province-wide in similar site conditions. Information is not available for a comparison of structural variability between occurrences.

Calamagrostis canadensis can be found in moist woods, meadows, wetlands, clearings and lakeshores throughout Alberta (Johnson *et al.* 1995) where it often forms dominant communities. Marsh reed grass grows in pine, spruce, and mixed wood forests north of 35° latitude from Newfoundland to Alaska (Lieffers and Macdonald 1993), and is the most common and widespread *Calamagrostis* species in North America (USDA 1988). In the United States, marsh reed grass is documented as a water loving plant commonly growing on imperfectly to moderately well-drained sites; soils are often peat-based, and have a pH as low as 3.5 (USDA 1988). Schneider *et al.* (1997) describe this community type as occurring in the grasslands of Montana.

Calamagrostis stricta - *Juncus balticus* - *Potentilla anserina* - *Aster hesperius*

Narrow reed grass – Wire rush – Silverweed – Western willow aster

Preliminary Rank: S4

In the Central Parkland, this community is described by Griffiths *et al.* (1996) from one area near Lister Lake, Alberta. It is described as a moist, saline meadow and represents the transition between moist meadow and willow shrubbery. Species diversity averages 10-25 species per 10x10 m plot and can include *Glaux maritima*, *Primula incana* and *Aster falcatus*. Soils, site conditions and species coverages are not discussed. This community is considered common on shores in the Central Parkland but it is possible it has been converted to “improved” pasture in some areas, hence was given an S4 rank (G. Griffiths, pers comm.). This community is not documented elsewhere in the literature, so an assessment of structural variability between occurrences is not available.

DISTICHLIS STRICTA HERBACEOUS ALLIANCE

Distichlis stricta

Salt grass

Preliminary Rank: S4

In the Central Parkland, this community is documented from one site in the Rumsey area by Fehr (1982), and occurs generally as a hypersaline dry meadow in the Parkland and Grassland Regions of Alberta (Wallis 1990). This community occurs as a wide band of vegetation around saline sloughs, occupying areas which are less saline than areas occupied by *Puccinellia nuttalliana*, forming almost pure stands with a basal cover of > 20% (Wallis 1990). *Glaux maritima* can be found frequently in this community type. *Distichlis stricta* is ranked S4 in Alberta (ANHIC 2001). Its occurrence as a provincial community type is not well represented in the literature. As such, an assessment of structural variability between occurrences is not available.

Salt grass is distributed across western North America and is generally restricted to low-lying alkaline or saline areas. It is a dominant species on saline areas occurring on a variety of soil textures and moisture levels as far south as Oklahoma (Dodd and Coupland 1966) and can be found as far north as Wood Buffalo National Park (Schwartz and Wein 1997). In Colorado and Wyoming, salt grass covers an estimated 500,000 ha (Mueller *et al.* 1985). It occupies much less extensive areas in the warm desert region south of Utah, but it is prominent in the most saline parts of desert meadows (Dodd and Coupland 1966).

Natureserve (2001) describes this community in semi-arid and arid western North America, from southern Saskatchewan to Mexico. Stands are found in lowland habitats such as playas, swales and terraces along washes that are typically intermittently flooded. The flooding is usually the result of highly localized thunderstorms. This association may also occur in other flood regimes (temporarily, seasonally, and semipermanently). Soil texture ranges

from clay loam to sandy clay. Vegetation cover is sparse to dense and is dominated by *Distichlis stricta* occurring in nearly pure stands. Minor cover of associated graminoids may include *Muhlenbergia asperifolia*, *Hordeum jubatum*, *Agropyron smithii*, *Sporobolus airoides*, *Carex filifolia*, *Eleocharis palustris*, *Puccinellia nuttalliana*, and *Juncus balticus*. Associated forbs, such as *Iva axillaris*, *Helianthus* spp., *Aster* spp. (from lower salinity sites), *Salicornia rubra*, *Triglochin maritima*, and *Suaeda* spp., may also be present. Shrubs are rare, but scattered *Atriplex canescens* and *Sarcobatus vermiculatus* may be present.

Distichlis stricta - Agropyron smithii

Salt Grass – Western wheat grass

Preliminary Rank: S4

This community type is noted by Wallis (1990) as a saline dry meadow type occurring generally within the Parkland and Grassland Regions of Alberta. This community is transitional between the highest, driest portion of the halophytic communities, and the semi-halophytic communities found further upslope. Soil types are not discussed. Common associates include *Aster ericoides* and *Grindelia squarrosa*. The inclusion of *Agropyron smithii* suggests drier site conditions and is thus differentiated from the *Distichlis stricta* community type. *Distichlis stricta - Agropyron smithii* is less abundant than the *Distichlis stricta* community type, but is still widespread in the Central Parkland Natural Subregion and Grassland Region of Alberta (C. Wallis, pers comm.). This community type is not well represented in the literature, and we found no instances where cover values were recorded allowing for a comparison of species variability between occurrences of this type.

ELYMUS CANADENSIS HERBACEOUS ALLIANCE

Elymus canadensis - Poa pratensis - Glycyrrhiza lepidota

Canada wild rye – Kentucky bluegrass – Wild licorice

Preliminary Rank: S4

In the Central Parkland, this community type is described by Griffiths *et al.* (1997) from one site in the Blackfoot Provincial Recreation Area. Coverages of the dominant species are not discussed, nor are soil types. This dry grassland community is found on south-facing, steep slopes (river valleys or steep hills) where shrubs are relatively sparse and, when present, consist mainly of *Symphoricarpos occidentalis*. Shrubs are overtopped by grasses and wild licorice. This community is not considered uncommon but it is found on steep south-facing slopes potentially limiting its overall distribution (G. Griffiths pers comm.). Because this community is only described from one site a comparison of species and structural variability between occurrences is not possible.

KOELERIA MACRANTHA HERBACEOUS ALLIANCE

Koeleria macrantha - *Agropyron smithii*

June grass – Western wheat grass

Preliminary Rank: SU

In the Central Parkland, this community is described by Wroe (1971) from one area near Dry Island Buffalo Jump, and by Fehr (1982) from one area near Wainwright, Alberta. This grassland community type occupies xeric sites on level areas and lower slopes with Black and Brown Solonetz soils. Coverages of the dominant and associated species are not provided, and moisture and nutrient conditions are not discussed. Common associates include *Selaginella densa*, *Carex stenophylla*, *Poa secunda*, *Grindelia squarrosa* and *Distichlis stricta*. It is considered a minor community type in the Rumsey area (Fehr 1982) but is not well documented in the literature, so the provincial abundance and structural variability between occurrences are not clear.

A similar community type is noted from the northern and western Great Plains, Rocky Mountains, and the interior western United States (Natureserve 2001). This community type is described as a *Agropyron smithii* Herbaceous Vegetation with *Koeleria macrantha* achieving local dominance (at unspecified coverages). It can be found on level to gently sloping terrain, on alluvial fans, swales, river terraces, floodplains, valley floors and basins. The soils are clay, clay loam, and silt loam. *Agropyron smithii* can dominate the moderate to dense (40-100% cover) herbaceous canopy that grows 0.5-1 m tall. Other graminoids that co-occur and may achieve local dominance are *Eleocharis palustris*, and *Poa* spp. Other associated species can include *Artemisia ludoviciana*, *Eriogonum* spp., *Bouteloua gracilis*, *Nassella viridula*, and *Stipa comata*. Shrubs and dwarf-shrubs are rare in this community. Natureserve (2001) refers the reader to Thilenius *et al.* (1995), Baker (1984a), and Baker and Kennedy (1985).

STIPA SPP. HERBACEOUS ALLIANCE

Stipa comata - *Koeleria macrantha* - *Agropyron smithii*

Needle and thread grass – June grass – Western wheat grass

Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from three sites in Dillberry Lake Provincial Park. A structural description of this community is not provided, but it can be found on stabilized dune areas with little sign of recent erosion and visible evidence of soil formation. It occurs on south to south-west facing slopes with Orthic Dark Brown soils, rapid drainage, a subxeric moisture regime, and

submesotrophic nutrient conditions. Forbs and grasses that can be present include *Solidago missouriensis*, *Thermopsis rhombifolia*, *Calamovilfa longifolia* and *Agropyron trachycaulum*. The three dominant species are provincially common but their occurrence together as a community type is not clear. The provincial abundance of this type is unknown and it is not documented elsewhere in the literature.

Stipa comata - *Koeleria macrantha* - *Agropyron trachycaulum* - *Agropyron smithii* - *Carex obtusata*

Needle and thread grass – June grass – Slender wheat grass – Western wheat grass – Blunt sedge

Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from five sites in the north section of Dillberry Lake Provincial Park. It occurs on south slopes and flat, upland areas associated with hummocky, glaciofluvial landscape. It is found in xeric, rapidly drained areas with Orthic Dark Brown soils, and submesotrophic nutrient conditions. Coverages of dominants are not provided, but common associates include *Artemisia frigida*, *Petalostemon purpureum* and *Calamovilfa longifolia*. The presence of blunt sedge (*Carex obtusata*) sets this community apart from other communities found on eolian deposits throughout the grasslands (Meijer and Karpuk 1999), namely the *Stipa comata* - *Koeleria macrantha* - *Agropyron smithii* community described above. This community is not documented elsewhere in the literature and the provincial abundance of this community is not clear.

Stipa comata - *Poa sandbergii* - *Carex obtusata*

Needle and thread grass – Sandberg bluegrass - Blunt sedge

Preliminary Rank: SU

In the Central Parkland, this community is documented by Meijer and Karpuk (1999) from one site in Dillberry Lake Provincial Park. A description of dominant species coverages is not provided. This community is found in morainal depressions with imperfect drainage, subhygric moisture conditions and a mesotrophic nutrient regime. Soils are Gleyed Solonchic Black Chernozems with a clay or clay-loam texture. Common associates include *Achillea millefolium*, *Antennaria parviflora*, *Astragalus spp.* and *Anemone cylindrica*. The presence of *Grindelia squarrosa* in this community indicates saline conditions. This community was not found elsewhere in the literature and the provincial abundance is not clear.

Stipa comata - *Stipa viridula* - *Festuca hallii*

Needle and thread grass – Green needle grass – Plains rough fescue

Preliminary Rank: S3

In the Central Parkland, this community is documented by Bradley and Bradley (1977) from one area near the Neutral Hills south of Provost, Alberta. This community is described as a common grassland community occurring on slopes with little to no grade and various aspects. The three species form a co-dominant, fairly even mixture of *Stipa* and *Festuca*. Soil and site conditions are not discussed. This community is ranked S3 because contains rough fescue (considered provincially threatened), and is a community type that has been reduced drastically in area by agricultural development (A.H. Weerstra pers comm.). It is not well represented in the literature, its provincial abundance is not known and an assessment of structural variability between occurrences is not available.

Stipa curtisetata - Artemisia frigida

Western porcupine grass – Pasture sagewort

Preliminary Rank: S4

In the Central Parkland, this community is described from three areas; one area near Rumsey by Fehr (1982), one area near Dry Island Buffalo Jump Provincial Park by Wroe (1971), and one area near Kinsella, Alberta by Anderson (1972). *A. frigida* can have a wide range of cover values (15-54%) and a frequency of 80-100% between sites (Wroe 1971). Together the dominant species have roughly 70% cover. Anderson (1972) describes a significant (but unspecified) coverage of *Agropyron trachycaulum* in this community. This type occupies the area from the tops of the hills to the intermediate slope position and is confined to south and south west-facing, xeric and subxeric sites on the crests and upper slopes of hills. Soils are rapidly and well drained Orthic, Calcareous and Regosolic Dark Brown Chernozems. Other associated species include *Selaginella densa*, *Festuca scabrella*, *Thermopsis rhombifolia*, *Koeleria cristata* and *Anemone patens* at unspecified coverages. This community is considered common but it is restricted to south-facing slopes (Wroe 1971, Fehr 1982), possibly due to over-grazing on north-facing slopes.

Stipa curtisetata - Bouteloua gracilis

Western porcupine grass – Blue grama

Preliminary Rank: S5

This community is described by Coupland (1950) from 43 sites in the Dark-Brown soil zone which includes the majority of the central and southern Central Parkland subregion of Alberta and Saskatchewan. This community occurs on deep glacial deposits and is also prominent on coarse textured soils in drier areas. It is the characteristic climax community occurring on medium-textured soils developed on undifferentiated glacial till deposits in the moister part of the brown soil zone and the drier part of the dark brown soil zone. In drier areas this community occurs in lower-slope positions and in moister areas occurs on upper to top-slope positions. Specific site

conditions are not discussed but appear to be variable in moisture and soil texture. An assessment of species and structural variability between occurrences is not available. Associated species, including *Artemisia frigida* and *Gutierrezia sarothrae*, can be present but are scarce and low growing. Shrubs including *Symphoricarpos occidentalis*, *Rosa spp.* and *Elaeagnus commutata* are present in areas of increased moisture penetration but when present are not abundant (less than 1%). The largest portion of the Mixed Prairie area in Canada is occupied by the *Stipa - Bouteloua* association (Coupland 1950).

Stipa viridula - Poa sandbergii - Carex obtusata
Green needle grass – Sandberg blue grass – Blunt sedge
Preliminary Rank: SU

In the Central Parkland, Meijer and Karpuk (1999) describe a single occurrence of this community in Dillberry Lake Provincial Park. A structural description with coverages of the dominant species is not provided, but this community is found in an area of imperfect drainage where the moisture regime is subxeric and nutrient conditions are mesotrophic. Soils are Gleyed Solonetzic Black Chernozems. Other species present include *Grindelia squarrosa*, *Achillea millefolium*, *Antennaria parviflora*, *Bromus inermis* and *Koeleria macrantha*. The presence of *Grindelia squarrosa* indicates potentially saline conditions. This community is not found elsewhere in the literature and an assessment of provincial abundance is not available.

Temperate or subpolar grasslands on sandy substrate

CALAMOVILFA LONGIFOLIA SANDY HERBACEOUS ALLIANCE

Calamovilfa longifolia - Artemisia frigida - Koeleria macrantha - Festuca saximontana - Festuca hallii / Selaginella densa
Sand grass – Pasture sagewort – June grass – Rocky mountain fescue – Plains rough fescue / Prairie Selaginella
Preliminary Rank: S4

In the Central Parkland, this community is described by Fehr (1984) from ten sites near Wainwright, Alberta. It is described as a grassland community where the herb-dwarf shrub cover ranges from 40-70% on xeric, rapidly drained, sand flats and outwash plains with Dark Brown Chernozemic soils (Fehr 1984). Moisture conditions are xeric to subxeric and nutrient conditions are not discussed. Associated species include *Opuntia fragilis* and *Rhus radicans*. This grassland community is found frequently within the Central Parkland upon undulating sand landscapes (H. Loonen pers. comm.). It is considered a common grassland community in the sandhills of the Central Parkland (Fehr 1984) but is restricted in range to sandy areas (A.H. Weerstra pers comm.). This community is not found elsewhere in the literature.

Natureserve (2001) describes a *Calamovilfa longifolia* - *Carex inops* ssp. *heliophila* (Herbaceous Vegetation) prairie sandreed grassland associated with *Koeleria macrantha*, *Schizachyrium scoparium*, and *Stipa comata* occurring on sandy slopes in the Great Plains of the United States and Canada. This community has two similar dominants (*Calamovilfa longifolia* and *Koeleria macrantha*) and similar site conditions, but associated species and a *Carex inops* co-dominant differentiate this community from Central Parkland types. The Natureserve website refers the reader to Bourgeron and Engelking (1994), Driscoll *et al.* (1984), Hansen (1985), Hansen *et al.* (1984), Hansen and Hoffman (1988), Hansen and Whitman (1938), Hirsch (1985), and USFS (1992) for information on *Calamovilfa* types described from the USA, all of which appear to be different in composition to the Parkland type described above.

MUHLENBERGIA ASPERIFOLIA SANDY HERBACEOUS ALLIANCE

Muhlenbergia asperifolia - *Scirpus nevadensis* - *Distichlis stricta*
Scratch grass – Nevada bulrush – Salt Grass
Preliminary Rank: S2

This community is described by Wallis (1990) as a hypersaline dry meadow type occurring generally within the Grassland and Parkland Regions of Alberta. In this type *Muhlenbergia asperifolia* occasionally dominates to the exclusion of *Scirpus nevadensis* and *Distichlis stricta* but there is usually a mixed community of these three species (Wallis 1990). A comparison of species and structural variability between occurrences is not possible. Specific soil types and site conditions are not discussed, but this community occurs where sandy soils have been disturbed. *Muhlenbergia asperifolia* is currently ranked S2 on the Alberta Vascular Plant Tracking List (ANHIC 2001) and *Scirpus nevadensis* is not considered common in the province. Although not documented, the frequency of these two species occurring together as a community would most likely be very localized (Wallis 1990, C. Wallis pers comm.). This community is not described elsewhere in the literature.

PUCCINELLIA NUTTALLIANA SANDY HERBACEOUS ALLIANCE

Puccinellia nuttalliana - *Distichlis stricta*
Nuttall's salt meadow grass – Salt grass
Preliminary Rank: S4

This community is described by Wallis (1990) as a hypersaline dry meadow type occurring generally within the Grassland and Parkland Regions of Alberta. It is described as occurring on open, sandy sites ecotonal between bands of *Puccinellia nuttalliana* and *Scirpus nevadensis* with the dominant

species occurring at unspecified coverages. Specific soil and site conditions are not discussed and an assessment of structural variability between occurrences is not available. Associated species include *Hordeum jubatum* and *Agropyron smithii*. This is considered a common community in Alberta (C. Wallis pers comm.) and continuous stands of *Puccinellia nuttalliana* are commonly found associated with *Distichlis stricta* in British Columbia (Parish *et al.* 1996). *Distichlis stricta* is ranked S4 in Alberta, so this is reflected in the community rank. This community is not found elsewhere in the literature.

SPARTINA GRACILIS HERBACEOUS ALLIANCE

Spartina gracilis - *Agropyron smithii*
Alkali cord grass – Western wheat grass
Preliminary Rank: S2S3

This community is described by Wallis (1990) as a saline dry meadow type occurring generally within the Grassland and Parkland Regions of Alberta. Specific site and soil conditions are not discussed, but this community occurs on open, sandy sites between driest portion of the (unspecified) halophytic communities and the (unspecified) semi-halophytic communities occurring further upslope (Wallis 1990). This community is not represented elsewhere in the literature. This type is not considered common and grows in narrow bands in very specific saline/sandy habitat types limiting its extent (C. Wallis, pers comm.). Information is not available to assess species or structural variability between occurrences.

STIPA CURTISETA SANDY HERBACEOUS ALLIANCE

Stipa curtisetia - *Koeleria macrantha* - *Agropyron trachycaulum* / *Cladina mitis*
Western porcupine grass – June grass – Slender wheat grass / Yellow reindeer lichen
Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from three sites in Dillberry Lake Provincial Park. Lichen cover averages 72%, with grass cover inversely related to lichen cover, ranging from 6 to 25%. Forb cover is limited. It is associated with shallow, south-facing slopes in sandy areas of dune fields. It is found on south to south-east aspects with rapid drainage and submesotrophic nutrient conditions. Moisture conditions are xeric and soils are Ortho Regisolic to Orthic Dark Brown Regisols. Associated species include *Arctostaphylos uva-ursi*, *Juniperus horizontalis* and *Symphoricarpos occidentalis* at unspecified coverages. This community is not found elsewhere in the literature, so provincial abundance of this community type is not known.

Temperate or subpolar seasonally flooded grasslands

HORDEUM JUBATUM SEASONALLY FLOODED HERBACEOUS ALLIANCE

Hordeum jubatum

Foxtail barley

Preliminary Rank: S4

In the Central Parkland of Alberta, this community type is described by Fehr (1982) from near Rumsey, and by Wallis (1990) as a saline dry meadow type occurring generally within the Parkland and Grassland Regions of Alberta. This community type occurs as a wide band of vegetation around saline sloughs. It characterizes sites on higher margins of the wetlands where more permanent cover has been destroyed by periodic flooding and desiccation (Wallis 1990). In Alberta, *Hordeum jubatum* forms almost pure stands with covers up to 40%. This community is described elsewhere (see below) and is likely not uncommon. An assessment of structural variability between occurrences of this type in Alberta is not possible with current information. This community is ranked S4 to coincide with its global rank assigned by The Nature Conservancy (Natureserve 2001).

Natureserve (2001) describes a similar community type with similar site conditions reported from Montana, Utah, generally from the Great Plains of the United States, and within western Canada (ranked G4 by The Nature Conservancy). However, references for this community type in the literature are not strong (undergraduate projects, draft reports). The vegetation is dominated by short and medium-tall graminoids (unspecified species) with a total vegetation cover of nearly 100%. Shrubs are usually absent from this community (Natureserve 2001). In Utah, stands are found in lowlands with moderately to strongly saline soils. The topography is flat and the soils are often flooded or saturated in the spring. In Montana, this community is a seasonally flooded subsaline wetland type, and a drawdown phase of seasonal and semipermanent brackish and subsaline marshes (Heidel *et al.* 2000, Thompson 1994).

PUCCINELLIA NUTTALLIANA HERBACEOUS ALLIANCE

Puccinellia nuttalliana

Nuttall's salt meadow grass

Preliminary Rank: S3

In the Central Parkland, this community is described by Fehr (1982) from one site in the Rumsey area, and as occurring generally as a saline emergent marsh or a hypersaline dry meadow within the Parkland and Grassland Regions of Alberta by Wallis (1990). This community is typified by almost pure stands of *Puccinellia nuttalliana* with a coverage of < 20%. It can be found on fine, sandy loam to heavy clay textured saline meadow soils, and subhydric saline depressions, where it occurs in a wide band around saline sloughs (Fehr 1982,

Wallis 1990). Specific site conditions are not discussed. Other species that can be present include *Triglochin maritima*, *Salicornia europaea*, *Scirpus nevadensis*, *Distichlis stricta*, *Suaeda calceoliformis* and *Hordeum jubatum*.

As a species, *Puccinellia nuttalliana* can be found throughout central and southern Alberta (Moss 1983), but it is not well represented in the Alberta literature as a community type. Although not formally considered rare in the province, this community type is ranked G3 by The Nature Conservancy (Natureserve 2001) because it has specific habitat needs; this community requires moist soils of intermediate salinity in seasonally wet meadows, and has been impacted by water diversion, livestock grazing, and land conversion in many places (Ungar 1974).

Elsewhere, similar communities of *Puccinellia nuttalliana* are found in the Great Plains and intermountain region, extending from Alberta and Saskatchewan to Nebraska, and west to Montana, Utah and California (Heidel *et al.* 2000). This community is characterized by the dominance of *Puccinellia nuttalliana* in the graminoid layer. Trace amounts of *Distichlis stricta* or *Hordeum jubatum* can be associated with this community. The forb layer is relatively sparse and is often composed of *Salicornia rubra* or *Triglochin maritima*. Topographic position and saline conditions of these communities are very similar to Central Parkland types.

PHALARIS ARUNDINACEA HERBACEOUS ALLIANCE

Phalaris arundinacea - *Carex atherodes* - *Calamagrostis stricta*

Narrow reed grass – Awned sedge - Reed canary grass

Preliminary Rank: S5

This community is described from one site near Sylvan Lake on the western edge of the Central Parkland by Griffiths and Griffiths (1987a). Quadrat coverages of *Phalaris arundinacea* and *Carex atherodes* were approximately 25-50%, with *Calamagrostis stricta* at approximately 6-15%. This community is considered “mature” and occurs along shorelines beyond the limits of willow growth where water depth does not exceed 20 cm. Soils are Hydric Humisols and deadfall is present at trace amounts. Associated species include *Carex aquatilis* and, where water is deeper, *Sagittaria cuneata* and *Eleocharis palustris* all at levels of <5%. Adjacent communities further from the water include “willow shrubbery”, *Typha latifolia* followed by a broad, monoculture zone of *Scirpus validus*. This community is considered a common community type around lakes and sloughs in Alberta (G. Griffiths pers comm.) but is not well represented in the literature. Because this type is only described from one site, an assessment of species and structural variability between occurrences is not possible.

Temperate or subpolar intermittently flooded grasslands

CALAMAGROSTIS INEXPANSA INTERMITTENTLY FLOODED HERBACEOUS ALLIANCE

Calamagrostis inexpansa - *Juncus balticus*

Northern reed grass – Wire rush

Preliminary Rank: S5

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from two sites in Dillberry Lake Provincial Park. Because this type is only described from two sites in the same area, an assessment of species and structural variability between occurrences is not possible. Coverages for dominant species are not provided. This community is associated with receding lakeshores Orthic Regosolic and Rego Gleysolic soils in poorly drained sites with mesic to hygric moisture regimes and permesotrophic nutrient conditions. Associated species include *Populus balsamifera*, *Salix bebbiana*, *Salix candida*, *Solidago canadensis* and *Carex praegracilis*. The *Calamagrostis inexpansa* – *Juncus balticus* community type is not described elsewhere in the literature.

HORDEUM JUBATUM INTERMITTENTLY FLOODED HERBACEOUS ALLIANCE

Hordeum jubatum - *Puccinellia nuttalliana*

Foxtail barley - Nuttall's salt meadow grass

Preliminary Rank: S4

In the Central Parkland, this community is described by Griffiths *et al.* (1996) from one site near Lister Lake, Alberta. It is a patchy grass community that occurs under beach ridges exposed for more than two years. Specific site and soil conditions are not discussed. *Hordeum jubatum* is dominant or co-dominant and can almost reaches monocultures in some areas. Hydrological regime differentiates this community from the *H. jubatum* community described earlier (i.e. this community is not seasonally flooded). Associated species can include *Scirpus pungens*, *Eleocharis palustris*, *Juncus balticus*, *Rumex maritimus*, *Sonchus arvensis*, *Aster brachyactis*, *Ranunculus cymbalaria* and *Taraxacum officinale* at unspecified coverages. This community is not threatened, but somewhat limited in extent as it is found only from oligohaline lakes with fluctuating water levels (e.g. Beaverhill Lake, G. Griffiths pers comm.). Because it is only described from one site, an assessment of species and structural diversity between occurrences is not possible.

(Herbaceous)

Composites

Oblong, spatulate-leaved

Temperate or subpolar composite grasslands

ANTENNARIA PARVIFOLIA HERBACEOUS ALLIANCE

Antennaria parvifolia - *Artemisia frigida* - *Koeleria macrantha*

Small leaved everlasting – Pasture sagewort – June grass

Preliminary Rank: SU

In the Central Parkland, this community is described by Fehr (1984) from ten sites near Wainwright (David Lake), Alberta. Fehr (1984) estimated percent cover for the three main species at 35%, 20%, and 20%, respectively. Other species present can include *Juncus balticus* (15%), *Carex spp.* (10%) and *Poa spp.* (10%). The variability in species and structure between occurrences is not reported. This community can be found on hilltops and dry areas. Specific soil and site conditions are not discussed. The absence of important forage species (*Agropyron trachycaulum*, *Bouteloua gracilis*, etc.) is a reflection of the heavy grazing this community type receives. This community is not reported elsewhere in the literature and its provincial abundance is unknown.

(Herbaceous)

(Composites)

Twice-pinnate leaves

Temperate or subpolar composite grasslands

ARTEMISIA FRIGIDA HERBACEOUS ALLIANCE

Artemisia frigida / *Koeleria macrantha* - *Carex obtusata*

Pasture sagewort / June grass – Blunt sedge

Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from two sites in Dillberry Lake Provincial Park. Coverages of the dominant species are not provided, so an assessment of variability between occurrences is not possible. It is described as being closely associated with south-facing slopes of glaciofluvial origin. It is found on xeric to very xeric sites with very rapid drainage and Orthic and Rego Dark Brown Chernozemic soils. Associated species can include *Symphoricarpos occidentalis*, *Heterotheca villosa*, *Thermopsis rhombifolia*, and *Agropyron trachycaulum* at unspecified amounts. This community is not represented elsewhere in the literature and its provincial abundance is not known.

(Herbaceous)

(Composites)

Leaves large, petioled

Temperate or subpolar permanently flooded composite wetlands

POLYGONUM AMPHIBIUM PERMANENTLY FLOODED HERBACEOUS ALLIANCE

Polygonum amphibium

Water smartweed

Preliminary Rank: S4

This wetland vegetation type is not formally documented from the Central Parkland of Alberta, but is known to occur in shallow water along the edges of ponds and lakes, with floating-leaved aquatic forbs covering roughly 30% of the water's surface (Kunze 1994). *Polygonum amphibium* often forms dense, nearly monotypic stands. *Lemna minor*, *Potamogeton natans*, *Sagittaria spp.*, *Spirodela polyrhiza* and *Wolffia spp.* are occasionally present. This community is currently on the Alberta Plant Community Tracking List (Allen 2001) and is ranked SU. Little information is available for Alberta communities. The species *Polygonum amphibium* is noted as widespread in Alberta (Moss 1983) and the occurrence of this species as a community type is not considered rare (C. Wallis pers comm.).

Johnson (1939, 1941) found this community in Colorado around montane lakes with significant sediment deposition. In Montana, Heidel *et al.* (2000) note this community from disturbed agricultural sites that are seasonally inundated and slightly brackish (ranked S? for Montana). It is also found around reservoirs, ponds and other wet areas in Montana.

Shrubland

Deciduous

Broad-leaved

Temperate or subpolar cold-deciduous shrubland

ELAEAGNUS COMMUTATA SHRUBLAND ALLIANCE

Elaeagnus commutata

Silverberry

Preliminary Rank: S2

In the Central Parkland, this community is described from three different sites; one site in Dillberry Lake Provincial Park by Meijer and Karpuk (1999), one site near Rumsey by Fehr (1982) and one site in Dry Island Buffalo Jump by Bradley and Bradley (1972). In this community *Elaeagnus commutata* forms a shrub layer of approximately 50% cover (Fehr 1982). This type occurs as low shrublands and shrubby meadows along the perimeter of saline lakes

adjacent to marshes or graminoid meadows. It occupies the middle and lower positions of mesic to submesic slopes, and is found in areas that are moderately to well drained with northern or southern aspects often in proximity to alkaline water bodies. This shrubland community possesses xeric moisture conditions and submesotrophic nutrient conditions with Orthic Dark Brown soils. Common associates include *Symphoricarpos occidentalis* (potentially reaching co-dominance in some cases according to Fehr (1982)), *Rosa acicularis*, *Carex spp.*, *Calamovilfa longifolia* and *Agropyron trachycaulum* at unspecified (but assumed trace) levels. Characteristic herbs can include *Hackelia americana*, *Galium boreale* and *Rorippa islandica* all at coverages <5%.

Fehr (1982) describes an *Elaeagnus commutata* - *Symphoricarpos occidentalis* community in the Rumsey area of Alberta. Meijer and Karpuk (1999) describe an *Elaeagnus commutata* community in Dillberry Lake Provincial Park. This shrubland community is typically well developed and dense with a poorly developed herb-dwarf shrub layer. In terms of abundance, Bradley and Bradley (1977) considered *E. commutata* to be more abundant as one goes south of the Rumsey area and this community is considered as common in the Central Parkland of Alberta (H. Loonen pers comm.). The species *Elaeagnus commutata* can be found throughout central and southern Alberta, but the occurrence of this species as a community type is not well documented. The Nature Conservancy (Natureserve 2001) ranks *Elaeagnus commutata* as G2, requiring an S2 rank for Alberta.

Elsewhere, similar communities of silverberry shrubland are reported in Montana east of the continental divide with shrub coverages around 10% and a *Stipa comata* understory of up to 70% (Heidel *et al.* 2000). Natureserve (2001) describes an *Elaeagnus commutata* shrubland, but it is unclear whether this is the same community found in Alberta because the site conditions and species coverages are not adequately described. *Elaeagnus commutata* Shrubland is considered a minor community type forming narrow stringers on stream terraces in northern Montana, east of the Continental Divide (Natureserve 2001).

Elaeagnus commutata / *Symphoricarpos occidentalis* - *Rosa woodsii* / *Poa palustris*
Silverberry / Snowberry – Common wild rose / Fowl bluegrass
Preliminary Rank: SU

In the Central Parkland, this community is described by Fehr (1984) from one site near Wainwright, Alberta. Unlike the *Elaeagnus commutata* cover type described above, this community has a tall shrub layer of *Symphoricarpos occidentalis* and a characteristic component of *Rosa woodsii*. This type can be found in subxeric and submesic locations where drainage is good (Fehr 1984). It has a well developed low shrub layer (75-80% cover) and occurs on Dark Brown Chernozemic soils. Other species occurring at levels less than

5% include *Achillea millefolium*, *Thermopsis rhombifolia*, *Disporum trachycarpum*, *Fragaria virginiana*, *Ribes oxyacanthoides* and *Galium boreale*. This community is only documented once in the literature, preventing any assessment of structural variability between occurrences, and its provincial abundance is therefore not certain.

POPULUS TREMULOIDES SHRUBLAND ALLIANCE

Populus tremuloides - *Amelanchier alnifolia* - *Prunus virginiana* / *Rosa* spp.
Trembling aspen – Saskatoon – Choke cherry / Rose spp.
Preliminary Rank: S5

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from three sites in Dillberry Lake Provincial Park associated with steep, north aspects. This community is a young, shrub-level successional stage of regenerating aspen in areas where the canopy has opened from natural disturbance (e.g. windthrow, beaver, natural forest edge). Drainage is rapid, moisture regimes are submesic, and nutrient conditions are typically submesotrophic to mesotrophic. Soils tend to be Rego Dark Brown Chernozems, Orthic Regosols and Cumulic Regosols. Other species present can include *Rubus idaeus*, *Ribes oxyacanthoides*, *Aralia nudicaulis*, *Smilacina stellata*, and *Schizachne purpurascens*. We have observed this community in upland clearings and well-drained, open woods in the Boreal and Foothills Regions of Alberta. This community was not found elsewhere in the literature so an assessment of structural variability between occurrences is not possible.

PRUNUS VIRGINIANA SHRUBLAND ALLIANCE

Prunus virginiana - *Amelanchier alnifolia* / *Agropyron trachycaulum* - *Poa pratensis*
Choke cherry – Saskatoon / Slender wheat grass – Kentucky blue grass
Preliminary Rank: S5

In the Central Parkland, this community is described by Griffiths *et al.* (1997) from one site in the Blackfoot Provincial Recreation Area. It is described as “dryland shrubbery”; cleared areas without aspen saplings where the low shrubs grow mixed with tall grasses. Soil conditions are not discussed, but this type was located on a steep, south-facing slope. This community is considered common in the parklands of the prairie provinces, but becomes restricted to south-facing slopes in forested areas where it is described by Griffiths *et al.* (1997). This dryland shrubbery can be distinguished from aspen-dominated shrubland and beaver-cleared areas by the absence of aspen suckers. Looman (1983) classifies this vegetation in the alliance “*Prunion melanocarphae*”, but does not discuss abundance or variation within occurrences.

ROSA ACCICULARIS SHRUBLAND ALLIANCE

Rosa acicularis – (shrub) / *Solidago spp.*

Prickly rose

Preliminary Rank: SU

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from one site in Dillberry Lake Provincial Park. A structural description is not provided for this community. It can be found on subxeric, west-facing slopes with rapid drainage, Rego Dark Brown soils, and submesotrophic nutrient conditions. Associated species can include *Amelanchier alnifolia*, *Prunus virginiana*, *Solidago spp.*, *Heterotheca villosa* and *Agropyron spp* at unspecified (but assumed trace) amounts. This rose-dominated community is not found elsewhere in the literature and its provincial abundance is not certain.

Rosa acicularis - *Rubus idaeus* / *Lathyrus venosus*

Prickly rose – Wild red raspberry / Purple pea vine

Preliminary Rank: S5

In the Central Parkland, this community is described by Timoney and Robinson (1998) from nine sites in the Blackfoot Provincial Recreation Area. This community forms when beavers, pathogens, or insect outbreaks remove large portions of the forest canopy allowing shrub layers to develop. Cover values for the dominant species are not provided. It is a beaver-dependent shrubland type typical of well drained slopes near beaver ponds. It is found in areas of submesic to subhygric moisture regimes, in Orthic Grey Luvisolic soils. Associated species include *Symphoricarpos occidentalis*, *Amelanchier alnifolia*, *Prunus virginiana*, *Salix bebbiana*, and *Calamagrostis canadensis* at unspecified (but assumed trace) amounts. It is differentiated from the *Rosa acicularis* – (shrub) / *Solidago spp.* community described above because this community occurs in moister site conditions, different soil types (i.e. not Rego Dark Brown) and it is characterized by the presence of *Rubus idaeus* as a dominant species. We have observed this community on similar site conditions associated with mature aspen forest openings in the Central and Dry Mixedwood subregions of Alberta. Although a formal assessment of its provincial abundance is not available, this community should not be considered a rare type.

SALIX SPP. SHRUBLAND ALLIANCE

Salix petiolaris / (low shrubs and herbs)

Basket willow

Preliminary Rank: S5

In the Central Parkland, this community is described by Fehr (1982) from one site near Rumsey, and by Wroe (1971) from one site near Dry Island Buffalo Jump, Alberta. A description of the structure and cover of the dominant species is not provided. It occupies moist banks around sloughs, forms rings or clumps around mesic depressions and is generally restricted to moist areas with rolling topography. It occurs in mesic to hydric moisture regimes with moderately to poorly drained Rego and Orthic Humic Gleysolic soils. This community is surrounded to varying degrees by *Populus tremuloides* forest (community type not specified). *Salix lutea* and *Populus balsamifera* are sometimes present as trees. Associated shrubs and herbs can include *Rosa woodsii*, *Amelanchier alnifolia*, *Symphoricarpos occidentalis*, *Poa palustris*, *Epilobium angustifolium*, *Beckmannia syzigachne*, *Calamagrostis spp.*, *Carex rostrata* and *Carex atherodes* at unspecified levels.

Fehr (1982) describes a *Salix petiolaris* / *Rosa woodsii* / *Sonchus spp.* community from near Rumsey, Alberta, and Wroe (1971) describes a *Salix petiolaris* / *Rosa woodsii* / *Sonchus arvensis* community from near Dry Island Buffalo Jump Provincial Park. Both types occur on identical site conditions with the same associated species and are considered the same community type in this report. This community is not documented elsewhere in the literature. *Salix petiolaris* is common in central Alberta (Moss 1983) and when present often forms communities as thickets or clumps in wet areas, but a formal assessment of abundance is not available for this community.

SYMPHORICARPOS OCCIDENTALIS SHRUBLAND ALLIANCE

Symphoricarpos occidentalis

Buckbrush

Preliminary Rank: S4

In the Central Parkland, this community is described from several locations; two sites in Dillberry Lake Provincial Park (Meijer and Karpuk 1999), one site near Kinsella, one site near Rumsey (Fehr 1982), and one site near Dry Island Buffalo Jump (Wroe 1971). In the Central Parkland, this community is a dense shrubland ranging in size from 0.004 to 0.30 ha (Wroe 1971) where total shrub cover averages 80% (range 75-95%). A large quantity of dead stems of *Symphoricarpos occidentalis* can be an important component of this community. It occupies shallow depressions on the grassland, and in many instances positions of higher elevations adjacent to the aspen communities. This community can be found on north/east and south/west-facing slopes on rapidly drained sites with submesotrophic nutrient conditions, along the perimeters of naturally open woodlands or in association with fescue grasslands. It is associated with dry conditions, and Orthic and Rego Dark Brown Chernozemic soils. Associated species include *Elaeagnus commutata*, *Rosa acicularis*, *Prunus virginiana*, *Artemisia frigida* and *Koeleria*

macrantha. *Rosa woodsii* and *Ribes oxycanthoides* can also be present all at <10% cover.

Anderson and Bailey (1980) mention a similar community found in *Populus tremuloides* parkland from east-central Alberta associated with *Agropyron spp.*, *Stipa spartea* and *Elaeagnus commutata*. Fehr (1982) and Wroe (1971) both describe this community from the Rumsey area. All of these types possess the same site conditions and associated species and are all considered the same community type in this report. The Nature Conservancy (Natureserve 2001) has ranked this community type as G4, and it is therefore ranked S4 in Alberta.

Elsewhere, similar *Symphoricarpos occidentalis* communities are reported from Saskatchewan, Missouri, and Montana in mesic depressions, typically surrounded by upland grassland associations (Heidel *et al.* 2000). This community type has also been described for Montana, North Dakota (Hansen *et al.* 1984, Hansen *et al.* 1990) and is reported as common in the Saskatchewan parkland (Coupland *et al.* 1953). Natureserve (2001) describes a western snowberry shrubland from the western tallgrass and northern Great Plains of the United States and Canada where *Symphoricarpos occidentalis* is the predominant species in the shrub layer and at times forms almost monotypic stands. Associated species occurring at the fringe of this community include *Rosa woodsii*, *Rhus aromatica* and *Prunus virginiana*. The herbaceous layer is poorly represented where the shrubs are dense but can include *Artemisia ludoviciana*, *Solidago spp.*, *Poa pratensis* and *Achillea millefolium*. This type is frequently observed in heavily grazed meadows and prairies. It is tolerant of both grazing and fire (Hansen and Hoffman 1988), and is under no threat from human activities (Natureserve 2001).

Symphoricarpos occidentalis / *Antennaria parvifolia* - *Festuca hallii*
Buckbrush / Small leaved everlasting – Plains rough fescue
Preliminary Rank: SU

In the Central Parkland, this community is described from one site in the Wainwright Ecological Reserve by Pearson Timberline (1993). The inclusion of *Antennaria parvifolia* and *Festuca hallii* as community dominants differentiates this community from the *Symphoricarpos occidentalis* community described above. A description of coverage or structure of the dominant species is not provided. This community is found on sandy, outwash plains with Dark Brown Chernozem and Humic Gleysolic (calcareous) soils. Slopes can range from 0-10% in all directions, sites are well to moderately-well drained, and nutrient conditions are submesic to mesic. Other shrub species can include *Elaeagnus commutata* and *Rosa spp.* Forbs and grasses can include *Selaginella densa*, *Carex spp.* and *Stipa spartea*. This community is not represented elsewhere in the literature and its provincial abundance is unknown.

Temperate or subpolar permanently flooded cold-deciduous shrubland

BETULA PUMILA PERMANENTLY FLOODED SHRUBLAND ALLIANCE

Betula pumila / *Carex aquatilis*

Dwarf birch / Water sedge

Preliminary Rank: S5

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from one site in Dillberry Lake Provincial Park. A description of dominant-species structure or coverage is not provided. It is described as a classic fen possessing a floating mat of organic material greater than 140 cm in thickness. Soils are Hydric Humisols. Drainage is very poor, moisture conditions are hydric and nutrient conditions are eutrophic. Associated species include *Salix serissima*, *Carex aquatilis* and *Drepanocladus spp.* Floating fens are considered unusual in the Central Parkland (Meijer and Karpuk 1999) but we have observed this community in the Boreal Region of Alberta, and similar types are described elsewhere.

A *Betula pumila* – *Salix spp.* community type is described from southwestern Alberta by Archibald *et al.* (1996). This “dwarf birch / sedge / peat moss” community occurs in similar site conditions and can include *Larix laricina*, *Carex spp.* and *Sphagnum spp.* A *Betula pumila* – *Salix spp.* community type is described from the northern tallgrass prairie and adjacent prairie-forest border regions of the upper midwestern United States, where surface water pH is neutral to somewhat alkaline and has moderate nutrient levels (Natureserve 2001). It is typically found on relatively shallow peat. This community has a shrub layer with 25-70% cover, abundant herbaceous species, and sparse to abundant mosses. Associated species include *Calamagrostis canadensis*, *Calamagrostis stricta*, *Carex aquatilis*, *Carex lasiocarpa*, *Eriophorum angustifolium*, *Euthamia graminifolia*, *Lobelia kalmii*, *Lycopus uniflorus*, and *Triadenum fraseri*. Valid comparisons of this community type to the type found in the Central Parkland difficult without identifying the *Salix* to species.

SALIX PYRIFOLIA PERMANENTLY FLOODED SHRUBLAND ALLIANCE

Salix pyrifolia - *Betula neoalaskana* / *Ledum groenlandicum* - *Calamagrostis canadensis* / *Sphagnum warnstorffii*

Balsam Willow – Alaskan birch / Labrador tea – Marsh reed grass / Warnstorff's peat moss

Preliminary Rank: S5

In the Central Parkland, this community is described by Griffiths *et al.* (1997) from one site in the Blackfoot Provincial Recreation Area. This community is found in oligotrophic, perched morainal depressions and on bog margins

sometimes intergrading with taller swamp vegetation. Associated species, specific site conditions and soil types are not discussed, however, the dominance of balsam willow would indicate relatively oligotrophic conditions. Griffiths *et al.* (1997) describe this community as a Balsam Willow Carr; the term “Carr” used for fens with extensive shrub coverage but with few or no trees. This community is considered common throughout the Boreal and Parkland Regions of Alberta (Griffiths *et al.* 1997) but is not discussed elsewhere in the literature.

Salix pyrifolia / *Carex utriculata*
Balsam willow / Greater beaked sedge
Preliminary Rank: SU

In the Central Parkland, this community is described by Griffiths and Griffiths (1987b) from one site near Onoway, Alberta. This balsam willow Carr occurs with a dense and diverse herb coverage dominated by *Carex utriculata*. Variation in water levels prevents willow colonization from suppressing herb growth. Specific site and soil conditions are not discussed, but other herbaceous associates, depending on water depth variation, include *Glyceria grandis*, *Polygonum amphibium* and *Lemna minor*. This community is not well represented in the literature and its provincial abundance is not known.

Temperate or subpolar semi-permanently flooded cold-deciduous shrubland

***SALIX PETIOLARIS* SEMI-PERMANENTLY FLOODED SHRUBLAND**

Salix petiolaris - *Salix planifolia* / *Carex utriculata*
Basket willow – Tea-leaved willow / Greater beaked sedge
Preliminary Rank: S5

In the Central Parkland, this community is described by Griffiths and Griffiths (1987b) from one site near Onoway, Alberta. This community occurs in flooded areas where *Salix pyrifolia* is rare or entirely lacking. Specific site and soil conditions are not discussed and an estimation of the coverage of dominant species is not provided. Vegetation around the periphery of this community can include *Salix petiolaris*, *Carex aquatilis*, *Glyceria grandis*, *Scirpus microcarpus*, *Galium trifidum*, *Lysimachia thyrsiflora*, *Bidens cernua*, *Lemna minor* and *Calamagrostis canadensis* at unspecified levels. This community is considered as common around lakes and sloughs in Alberta (G. Griffiths pers comm.) but it is not found elsewhere in the literature.

Salix petiolaris / *Juncus balticus*
Basket willow / Wire rush
Preliminary Rank: S5

In the Central Parkland, this community is described by Griffiths *et al.* (1996) from one site near Lister Lake, Alberta. This community is described as willow shrubbery; a stable edaphic climax community on land with a water table too high for balsam poplar to reach maturity. Site and soil conditions are not discussed. Associated species include *Juncus balticus*, *Calamagrostis stricta* and *Poa pratensis* at unspecified levels. The moss *Drepanocladus aduncus* can sometimes be present. This community is considered common around lakes and sloughs in Alberta (G. Griffiths pers comm.) but it is not found elsewhere in the literature.

Temperate or subpolar cold-deciduous shrubland on sandy substrate

POPULUS TREMULOIDES SANDY SHRUBLAND ALLIANCE

Populus tremuloides / *Juniperus horizontalis* / *Cladina mitis*
Trembling aspen / Creeping juniper / Yellow reindeer lichen
Preliminary Rank: SU

In the Central Parkland, this community is described from three sites; one site near Wainwright (Fehr 1984) and three in Dillberry Lake Provincial Park (Renewable Resources Consulting Services 1974, Meijer and Karpuk 1999). Fehr (1984) noted a well-developed shrub layer (approximately 24% cover), aspen being the dominant species, averaging approximately 2m in height. *Juniperus horizontalis* averages 20% cover with *Cladina mitis* averaging 33% cover (Fehr 1984). This type is found on inter-dune depressions, on the upper positions of shallow, south-facing slopes with rapid drainage. Soils are Orthic Dark Brown to Orthic Regisols with xeric moisture regimes and submesotrophic nutrient conditions. Other shrub species with lower abundances (<7%) can include *Betula papyrifera*, *Rosa acicularis*, *Salix bebbiana*, *Gallium boreale*, *Artemisia campestris*, *Agropyron trachycaulum* and *Stipa curtiseta*. The herb-dwarf shrub layer averages 40% cover with *Juniperus horizontalis* dominant, but also containing *Selaginella densa* and *Calamovilfa longifolia*. Bryoides form an extensive ground mat in these stands (30-60% cover according to Fehr 1984) consisting of *Cladina mitis* (the dominant), *Cetraria nivalis*, *Cladonia uncialis*, *Cladonia pyxidata* and *Cetraria ericetorum*. This community is considered common on historically windblown and recently stabilized steep slopes and blowouts of the choppy sand hill terrain of the Central Parkland (H. Loonen, pers comm.), however, it is not well represented in the literature so its provincial abundance is not known.

SYMPHORICARPOS OCCIDENTALIS SANDY SHRUBLAND ALLIANCE

Symphoricarpos occidentalis - *Rosa arkansana* / *Juniperus horizontalis*
Buckbrush – Prairie rose / Creeping juniper
Preliminary Rank: SU

In the Central Parkland, this community is described by Renewable Resources Consulting Services (1975) from one site in Dillberry Lake Provincial Park. This community type is described from the grassland areas as scattered patches of *Symphoricarpos occidentalis* and *Rosa arkansana* with vigorous mats of creeping juniper. Specific site and soil information is not discussed. This community is described only briefly in the literature so its overall abundance is not clear and there is no information on community structure or dominant species abundance.

Sparse

Xeric sandy soil

Dunes and blowouts

Temperate or subpolar sparse graminoids on sand

Calamovilfa longifolia - *Sporobolus cryptandrus* - *Koeleria macrantha* - *Carex obtusata*

Sand grass – Sand dropseed – June grass – Blunt sedge

Preliminary Rank: S3

In the Central Parkland, this community is described by Meijer and Karpuk (1999) from three sites in the southern half of Dillberry Lake Provincial Park. This community can be found on sand dunes or active blowouts. It is generally located on south to west-facing aspects with Regosolic soils and sparse vegetation cover. Conditions are typically xeric with rapid drainage and oligotrophic to submesotrophic nutrient conditions. *Juniperus horizontalis* and *Heterotheca villosa* can also be present. This community is not well represented in the literature. It is restricted to active blowouts and consequently has a limited range (Meijer and Karpuk 1999).

Temperate or subpolar sparse cold-deciduous shrubs on sand

Populus tremuloides / *Juniperus horizontalis*

Trembling aspen / Creeping juniper

Preliminary Rank: SU

This regenerating aspen community was described from one site in Dillberry Lake Provincial Park on a shallow, south-facing slope with rapid drainage (Meijer and Karpuk 1999). It is found on Orthic Dark Brown soils with subxeric moisture conditions and submesotrophic nutrient conditions. Trace grasses, forbs and lichens can include *Solidago missouriensis*, *Calamovilfa longifolia*, *Carex obtusata* and *Cladina mitis*. This community is not found elsewhere in the literature and its provincial abundance is unknown.

Populus tremuloides - *Prunus virginiana* / *Juniperus horizontalis* / *Selaginella densa*
Trembling aspen – Choke cherry / Creeping juniper / Prairie Selaginella
Preliminary Rank: S3

This community type, described from one site in the Wainwright area, can be found on sand dunes with Regosolic soils, rapid drainage, and xeric to very xeric moisture regimes (Pearson Timberline 1993). Slopes can range from 0-50% on south or southwest aspects. Aspen growth is stunted to shrub height. *Carex siccata*, *Calamovilfa longifolia* and *Elymus canadensis* can also be present. This community is not found elsewhere in the literature. It has a limited range on south-facing sand dunes and is thus ranked S3. Because it is only documented from one site, and because the coverage and structure of the community is not described, an assessment of structural variability between occurrences is not possible.

Temperate or subpolar sparse evergreen shrubs on sand

Juniperus horizontalis / (*Koeleria macrantha*) / *Cladina mitis*
Creeping juniper / (June grass) / Yellow reindeer lichen
Preliminary Rank: S1S2

In the Central Parkland, this community is described by Fehr (1984) from one site near Wainwright, Alberta. Fehr (1984) estimated that *Juniperus horizontalis* forms roughly 16% cover, with *Koeleria macrantha* and *Cladina mitis* forming 1.8% and 27% cover, respectively. It is found in stable blowout areas of sand dunes. It occurs in xeric to very xeric areas with southern aspect, rapid drainage, sparse vegetative cover, and Orthic Regosolic soils. Other lichens present can be *Cetraria ericetorum* and *Cetraria nivalis*. The moss *Tortula ruralis* can also be present. Because it occurs on active blowouts in sand dune areas, this community is restricted in area. It is not found elsewhere in the literature and is currently on the ANHIC plant community tracking list with a rank of S1S2 (Allen 2001).

Lake and slough edges Saline wetlands

Saline wetland sparse herbaceous vegetation

Salicornia europaea
Samphire
Preliminary Rank: S2

This community is described by Wallis (1990) as occurring generally within the Grassland and Parkland Regions of Alberta, characteristic of saline Gleyed Regosolic soils at the edge of the non-vegetated portion of hypersaline emergent marshes. It is typified by a band of pure *S. europaea* with less than

25% cover. Associated species are *Hordeum jubatum*, *Triglochin maritima*, *Chenopodium rubrum* and *Suaeda calceoliformis*. This community is not well represented in the literature for Alberta and is included on the Alberta Plant Community Tracking List with a rank of S2 (Allen 2001).

The Nature Conservancy has ranked this community G2. They describe it as being associated with highly alkali wetlands or semipermanent alkali lakes in the northern Great Plains and Great Basin of the United States and adjacent Canada (Natureserve 2001). It often borders intermittent open water or is found on the exposed mud of alkali flats. *Salicornia europaea* may make up to 100% of the vegetation within these areas. Other associated species can include *Puccinellia nuttalliana*, *Distichlis spicata*, *Hordeum jubatum*, *Triglochin maritima*, *Chenopodium rubrum*, and *Suaeda calceoliformis*. This type occurs over a broad geographic range and has been recorded from Nebraska, South Dakota, North Dakota, Minnesota, Montana, Colorado, and north into Saskatchewan (Dodd and Coupland 1966). However, it has very specific habitat needs occurring on the exposed mud of alkali flats and there may be less than 50 occurrences rangewide (Ungar 1974c).

Scirpus nevadensis

Nevada bulrush

Preliminary Rank: S3

This community is described by Wallis (1990) as occurring generally within the Grassland and Parkland Regions of Alberta on sandy saline shores of emergent marshes. The structure, percent cover and variability of this community are not discussed in the literature. *Scirpus nevadensis* is considered uncommon and usually occurs with very open growth in the emergent marshes of the Grassland and Parkland regions. It is restricted to saline shores and a few sand plain landscapes, however, in certain areas it may form extensive stands (Wallis 1990, C. Wallis pers comm.) and is found as far south as Texas (Moss 1983). The occurrence of this species as a community type is not well documented, but it is restricted in extent given its saline habitat requirements.

Scirpus nevadensis - (*Triglochin maritima*)

Nevada bulrush – (Common cattail)

Preliminary Rank: S2S3

This community is described by Wallis (1990) as occurring generally within the Grassland and Parkland Regions of Alberta on sandy saline shores of emergent marshes. The structure, percent cover and variability of this community are not discussed in the literature. *Scirpus nevadensis* is considered uncommon and usually occurs with very open growth in the emergent marshes of the Grassland and Parkland regions. This is a localized community type confined to a few sand plain landscapes in Alberta; the

Triglochin component appears to be the rare feature (C. Wallis pers comm.) and the diagnostic component that differentiates it from the *Scirpus nevadensis* community described above. This community is included on the Alberta Plant Community Tracking List (Allen 2001) and is ranked S1S2 therein.

Suaeda calceoliformis

Western sea blight

Preliminary Rank: S5

This community is described by Wallis (1990) as occurring generally within the Grassland and Parkland Regions of Alberta as a hypersaline emergent marsh dominated by *C. calceoliformis* with a cover of <25% at the edge of non-vegetated saline wetlands. Common associated species are *Chenopodium rubrum*, *Chenopodium salinum*, *Puccinellia nuttalliana*, *Salicornia europaea*, *Triglochin maritima*, and *Scirpus pungens*. On the upslope side the community grades into grasslands. This community is considered as common around sloughs and saline shores throughout the Grassland and Parkland regions (C. Wallis pers comm.). Because it is only documented once in the literature an assessment of variability between occurrences is not possible.

Triglochin maritima

Seaside arrow grass

Preliminary Rank: S2?

This community is described by Wallis (1990) as occurring generally within the Grassland and Parkland Regions of Alberta as a saline emergent marsh characteristic of very fine sand to sandy clay saline meadow soils. It is typified by a band of pure *Triglochin maritima* with a basal cover < 20%. The most prevalent associated plants include *Hordeum jubatum*, *Distichlis stricta*, *Ranunculus cymbalaria*, *Chenopodium rubrum* and *Suaeda calceoliformis*. This community also occurs in active springs in Alberta but is very restricted in area extent (C. Wallis pers comm.). The plant species *Triglochin maritima* is common in saline depressions in Alberta (Moss 1983, Johnson *et al.* 1995), British Columbia (Parish *et al.* 1996) and Saskatchewan (Dodd and Coupland 1966) and the community type is described from north-central United States (e.g. Montana; Heidel *et al.* 2000). This community is currently on the ANHIC plant community tracking list (Allen 2001) and ranked S2?.

Elsewhere, this community corresponds to the *T. maritimus* community described from Montana by Heidel *et al.* (2000); a community present on fine-textured soils of saline and subsaline lakes, seeps and associated inlets. However, a formal comparison of this community with Alberta Parkland types is difficult because cover estimates are not provided by Heidel *et al.* (2000).

Crosswalk Table – Literature Comparisons.

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Arctostaphylos uva-ursi</i>	<i>Arctostaphylos uva-ursi</i> (Douglas and Bliss 1997; Cascade Mountains of Washington State), <i>Arctostaphylos uva-ursi</i> (Johnson <i>et al.</i> 1995; Northern British Columbia, Yukon, and Alaska)	Similar communities are in forested and alpine habitats, different in elevation and substrate than the sand community described in the Central Parkland by Meijer and Karpuk (1999).
<i>Juniperus horizontalis</i>	<i>Juniperus horizontalis</i> (Johnson <i>et al.</i> 1995; Moss 1983; Alberta Foothills and Mountains)	Similar communities are also described from sandy and rocky areas in the foothills and mountains of Alberta but elevation and climate would differentiate these communities from Central Parkland types.
<i>Juniperus horizontalis</i> - <i>Selaginella densa</i> - <i>Calamovilfa longifolia</i>	<i>Juniperus horizontalis</i> / <i>Koeleria macrantha</i> - <i>Artemisia frigida</i> (Thorpe and Goodwin 1993; Manito Sand Hills, Saskatchewan)	Similar substrate, almost all associated species similar. Prevailing conditions also similar.
<i>Acer negundo</i> / <i>Prunus virginiana</i>	<i>Acer negundo</i> / <i>Prunus virginiana</i> (Wallis 1977, South east Alberta; Bourgeron and Engelking 1994, Great Plains of the United States; Carsey <i>et al.</i> 2001, Colorado)	Similar communities are found in riparian and alpine areas. Also found in Colorado in stream reaches on larger rivers in low elevation canyons in the foothills and plateaus (generally steep gradient and coarse soils). Thought to occur in coulee bottoms in the Central Parkland of Alberta.
<i>Alnus tenuifolia</i> - <i>Salix discolor</i> / <i>Lonicera involucrata</i> - <i>Cornus stolonifera</i> - <i>Ribes hudsonianum</i> / <i>Plagiomnium cuspidatum</i>	River Alder Gully: <i>Alnus tenuifolia</i> - <i>Cornus stolonifera</i> - <i>Ribes spp.</i> (Beckingham <i>et al.</i> 1996, Midboreal Ecoregion of Saskatchewan)	The midboreal Saskatchewan community has the same site conditions and species assemblage, but the alder is not of tree height.
<i>Betula neolaskana</i> - <i>Populus balsamifera</i> / <i>Cornus stolonifera</i> / <i>Calamagrostis canadensis</i>	Dogwood <i>Populus balsamifera</i> - <i>Populus tremuloides</i> (Beckingham <i>et al.</i> 1996; Midboreal Ecoregion of Saskatchewan)	Midboreal Sask. Community has a lower birch component and a higher trembling aspen component, but similar topographic position and nutrient regime.
<i>Populus balsamifera</i> / <i>Salix discolor</i> - (<i>Betula occidentalis</i>)	Not described elsewhere in literature	
<i>Populus balsamifera</i> / <i>Salix petiolaris</i>	<i>Populus Balsamifera</i> / <i>Salix spp.</i> community MBMA4 (Willoughby and Downing 1995, Boreal Region of Alberta)	Similar site characteristics and dominant species, associated species are variable.
<i>Populus balsamifera</i> / <i>Viburnum opulus</i> / <i>Cornus stolonifera</i> / <i>Aralia nudicaulis</i>	Not described elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Populus tremuloides</i>	<i>Populus tremuloides</i> (Fehr 1982, Alberta; Wroe 1971, Alberta; Looman 1987a, Alberta).	All communities mentioned are upland aspen forest in parkland with similar understory assemblages and site conditions.
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> - <i>Betula papyrifera</i> / <i>Symphoricarpos occidentalis</i>	Not described elsewhere in literature	
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> / <i>Amelanchier alnifolia</i>	Aspen Consociation (Moss 1932), low-bush cranberry Aw (Beckingham and Archibald 1996, Boreal Alberta)	This community is clearly representative of Moss' (1932) "aspen consociation". The community described by Beckingham and Archibald (1996) has a higher cover of <i>Rosa acicularis</i> (~10%) but a similar herb layer without dominants.
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> / <i>Cornus stolonifera</i> - <i>Rosa acicularis</i> - <i>Viburnum edule</i> / <i>Aralia nudicaulis</i>	Not described elsewhere in literature	
<i>Populus tremuloides</i> - <i>Populus balsamifera</i> / <i>Symphoricarpos occidentalis</i>	Not described elsewhere in literature	
<i>Populus tremuloides</i> / <i>Corylus cornuta</i> / Low Shrubs and Herbs	<i>Populus tremuloides</i> / <i>Corylus cornuta</i> (Colorado, North Dakota, South Dakota, Wyoming, Saskatchewan), <i>Populus tremuloides</i> / <i>Corylus cornuta</i> / <i>Aster conspicuus</i> - <i>Disporum trachycarpum</i> - <i>Epilobium angustifolium</i> (Bork 1993, Central Alberta), <i>Populus tremuloides</i> / <i>Corylus cornuta</i> / <i>Oryzopsis asperifolia</i> (Bork 1993, Central Alberta), dogwood Pb-Aw, low-bush cranberry Aw (Beckingham and Archibald 1996, Boreal Alberta), Aspen - Beaked Hazelnut - Rose / Wild Peavine (community LBMIB3, Willoughby and Downing 1995, Mixedwood Boreal Alberta) and Aspen - Beaked Hazelnut - Rose / Wild Sarsaparilla (community LBMIB4, Willoughby and Downing 1995, Mixedwood Boreal Alberta); Aw / Rose - Hazelnut (Willoughby <i>et al.</i> 2000, Alberta Central Mixedwood).	All communities have similar site and dominant species characteristics. Understory cover varies as it is described in the <i>P. tremuloides</i> section of this report. Moss (1932) included this community type his "aspen consociation", and Looman (1987a) in his "Alliance <i>Populion tremuloides</i> in the association <i>Corylo-Populetum</i> (forests dominated with aspen having <i>Corylus</i> prominent in the shrub layer)."

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Populus tremuloides</i> / <i>Prunus virginiana</i> - (Shrub) / Low Herb	<i>Populus tremuloides</i> / <i>Populus tremuloides</i> - <i>Prunus virginiana</i> - <i>Amelanchier alnifolia</i> / <i>Juniperus horizontalis</i> - <i>Arctostaphylos uva-ursi</i> (Meijer and Karpuk 1999, Alberta), <i>Populus tremuloides</i> / <i>Rosa acicularis</i> - <i>Prunus virginiana</i> / <i>Carex siccata</i> (Fehr 1984, Alberta), <i>Populus tremuloides</i> / <i>Prunus virginiana</i> (Natureserve 2001, North Dakota); <i>Populus tremuloides</i> / <i>Amelanchier alnifolia</i> - <i>Symphoricarpos oreophilus</i> / <i>Calamagrostis rubescens</i> (Meuggler 1988; Meuggler and Campbell 1982, Idaho, Utah); <i>Populus tremuloides</i> / <i>Amelanchier alnifolia</i> - <i>Prunus virginiana</i> / <i>Symphoricarpos occidentalis</i>	These communities have similar soil and site conditions as the types described under the <i>Populus tremuloides</i> section of this report. However, understory diversity can vary and can include <i>Thalictrum venulosum</i> , <i>Campanula rotundifolia</i> , <i>Lathyrus ochroleucus</i> , and <i>Schizachne purpurascens</i> . In North Dakota, stands occur on well-drained loam soils. The canopy is moderately dense to dense. It is dominated by <i>Populus tremuloides</i> , with <i>Fraxinus pennsylvanica</i> as a common associate.
<i>Populus tremuloides</i> / <i>Prunus virginiana</i> - <i>Amelanchier alnifolia</i> - <i>Rosa acicularis</i> / <i>Gallium boreale</i>	<i>Populus tremuloides</i> / <i>Prunus virginiana</i> (Natureserve 2001; Girard et al. 1989; Schneider et al. 1997, Montana, South Dakota, Wyoming), <i>Populus tremuloides</i> / <i>Prunus virginiana</i> - <i>Rosa acicularis</i> (Fehr 1984, Alberta), <i>Populus tremuloides</i> / <i>Prunus virginiana</i> / <i>Corylus cornuta</i> - <i>Rosa acicularis</i> / <i>Symphoricarpos albus</i> (Miejer and Karpuk 1999, eastern Alberta).	All communities have similar site and dominant species characteristics. Understory cover varies as it is described in the community description section of this report. These community types are described under the <i>Populus tremuloides</i> Forest/woodland section.
<i>Populus tremuloides</i> / <i>Prunus virginiana</i> / <i>Solidago canadensis</i>	Not described elsewhere in literature	
<i>Populus tremuloides</i> / <i>Rosa acicularis</i> - <i>Rubus idaeus</i>	Aspen - Rose (Community LBMA3, Willoughby and Downing 1995, Boreal Alberta); Aw / Rose / Tall Forb (Willoughby et al. 2000, Alberta Central Mixedwood)	Communities have similar dominants, but detail is not sufficient for an adequate comparison of herb and shrub layer similarities.
<i>Populus tremuloides</i> / <i>Symphoricarpos occidentalis</i> / Herbs	<i>Populus tremuloides</i> / <i>Symphoricarpos occidentalis</i> (Wroe 1971, Alberta); <i>Populus tremuloides</i> / <i>Symphoricarpos occidentalis</i> / <i>Gallium boreale</i> (Smith and Kondla 1972, Alberta); <i>Populus tremuloides</i> / <i>Symphoricarpos occidentalis</i> / <i>Rubus pubescens</i> (Hilton 1970, Alberta).	All communities have similar site conditions and associated species. All are described from the Central Parkland Natural Subregion near Dry Island Buffalo Jump Provincial Park.
<i>Populus tremuloides</i> / <i>Viola adunca</i> - <i>Orthilia secunda</i>	Not described elsewhere in literature	
<i>Picea glauca</i> / <i>Cornus stolonifera</i> / <i>Climacium dendroides</i>	Not described elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Picea mariana</i> / <i>Ledum groenlandicum</i> - <i>Vaccinium vitis-idaea</i>	Shrubby Bog and Treed Bog: <i>Picea mariana</i> / <i>Ledum groenlandicum</i> / <i>Sphagnum</i> spp. (Beckingham et al. 1996, Midboreal Saskatchewan); <i>Picea mariana</i> / <i>Ledum groenlandicum</i> / <i>Pleurozium schreberi</i> (Griffiths et al. 1997, Alberta)	All have similar site characteristics and associated species. No differences.
<i>Pinus banksiana</i> / <i>Arctostaphylos uva-ursi</i>	<i>Pinus banksiana</i> (Johnson et al. 1995, Boreal Alberta, Sask., Manitoba), bearberry Pj (Beckingham and Archibald 1996, Boreal Alberta)	Sandy upland jack pine forest with similar understory and site conditions. Community described by Beckingham and Archibald (1996) has a higher cover of <i>Cladina</i> spp, <i>Vaccinium myrtilloides</i> , and <i>Alnus crispa</i> .
<i>Picea glauca</i> - <i>Populus tremulooides</i> - <i>Populus balsamifera</i>	Horsetail Pb-Sw (Beckingham and Archibald 1996)	Similar dominants but there is variation in understory species in both composition and cover (descriptions not adequate for formal comparison).
<i>Picea glauca</i> - <i>Populus tremulooides</i> / low shrub	<i>Picea glauca</i> - <i>Populus tremulooides</i> (Greenall 1996, Manitoba); Low Bush Cranberry Trembling Aspen - White Spruce (Beckingham et al. 1996, Midboreal Saskatchewan)	Similar forest - mature spruce with aspen understory, similar site conditions and successional stage, shrubs and herbs vary little between communities.
<i>Populus tremulooides</i> - <i>Populus balsamifera</i> - <i>Picea glauca</i> / <i>Rubus idaeus</i>	Aspen--balsam poplar / rose--raspberry / pink wintergreen, and aspen / red-osier dogwood (Beckingham 1993, Alberta).	These communities differ in their white spruce content and understory compositions (species can vary in type and abundance), but site conditions are similar.
<i>Populus tremulooides</i> (<i>Picea glauca</i>) / <i>Aralia nudicaulis</i> / <i>Cornus canadensis</i>	Not found elsewhere in literature	
<i>Senecio congestus</i>	Not found elsewhere in literature	
<i>Eleocharis quinquefolia</i>	<i>Eleocharis quinquefolia</i> (Heidel et al. 2000, Montana, North and South Dakota), <i>Eleocharis quinquefolia</i> (Carsey et al. 2001, Colorado)	The fens of the northern Central Parkland and the communities described from the states of Montana and the Dakotas differ in site conditions, climate, and in associated species. All communities are considered marl, rich fens in localized areas where mineral-rich groundwater flow emerges at the ground's surface.
<i>Agropyron trachycaulum</i> - <i>Hordeum jubatum</i> - <i>Elymus canadense</i>	Not found elsewhere in literature	
<i>Agrostis scabra</i> - <i>Achillea millefolium</i> - <i>Antennaria nitida</i>	Not found elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Bouteloua gracilis</i> - <i>Artemisia frigida</i>	<i>Bouteloua gracilis</i> (Biota Consultants, unpublished data)	Biota Consultants (unpublished data) found <i>B. gracilis</i> forming communities each with the following dominants: <i>Stipa comata</i> , <i>Koeleria macrantha</i> , <i>Festuca hallii</i> , <i>Agropyron smithii</i> , <i>Carex stenophylla</i> , and <i>Carex pensylvanica</i> . These communities were primarily grassland types, but could also contain a patchy, low shrub component consisting of either <i>Rosa woodsii</i> , <i>Symphoricarpos occidentalis</i> , or <i>Elaeagnus commutata</i> .
<i>Calamagrostis canadensis</i>	<i>Calamagrostis canadensis</i> (Schneider <i>et al.</i> 1997; Great Plains of the United States); <i>Calamagrostis canadensis</i> (Natureserve 2001, Colorado); <i>Calamagrostis canadensis</i> (Willoughby <i>et al.</i> 2000, Central Mixedwood Alberta), <i>Calamagrostis canadensis</i> (Carsey <i>et al.</i> 2001, Colorado).	<i>Calamagrostis canadensis</i> communities are described from the grasslands of the United States but the habitat conditions are not specified), so it's unclear as to how they would differ from communities described from Alberta in seasonally-inundated depressions, along the edges of water bodies, and in forest gaps. However, Alberta communities are similar to those described from Colorado by Carsey <i>et al.</i> 2001.
<i>Calamagrostis inexpansa</i> - <i>Juncus balticus</i>	Not found elsewhere in literature	
<i>Calamagrostis stricta</i> - <i>Carex atherodes</i> - <i>Phalaris arundinacea</i>	Not found elsewhere in literature	
<i>Calamagrostis stricta</i> - <i>Juncus balticus</i> - <i>Potentilla anserina</i> - <i>Aster hesperius</i>	Not found elsewhere in literature	
<i>Calamovilfa longifolia</i> - <i>Artemisia frigida</i> - <i>Koeleria macrantha</i> - <i>Festuca saximontana</i> - <i>F. hallii</i> / <i>Selaginella densa</i>	Not found elsewhere in literature	
<i>Carex atherodes</i> - <i>Calamagrostis inexpansa</i>	Not found elsewhere in literature	Seasonally flooded (differs from other <i>C. atherodes</i> communities in hydrological regime and associated species).
<i>Carex atherodes</i> - <i>Carex utriculata</i>	Not found elsewhere in literature	Intermittently flooded (differs from other <i>C. atherodes</i> communities in hydrological regime and associated species).
<i>Carex atherodes</i> - <i>Scolochloa festucacea</i> - <i>Eleocharis palustris</i> - <i>Typha latifolia</i>	Not found elsewhere in literature	Permanently flooded (differs from other <i>C. atherodes</i> communities in hydrological regime and associated species).
<i>Carex atherodes</i> - <i>Scolochloa festucacea</i> - <i>Poa palustris</i> - <i>Cirsium arvense</i>	Not found elsewhere in literature	Seasonally flooded (differs from other <i>C. atherodes</i> communities in hydrological regime and associated species).

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Carex atherodes</i> / <i>Lemna minor</i>	<i>Carex atherodes</i> (Natureserve 2001, Schneider <i>et al.</i> 1998, Great Plains of the United States)	This temporarily flooded awned sedge wet meadow occurs in the northern tallgrass prairie region of the United States and Canada. Stands occur on lowland sites that have standing water for several weeks each year. These sites are typically in depressions or basins but can be along streams and rivers. The water may be fresh or moderately saline. <i>Carex atherodes</i> may form essentially monotypic stands or just be the dominant species. Common associated species include <i>Alisma triviale</i> , <i>Symphotrichum lanceolatum</i> (= <i>Aster lanceolatus</i>), <i>Eleocharis palustris</i> , <i>Glyceria grandis</i> (in drier stands), <i>Mentha arvensis</i> , <i>Phalaris arundinacea</i> , <i>Polygonum amphibium</i> , <i>Scolochloa festucacea</i> , <i>Sium suave</i> , and <i>Sparganium eurycarpum</i> . Alberta communities can become pure stands of <i>Carex atherodes</i> .
<i>Carex</i> group - <i>Calamovilfa longifolia</i>	Not found elsewhere in literature	
<i>Carex</i> group - <i>Koeleria macrantha</i>	Not found elsewhere in literature	
<i>Carex</i> group - <i>Stipa curtiseta</i>	Not found elsewhere in literature	
<i>Carex pseudo-cyperus</i> - <i>Calla palustris</i>	<i>Carex pseudo-cyperus</i> - <i>Calla palustris</i> (Moss 1928, Alberta); <i>Carex pseudo-cyperus</i> - <i>Calla palustris</i> (G. Griffiths and L. Allen, pers comm., Alberta).	Both communities were found along water body edges, Moss' communities were found adjoining two lakes. Communities are also found near Elk Island National Park but site characteristics cannot be adequately compared (lack of detail in Moss 1928).
<i>Carex siccata</i> - <i>Calamovilfa longifolia</i>	Not found elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Distichlis stricta</i>	<i>Distichlis stricta</i> (Schneider <i>et al.</i> 1997, Great Plains of the United States); <i>Distichlis stricta</i> (Dodd and Coupland 1966, Saskatchewan)	Salt grass is a dominant species of saline areas as far south as Oklahoma (Dodd and Coupland 1966). Salt grass occupies much less extensive areas in the warm desert region south of Utah, but it is prominent in the most saline parts of desert meadows (Dodd and Coupland 1966). Dodd and Coupland (1966) noted this community occurring on a variety of soil textures and moisture levels. Wallis (1990) described this community as occupying areas which are less saline than areas occupied by <i>Puccinellia nuttalliana</i> , forming almost pure stands with a basal cover of > 20%.
<i>Distichlis stricta - Agropyron smithii</i>	<i>Distichlis stricta - Agropyron spp.</i> (Dodd and Coupland 1966, Saskatchewan)	Same semi-halophytic or halophytic site conditions. Saskatchewan community varies in terms of basal coverage ranging from 10-30% of <i>Distichlis</i> cover, with significantly higher graminoid cover of associated species (up to 100% basal coverage).
<i>Eleocharis palustris</i>	<i>Eleocharis palustris</i> (Schneider <i>et al.</i> 1997, Great Plains United States); <i>Eleocharis palustris</i> (Heidel <i>et al.</i> 2000, Montana);	Site conditions not described in detail for an adequate comparison in Schneider <i>et al.</i> (1998). In other instances this community occurs in slightly to moderately brackish wetlands, including prairie potholes and riparian settings that are seasonally flooded. It is well represented in the central and northern Great Plains, possibly extending into the Southwest and the Pacific Northwest states, and is a major type at low elevations throughout Montana (Heidel 2000, Hansen <i>et al.</i> 1995).
<i>Hordeum jubatum</i>	<i>Hordeum jubatum</i> (Dodd and Coupland 1966, Saskatchewan); <i>Hordeum jubatum</i> (Natureserve 2001, Great Plains of the United States)	Similar communities: The foxtail barley herbaceous vegetation is a seasonally flooded subsaline wetland type (Thompson 1994), and a drawdown phase of seasonal and semipermanent brackish and subsaline marshes in Montana (Heidel <i>et al.</i> 2000). In Saskatchewan it occurs as an almost pure stand over extensive portions of large, flat saline depressions, or as a ring around smaller depressions.

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Hordeum jubatum</i> - <i>Puccinellia nuttalliana</i>	<i>Hordeum jubatum</i> - <i>Puccinellia nuttalliana</i> (Walker and Coupland 1970, Saskatchewan)	Hordeum jubatum occurs in variable composition with <i>Puccinellia nuttalliana</i> , rarely as a dominant (as found in Alberta) around sloughs in Saskatchewan.
<i>Koeleria macrantha</i> - <i>Agropyron smithii</i>	<i>Koeleria macrantha</i> - <i>Agropyron smithii</i> (Wroe 1971, Alberta); <i>Koeleria macrantha</i> - <i>Agropyron smithii</i> (Fehr 1982, Alberta)	These two communities appear very similar; same soil type, same associated species, same hydrological regime, and same aspect. Both are located within the central parkland subregion of Alberta.
<i>Muhlenbergia asperifolia</i> - <i>Scirpus nevadensis</i> - <i>Distichlis stricta</i>	Not found elsewhere in literature	
<i>Puccinellia nuttalliana</i>	<i>Puccinellia nuttalliana</i> (Fehr 1982, Alberta); <i>Puccinellia nuttalliana</i> (Wallis 1990, Alberta); <i>Puccinellia nuttalliana</i> (Heidel 2000, Montana, Utah, California).	For both descriptions in Alberta this community is found on fine, sandy loam to heavy, clay saline meadow soils, and subhydic saline depressions, where it occurs in a wide band around saline sloughs. In Montana it is associated with seasonally saturated saline soils in wet meadows, saline seeps and the flats associated with saline alkali lakes. Utah and California communities are not described in detail for adequate comparisons.
<i>Puccinellia nuttalliana</i> - <i>Distichlis stricta</i>	<i>Puccinellia nuttalliana</i> - <i>Distichlis stricta</i> (Parish et al. 1996, Interior British Columbia).	Described from low to mid elevations in alkali areas in interior BC. Probably similar site and species composition but with differences in elevation and climate compared to Alberta communities (detail not sufficient for formal comparison).
<i>Ruppia maritima</i>	<i>Ruppia maritima</i> (Wallis 1990, Alberta); <i>Ruppia maritima</i> (Heidel et al. 2000, Montana)	Wallis (1990) mentions this community for Alberta on unvegetated shores, where rooted vegetation is absent, and salinity is between 3 and 50 parts per million. This community type occurs in the glaciated plains region of Alberta, Saskatchewan, Montana and North Dakota. It is typically in closed-basin watersheds, and often part of glacial outwash channels (Heidel et al. 2000).
<i>Scirpus acutus</i>	<i>Scirpus acutus</i> (Wallis, Alberta); <i>Scirpus acutus</i> (Meijer and Karpuk 1999, Alberta); <i>Scirpus acutus</i> - <i>Scirpus validus</i> (Griffiths et al. 1997, Alberta); <i>Scirpus acutus</i> (Thompson 1994, Montana); <i>Scirpus acutus</i> (Heidel et al. 2000, Montana).	There is potential for soil and moisture conditions to vary between these community types. Descriptions are not detailed enough for formal comparisons.

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Scirpus paludosus</i>	<i>Scirpus paludosus</i> (Wallis 1990, Alberta); <i>Scirpus paludosus</i> (Fehr 1982, Alberta).	Both communities are described as a wide band around saline sloughs, but other characteristics are not described in detail for a formal comparison. These are probably the same community type.
<i>Scirpus paludosus</i> - <i>Puccinellia nuttalliana</i> - <i>Hordeum jubatum</i> - <i>Distichlis stricta</i>	Not described elsewhere in literature	
<i>Scirpus pungens</i>	<i>Scirpus pungens</i> (Heidel <i>et al.</i> 2000, Montana)	Not well described from Alberta, but occurs as bands around saline sloughs in Montana in the Intermountain Basin and the western and northern Great Plains of the United States.
<i>Scirpus pungens</i> - <i>Hordeum jubatum</i> - <i>Puccinellia nuttalliana</i>	Not described elsewhere in literature	
<i>Spartina gracilis</i> - <i>Agropyron smithii</i>	Not described elsewhere in literature	
<i>Stipa comata</i> - <i>Koeleria macrantha</i> - <i>Agropyron smithii</i>	Not described elsewhere in literature	
<i>Stipa comata</i> - <i>Koeleria macrantha</i> - <i>Agropyron trachycaulum</i> - <i>Agropyron smithii</i> - <i>Carex obtusata</i>	Not described elsewhere in literature	
<i>Stipa comata</i> - <i>Poa sandbergii</i> - <i>Carex obtusata</i>	Not described elsewhere in literature	
<i>Stipa comata</i> - <i>Stipa viridula</i> - <i>Festuca hallii</i>	Not described elsewhere in literature	
<i>Stipa curtiseta</i> - <i>Artemisia frigida</i>	<i>Stipa curtiseta</i> - <i>Artemisia frigida</i> (Wroe 1971 and Fehr 1982, Alberta); <i>Stipa curtiseta</i> - <i>Agropyron trachycaulum</i> - <i>Artemisia frigida</i> (Anderson 1972, Alberta)	All communities have similar topographic positions and hydrological regimes. Variation in species associations can occur with <i>Koeleria cristata</i> , <i>Artemisia frigida</i> , and <i>Anemone patens</i> var. <i>wolfgangiana</i> . Grazing pressure can vary in the community described by Anderson (1972).
<i>Stipa curtiseta</i> - <i>Bouteloua gracilis</i>	Not found elsewhere in literature	
<i>Stipa curtiseta</i> - <i>Festuca hallii</i>	<i>Stipa curtiseta</i> - <i>Festuca hallii</i> (Fehr 1982, Alberta); <i>Stipa curtiseta</i> - <i>Festuca hallii</i> (Wroe 1971, Alberta); <i>Stipa spp.</i> - <i>Festuca spp.</i> (Vujanovic and Wein 1997, Alberta).	The communities described by Fehr (1982) and Wroe (1971) are essentially identical. Vujanovic and Wein (1998) do not provide species information for a formal comparison, so this community might be completely different from theirs.
<i>Stipa curtiseta</i> - <i>Koeleria macrantha</i> - <i>Agropyron trachycaulum</i> / <i>Cladina mitis</i>	Not found elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Stipa viridula</i> - <i>Poa sandbergii</i> - <i>Carex obtusata</i>	Not found elsewhere in literature	
<i>Typha latifolia</i>	Marsh: <i>Typha latifolia</i> / <i>Carex</i> spp. (Beckingham et al. 1996, Midboreal Ecoregion of Saskatchewan); <i>Carex aquatilis</i> - <i>Carex rostrata</i> / <i>Typha latifolia</i> (Thorpe and Goodwin 1993, Manito Sand Hills Saskatchewan); <i>Typha latifolia</i> (Griffiths et al. 1995, Alberta); <i>Typha latifolia</i> (Heidel et al. 2000, Montana).	Site conditions are similar in all communities, but associated species can vary. Griffiths et al. (1995) describe a community with varying amounts of <i>Lemna trisulca</i> . Thorpe and Goodwin (1993) describe a community with co-dominants of <i>Carex</i> spp., as does Beckingham et al. (1996).
<i>Antennaria parvifolia</i> (<i>nitida</i>) - <i>Artemisia frigida</i> - <i>Koeleria macrantha</i>	Not found elsewhere in literature	
<i>Carex aquatilis</i> - <i>Carex utriculata</i>	<i>Carex aquatilis</i> (Schneider et al. 1997, Great Plains United States); <i>Carex aquatilis</i> - <i>Carex utriculata</i> (Naturreserve 2001, Colorado)	In Alberta this community occurs in wet, morainal depressions and along the shores of lakes and ponds representing the interface between water sedge fen and aquatic, open water vegetation. In Colorado, this common wetland vegetation generally occurs in small to moderate-sized patches in very shallow, slow-moving to still water, or on saturated soils near low-order streams, lakes, and backwater areas of larger rivers where it is recognized by the presence of both <i>Carex aquatilis</i> and <i>Carex utriculata</i> in roughly equal proportions in the herbaceous layer.
<i>Elymus canadensis</i> - <i>Poa pratensis</i> - <i>Glycyrrhiza lepidota</i>	Not found elsewhere in literature	
<i>Polygonum amphibium</i>	<i>Polygonum amphibium</i> (Kunze 1994, Washington State); <i>Polygonum amphibium</i> (Johnson 1939, 1941, Colorado); <i>Polygonum amphibium</i> (Heidel et al. 2000, Montana).	In Colorado, <i>Polygonum amphibium</i> often forms dense, nearly monotypic stands. <i>Lemna minor</i> , <i>Potamogeton natans</i> , <i>Sagittaria</i> spp., <i>Spirodela polyrrhiza</i> , and <i>Wolffia</i> spp. are occasionally present. Johnson (1939, 1941) found this community in Colorado montane lakes with significant sediment deposition at a higher elevation than Alberta types (and with differing, adjacent, montane community types). Heidel et al. (2000) describe a similar community to Alberta's occurring in brackish, seasonally inundated wetlands with recent disturbance.
<i>Potentilla anserina</i> - <i>Distichlis stricta</i>	Not found elsewhere in literature	
<i>Sparganium eurycarpum</i>	Not found elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Betula pumila</i> / <i>Carex aquatilis</i>	<i>Shrubby rich fen: Betula pumila</i> - <i>Carex</i> spp. (Beckingham <i>et al.</i> 1996, Midboreal Saskatchewan); <i>Betula pumila</i> - <i>Salix</i> spp. (Schneider <i>et al.</i> 1997, Natureserve 2001, Dakotas, Minnesota); <i>Betula pumila</i> - <i>Salix</i> spp. / <i>Carex</i> spp. (Beckingham <i>et al.</i> 1996, Archibald <i>et al.</i> 1996, West-central Alberta, South-western AB).	All communities potentially differ in the species of <i>Carex</i> present. This is not always identified to species, therefore, a formal comparison is not possible. Communities in the Dakotas and Minnesota have significantly less open water cover than those in Alberta and elsewhere.
<i>Elaeagnus commutata</i>	<i>Elaeagnus commutata</i> (Meijer and Karpuk 1999, Alberta); <i>Elaeagnus commutata</i> - <i>Symphoricarpos occidentalis</i> (Fehr 1982, Alberta); <i>Elaeagnus commutata</i> - <i>Symphoricarpos occidentalis</i> (Bradley and Bradley 1977, Alberta); <i>Elaeagnus commutata shrubland</i> (Heidel <i>et al.</i> 2000, Natureserve 2001, Montana).	Meijer and Karpuk (1997) described Silverberry communities from Dillberry Lake Provincial Park as low shrublands and shrubby meadows along the perimeter of saline lakes adjacent to marshes or graminoid meadows. Fehr (1982) and Bradley and Bradley (1972) describe an <i>Elaeagnus commutata</i> - <i>Symphoricarpos occidentalis</i> community occurring in middle and lower positions of mesic to submesic slopes, typically well developed and dense with a poorly developed herb-dwarf shrub layer. <i>Elaeagnus commutata</i> Shrubland is a minor type forming narrow stringers on stream terraces in northern Montana, but it is unclear if this is the same community as found in Alberta.
<i>Elaeagnus commutata</i> / <i>Symphoricarpos occidentalis</i> - <i>Rosa woodsii</i> / <i>Poa patustris</i> <i>Populus tremuloides</i> - <i>Amelanchier alnifolia</i> - <i>Prunus virginiana</i> / <i>Rosa</i> spp.	Not found elsewhere in literature <i>Populus tremuloides</i> / <i>Amelanchier alnifolia</i> / Tall Forbs Forest (Natureserve 2001; Meuggler 1988, Idaho, Nevada, Utah); <i>Populus tremuloides</i> / <i>Amelanchier alnifolia</i> (Meuggler and Campbell 1982, Colorado, Montana, Wyoming, Ontario); Aw / Saskatoon - Rose (Willoughby <i>et al.</i> 2000, Alberta Central Mixedwood)	All mature forest with varying understory. Descriptions are not detailed enough for a formal comparison.

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Populus tremuloides</i> / <i>Juniperus horizontalis</i> / <i>Cladina mitis</i>	<i>Populus tremuloides</i> / <i>Juniperus horizontalis</i> / <i>Cladina mitis</i> (Fehr 1984, Alberta); <i>Populus tremuloides</i> / <i>Juniperus horizontalis</i> / <i>Cladina mitis</i> (Meijer and Karpuk 1999, Renewable Resources Consulting 1974, Alberta).	This community is found in moderately well to rapidly drained inter-dune depressions, on Orthic Regisol soils with xeric moisture regimes (Fehr 1984). Meijer and Karpuk (1999) and Renewable Resources Consulting (1974) found this community on the upper positions of shallow, south-facing slopes in areas with rapid drainage, Orthic Dark Brown soils, xeric moisture regimes, and submesotrophic nutrient conditions. Forb diversity and cover is limited.
<i>Prunus virginiana</i> - <i>Amelanchier alnifolia</i> / <i>Agropyron subsecundum</i> - <i>Poa pratensis</i>	Not found elsewhere in literature	
<i>Rosa acicularis</i>	Not found elsewhere in literature	
<i>Rosa acicularis</i> - <i>Rubus idaeus</i> / <i>Lathyrus venosus</i>	Not found elsewhere in literature	
<i>Salix petiolaris</i>	<i>Salix petiolaris</i> (Fehr 1982, Alberta); <i>Salix petiolaris</i> / <i>Rosa woodsii</i> / <i>Sonchus arvensis</i> (Wroe 1971, Alberta).	Fehr (1982) describes this community as occurring in a mesic to hydric moisture regime in depressions with moderately to poorly drained, Rego and Orthic Humic Gleysol soils. Wroe (1971) describes a <i>Salix petiolaris</i> / <i>Rosa woodsii</i> / <i>Sonchus arvensis</i> community from the central parkland forming rings around depressions, occurring as clumps around depressions, and restricted generally to moist areas with rolling topography. Soils are Rego-humic Gleysols, both carbonated and salinized.
<i>Salix petiolaris</i> - <i>Salix planifolia</i> / <i>Carex utriculata</i>	Not found elsewhere in literature	
<i>Salix petiolaris</i> / <i>Juncus balticus</i>	Not found elsewhere in literature	
<i>Salix pyrifolia</i> - <i>Betula neoalaskana</i> / <i>Ledum groenlandicum</i> - <i>Calamagrostis canadensis</i> / <i>Sphagnum warnstorffi</i>	Not found elsewhere in literature	
<i>Salix pyrifolia</i> / <i>Carex utriculata</i>	Not found elsewhere in literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Symphoricarpos occidentalis</i>	<i>Symphoricarpos occidentalis</i> (Anderson 1972, Alberta); <i>Symphoricarpos occidentalis</i> (Fehr 1982, Alberta); <i>Symphoricarpos occidentalis</i> (Wroe 1971, Alberta); <i>Symphoricarpos occidentalis</i> (Meijer and Karpuk 1999, Alberta); <i>Symphoricarpos occidentalis</i> (Heidel <i>et al.</i> 2000, Montana).	Anderson (1972) describes this community as occupying the shallow depressions on the grassland and in many instances positions of higher elevations adjacent to the aspen communities. A large quantity of dead stems of <i>Symphoricarpos occidentalis</i> can be an important component of this community. Meijer and Karpuk (1999) describe this community associated with dry conditions, Orthic and Rego Dark Brown Chernozemic soils, and found on north-east and south-west-facing slopes in rapidly drained areas with submesotrophic conditions along the perimeters of naturally open woodlands. This community is also reported from Saskatchewan, Missouri, and Montana found in mesic depressions, typically surrounded by upland grassland associations (Heidel <i>et al.</i> 2000). A buckbrush shrub community type has been described for Montana, North Dakota, and Alberta (Anderson <i>et al.</i> 1980, Hansen <i>et al.</i> 1984, Hansen <i>et al.</i> 1990). Buckbrush often forms dense stands with little understory.
<i>Symphoricarpos occidentalis</i> - <i>Rosa arkansana</i> / <i>Juniperus horizontalis</i>	Not found elsewhere in Literature	
<i>Symphoricarpos occidentalis</i> / <i>Antennaria parvifolia</i> (nitida) - <i>Festuca hallii</i>	Not found elsewhere in Literature	
<i>Populus tremuloides</i> / <i>Juniperus horizontalis</i>	Not found elsewhere in Literature	
<i>Juniperus horizontalis</i> / (<i>Koeleria macrantha</i>) / <i>Cladina mitis</i>	Not found elsewhere in Literature	
<i>Calamovilfa longifolia</i> - <i>Sporobolus cryptandrus</i> - <i>Koeleria macrantha</i> - <i>Carex obtusata</i>	Not found elsewhere in Literature	
<i>Salicornia europaea</i>	Not found elsewhere in Literature	

Plant Communities of Alberta's Central Parkland

Central Parkland Community	Similar Community (Citation and Location)	Comments
<i>Scirpus nevadensis</i>	<i>Scirpus nevadensis</i> (Carsey <i>et al.</i> 2001, Colorado)	Species diversity in Colorado sites is higher. <i>Scirpus nevadensis</i> provides 10 to 80% cover; other species include <i>Juncus balticus</i> var. <i>montanus</i> (mountain rush, up to 5%), <i>Spartina gracilis</i> (alkali cordgrass, up to 10%), <i>Scirpus pungens</i> (common threesquare <5%), and <i>Hordeum jubatum</i> (<10%). Forb cover is minimal.
<i>Scirpus nevadensis</i> - (<i>Triglochin maritima</i>)	Not found elsewhere in Literature	
<i>Suaeda calceoliformis</i>	Not found elsewhere in Literature	
<i>Triglochin maritima</i>	<i>Triglochin maritima</i> – <i>Triglochin palustre</i> (Carsey <i>et al.</i> 2001, Colorado); <i>Triglochin maritima</i> (Dodd and Coupland 1966, Saskatchewan)	Documented from high-elevation intermountain parks in Colorado where soils are poorly drained, deep, saline and alkaline. Dodd and Coupland (1966) describe this community in Saskatchewan on coarse-textured (very fine sandy to sandy clay) saline meadow soils near the center of saline depressions. They noted that this species borders areas devoid of vegetation in the lowest part of the depression
<i>Populus tremuloides</i> - <i>Prunus virginiana</i> / <i>Juniperus horizontalis</i> / <i>Selaginella densa</i>	Not found elsewhere in Literature	

Discussion

The identification of community types in the Central Parkland Subregion of Alberta has been done primarily through qualitative, descriptive means, and with this type of information it is difficult to determine abundance or ecological effectiveness of described community types. In the literature, community descriptions vary from superficial to detailed, and in most instances there is not enough information to adequately describe community structure, let alone estimate a conservation rank (i.e. distribution and abundance). Several communities found in the Central Parkland Subregion can also be found in higher abundance in grassland, boreal or foothill regions. On a provincial scale, these would be considered without threat, but between natural regions the same community is most likely subject to different microclimates, animal assemblages (e.g. herbivory) and human use, hence serving two different functions on the landscape. It would be a difficult task to determine ecological differences between the same community type across natural regions, but it should be noted that some S5 communities described from the Central Parkland could be very different (through area, edge and species use effects), and serve very different ecological functions relative to similar community types found in other natural regions.

An estimation of patch size to augment measures of occurrence seems critical for an area as fragmented as the central parkland subregion of Alberta. Certainly an aspen forest relegated to marginal substrate or completely fragmented as a fencerow is not comparable to a similar forest type further north. With current information, a rating of S5 for an aspen community described from the Central Parkland Subregion says little about its value as habitat for plant and animal species endemic or specific to this subregion. We found only one source that gave quantitative values on patch size for forest patches in the Central Parkland. Given that the ranking guidelines are based primarily on the number of occurrences, an estimation of patch size, complete with some estimate of variance, seems critical for this process. There is a large body of information on area and edge effects for both plants and animals, something that should not be ignored when considering community "occurrences". These types of spatial data are readily available for other areas of the province and similar techniques should be used in the Central Parkland (e.g. simple GIS AVI polygon mapping).

Community description styles vary considerably between authors. This is to be expected considering the available information spans almost 70 years, and there has not been a standardized sampling method for either qualitative or empirical data collection. Standardized methods for describing communities are now prevalent (see Grossman *et al.* 1998), and well-described communities from historical data sets should be revisited in the field to apply these standards. Communities can appear similar in superficial descriptions, but differ markedly in soil type, aspect, and moisture regime for example, information that is currently not readily available in most cases. We caution the reader against information gaps such as these when making conclusions regarding community similarities outlined throughout this report.

The Central Parkland Natural Subregion of Alberta is the most impacted area of the province, and native plant communities exist only in remnants. In fact, upwards of 95% of the area is cultivated and converted for human agricultural, urban and industrial use. It would appear that a vegetation survey and digital mapping exercise locating native plant community remnants would not be a large undertaking across 5% of the region, even if these were evenly or randomly distributed. The amount of change on this landscape, the change in agricultural land boundaries, gas pipelines, highways and new urban developments makes the use and reliability of historical plant community descriptions tenuous. This report makes extensive use of historical community accounts, some dating back to the early 1900's. Our effort here represents a preliminary estimation of what is potentially still in the Central Parkland and, as it is based on existing literature, is most certainly not complete or necessarily representative of the current state of Central Parkland plant community diversity.

Not all formation classes are represented in this report. We found information on Forest/Woodland, Shrubland, Dwarf-shrubland, Herbaceous, and Sparse vegetation; we did not find information on nonvascular communities. Lichens and particularly mosses form important components of vascular-dominated communities in this region and undoubtedly form independent communities that would be classified into the nonvascular formation class; particularly in areas of the Central Parkland where exposed rock or glacial erratics are common (e.g. kame moraines). These community types should be considered in future surveys of this area.

Personal Communications

Lorna Allen, Biodiversity Biologist, Alberta Community Development, Parks and Protected Areas, Operations and Resource Management Branch, Government of Alberta, Edmonton Alberta.

Dr. Graham Griffiths, Biologist, Sherwood Park, Alberta.

Harry Loonen, Public Lands Specialist, Wainwright Public Lands Division, Sustainable Resource Development, Government of Alberta, Wainwright, Alberta.

Dr. Cliff Wallis, Biologist, Cottonwood Consultants, Calgary, Alberta.

Anne Holcroft Weerstra, Biologist, Biota Consultants, Cochrane, Alberta.

Literature Cited

Acharya, S.N., C.B. Chu, and R.Hermesh. 1989. Effects of population, environment and their interaction on Saskatoon berry (*Amelanchier alnifolia* Nutt.) seed germination. *Can. J. Bot.* 62:858-864.

Acton, D.F. G.A. Padbury and C.T. Stushnoff. 1998 *The Ecoregions of Saskatchewan*. Regina: Canadian Plains Research Center.

- Agee, J.K. 1993. Fire ecology of Pacific Northwest forests. Island Press. Washington, D.C. 493 pp.
- Alberta Environmental Protection. 1994b. Alberta protected areas system analysis (1994). Report 3: Revised by P. Achuff, December 7, 1994, Edmonton, Alberta.
- Alberta Environmental Protection. 1997. The parkland natural region of Alberta. One of a series of reports prepared for the Special Places 1000 Provincial Coordinating Committee. Alberta Environmental Protection, Natural Resource Service, Recreation and Protected Areas Division, Natural Heritage Planning and Evaluation Branch, Edmonton, Alberta.
- Allen, L. 2000. Alberta Natural Heritage Information Center Preliminary Plant Community Tracking List, Alberta Community Development, Edmonton, Alberta.
- Anderson, M.L. 1972. The effect of fire on grasslands in the Alberta Aspen Parkland. Master of Science Thesis. Dept. of Plant Science, University of Alberta, Edmonton, Alberta.
- Anderson, H.G. and A.W. Bailey 1980. Effects of annual burning on grassland in the aspen parkland of east-central Alberta. Canadian Journal of Botany. 58: 985-996.
- ANHIC, 2001. Alberta Natural Heritage Information Center Vascular Plant Tracking List, Updated March 19, 2001. Alberta Sustainable Resource Development, Public Lands Division, Edmonton, Alberta. Available on-line at <http://www.cd.gov.ab.ca/preserving/parks/anhic/vastrak.asp>
- Baker, W. L. 1984a. A preliminary classification of the natural vegetation of Colorado. Great Basin Naturalist 44(4):647-676.
- Baker, W. L., and S. C. Kennedy. 1985. Presettlement vegetation of part of northwestern Moffat County, Colorado, described from remnants. Great Basin Naturalist 45(4):747-777.
- Beckingham, J. 1993. Ecosystem associations of northern Alberta. Alberta Forest Service, Timber Management Branch, Edmonton, AB.
- Beckingham, J.D., D.G. Nielsen, and V.A. Futoransky. 1996. Field guide to ecosites of the mid-boreal ecoregions of Saskatchewan. Canadian Forest Service, Northwest Region, Northern Forestry Center, Edmonton, Alberta.
- Bird, R.D. 1961. Ecology of the aspen parkland of western Canada in relation to land use. Canada Agriculture Publication 1066 (II), I, X, 155p.

- Bourgeron, P. S., and L. D. Engelking, editors. 1994. A preliminary vegetation classification of the western United States. Unpublished report. The Nature Conservancy, Western Heritage Task Force, Boulder, CO. 175 pp. plus appendix.
- Bradley, Lynn and Cheryl Bradley. 1977. Aspen Groveland Resource Assessment: Neutral Hills Area. Parks Planning and Design Branch, Alberta Recreation, Parks and Wildlife, Edmonton, Alberta.
- Budd, A.C. 1987. Budd's flora of the Canadian prairie provinces. Research Branch Agriculture Canada. Publ. 1662. Ottawa, ON.
- Carsey, K., D. Cooper, K. Decker, and G. Kittel. 2001. Comprehensive statewide wetlands classification and characterization; Wetland plant associations of Colorado. Prepared for Colorado Department of Natural Resources, Colorado Natural Heritage Program, College of Natural Resources, Fort Collins, CO.
- Coupland, R.T. and T.C. Brayshaw 1953. The fescue grassland in Saskatchewan. *Ecology*. 34(2): 386-405.
- Crack, S., and B.J. Danielson. 1974. An Ecological Study of Beauvais Lake Provincial Park. Alberta Dep't of Lands and Forests.
- Dodd, J.D. and R.T. Coupland. 1966. Vegetation of saline areas in Saskatchewan. *Ecology*. 47:958-967.
- Douglas, G. W., and L. C. Bliss. (1977). Alpine and high subalpine plant communities of the North Cascades Range, Washington and British Columbia. *Ecological Monographs* 47:113-150
- Driscoll, R. S., D. L. Merkel, D. L. Radloff, D. E. Snyder, and J. S. Hagihara. 1984. An ecological land classification framework for the United States. USDA Forest Service. Miscellaneous Publication No. 1439. Washington, DC. 56 pp.
- Fehr, Alan W. 1982. The Candidate Rumsey Ecological Reserve: A Biophysical Inventory. Alberta Energy and Natural Resources, Public Lands Division, Edmonton, Alberta.
- Fehr, Alan W. 1984. Wainwright Study Area: A Biophysical Inventory. Alberta Energy and Natural Resources, Public Lands Division, Edmonton, Alberta.
- Gersib, R. A., and G. Steinauer. 1991. A biological inventory and general assessment of eastern Nebraska saline wetlands in Lancaster and southern Saunders counties. *Transactions of the Nebraska Academy of Sciences*, Volume VIII:37-44.

- Girard, M. M., H. Goetz, and A. J. Bjugstad. 1989. Native woodland habitat types of southwestern North Dakota. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. Research Paper RM-281. Fort Collins, CO. 36 pp.
- Greenall, J. A. 1996. Manitoba's terrestrial plant communities. Manitoba Conservation Data Center MS Report 96-02.
- Griffiths, D.E., and G.C.D. Griffiths. 1987a. Biophysical survey of Sylvan Lake Natural Area, Alberta. Public Lands Division, Alberta Forestry, Lands and Wildlife, Edmonton, Alberta.
- Griffiths, D.E., and G.C.D. Griffiths. 1987b. Biophysical survey of Bilby Natural Area. Alberta Forestry, Lands and Wildlife Information Center, Edmonton, Alberta.
- Griffiths, C.D., D.E. Griffiths, and W.S. Taylor. 1996. Biophysical and significant areas inventory of the Lister Lake study area (1995). Report prepared for Resource Data Division, Alberta Environmental Protection, Edmonton, Alberta.
- Griffiths, G.C.D., D.E. Griffiths, and R. Priddle. 1997. Floristic and rare plant survey of the Blackfoot Provincial Recreation Area (1996). Natural resource service/Parks, Parkland Region, Alberta Environmental Protection, Rimbey, Alberta.
- Grossman, D.H., K.L. Goodin, and C.L. Reuss. 1994. Rare plant communities of the conterminous United States, an initial survey. The Nature Conservancy, Arlington, VA. 619p.
- Grossman, D.H., D. Faber-Langendoen, A.S. Weakley, M. Anderson, P. Bourgeron, R. Crawford, K. Goodin, S. Landaal, K. Metzler, K. Patterson, M. Pyne, M. Reid, and L. Sneddon. 1998. International classification of ecological communities: terrestrial vegetation of the United States. Volume I. The National Vegetation Classification System: development, status, and applications. The Nature Conservancy, Arlington, Virginia, USA.
- Hansen, P, K. Boggs, R. Pfister, and J. Joy. 1990. Classification and management of riparian and wetland sites in central and eastern Montana. Missoula, MT: University of Montana, School of Forestry, Montana Forest and Conservation Experiment Station, Montana Riparian Association. 279 p.
- Hansen, P. L. 1985. An ecological study of the vegetation of the Grand River/Cedar River, Sioux, and Ashland districts of the Custer National Forest. Unpublished dissertation. South Dakota State University. 257 pp.
- Hansen, P. L., and G. R. Hoffman. 1988. The vegetation of the Grand River/Cedar River, Sioux, and Ashland districts of the Custer National Forest: A habitat type classification. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. General Technical Report RM-157. Fort Collins, CO. 68 pp.

- Hansen, P. L., G. R. Hoffman, and A. J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: A habitat type classification. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. General Technical Report RM-113. Fort Collins, CO. 35 pp.
- Hansen, P. L., R.D. Pfister, K. Boggs, J. Cook, J. Joy, and D.K Hinckley. 1995. Classification and management of Montana's riparian sites. Montana Forest and Conservation Experiment Station, School of Forestry, Misc. Pub. No. 43. University of Montana, Missoula. 646 pp.
- Hansen, P.L. and G.R. Hoffman. 1988. The vegetation of the Grand River/Cedar River, Sioux, and Ashland Districts of the Custer National Forest: a habitat type classification. Gen. Tech. Rep. RM-157. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 68 p.
- Hansen, P.L., G.R. Hoffman, and A.J. Bjugstad. 1984. The vegetation of Theodore Roosevelt National Park, North Dakota: a habitat type classification. Gen. Tech. Rep. RM-113. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 35 p.
- Hanson, H. C., and W. Whitman. 1938. Characteristics of major grassland types in western North Dakota. *Ecological Monographs* 8:58-114.
- Hardy and Associates 1986. Ecological land classification of Elk Island National Park. Hardy and Associates (1978) Ltd. Calgary, Alberta.
- Heidel, B., S. V. Cooper, and C. Jean. 2000. Plant species of special concern and plant associations of Sheridan County, Montana. Report to the U.S. Fish and Wildlife Service. Montana Natural Heritage Program, Helena. 22 pp. plus appendices.
- Hilton, J.E. 1970. Forage production and utilization in the aspen parkland of Alberta following aerial application of 2,4-D and 2,4,5-T. MSc Thesis, Department of Plant Science, University of Alberta, Edmonton, Alberta.
- Hirsch, K. J. 1985. Habitat type classification of grasslands and shrublands of southwestern North Dakota. Ph.D. thesis. North Dakota State University, Fargo.
- Hoffman, G. R., and R. R. Alexander. 1987. Forest vegetation of the Black Hills National Forest of South Dakota and Wyoming: A habitat type classification. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. Research Paper RM-276. Fort Collins, CO. 48 pp.

- IISD. 1994. Sustainability of Canada's agri-food system – a prairie perspective. Published by the International Institute for Sustainable Development (IISD) for the Faculty of Agriculture and Food Studies, University of Manitoba.
- Johnson, K. R. 1939. Plant ecology of northwestern Colorado lakes and surrounding areas. Unpublished dissertation. University of Colorado, Boulder. 138 pp.
- Johnson, K. R. 1941. Vegetation of some mountain lakes and shores in northwestern Colorado. *Ecology* 22:306-316.
- Johnston, B.C. 1987. Plant associations of Region Two: Potential plant communities of Wyoming, South Dakota, Nebraska, Colorado, and Kansas. R2-ECOL-87-2. USDA Forest Service, Rocky Mountain Region. Lakewood, CO. 429 pp.
- Johnson, D., L. Kershaw, A. MacKinnon, and J. Pojar. 1995. Plants of the western boreal forest and aspen parkland. Lone Pine Publishing, Edmonton, Alberta.
- Kunze, L. M. 1994. Preliminary classification of native, low elevation, freshwater wetland vegetation in western Washington. Washington State Department of Natural Resources, Natural Heritage Program. 120 pp.
- Lieffers, V. J. and S. E. Macdonald. 1993. Ecology of and control strategies for *Calamagrostis canadensis* in boreal forest sites. *Can. J. For. Res.* 23:2070-2077.
- Little, E.L., Jr. 1979. Checklist of United States trees (native and naturalized). *Agric. Handb.* 541. Washington, DC: U.S. Department of Agriculture, Forest Service. 375 p.
- Looman, J. 1983. The vegetation of the Canadian provinces IV. The woody vegetation, Part 1. *Phytocoenlogia* 11:297-330.
- Looman, J. 1987a. The vegetation of the Canadian provinces IV. The Woody vegetation, Part 3, Deciduous woods and forests. *Phytocoenlogia* 15 (1):51-84
- Looman, J. 1987b. The vegetation of the Canadian provinces IV. The Woody vegetation, Part 4, Coniferous forests. *Phytocoenlogia* 15 (3): 289-327.
- Lynch, D. 1955. Ecology of the aspen groveland in Glacier County, Montana. *Ecological Monographs* 25:321-344.
- Marschner, F.J. 1974. The original vegetation of Minnesota. A map compiled in 1930 by F. J. Marschner from U.S. General Land Office Survey Notes and published in 1974 under the direction of M.L. Heinselman of the USDA Forest Service. St. Paul, MN: Cartography Laboratory of the Department of Geography, University of Minnesota. 1 map (1:500,000) and accompanying text.

- Meijer, M. and E. Karpuk. 1999. Dillberry Lake Provincial Park Biophysical Inventory. Resource Data Division, Alberta Environment, Edmonton.
- Metcalf, F. P. 1931. Wild duck foods of North Dakota lakes. USDA Tech. Bull. 221. 72 pp.
- Miller, John Gage. 1978. An ecological study of creeping juniper (*Juniperus horizontalis* Moench.) in Montana. Bozeman, MT: Montana State University. 154 p. Thesis.
- Mueggler, W. F. 1988. Aspen community types of the Intermountain Region. USDA Forest Service General Technical Report INT-250. Intermountain Research Station, Ogden, UT. 135 pp.
- Mueggler, W. F., and R. B. Campbell, Jr. 1982. Aspen community types on the Caribou and Targhee national forests in southeastern Idaho. USDA Forest Service, Intermountain Forest and Range Experiment Station. Research Paper INT-294. Ogden, UT. 32 pp.
- Moss, E.H. 1928. The vegetation of Alberta II. The swamp, moor and bog forest vegetation of central Alberta. *Journal of Ecology* 16:19-70.
- Moss, E.H., 1932. The vegetation of Alberta. IV. The poplar association and related vegetation of central Alberta. *Journal of Ecology* 20: 380-415.
- Moss, E.H. and J.A. Campbell. 1948. The Fescue Grassland of Alberta. *Can. J. of Research*. Vol. 25, Sec. C.
- Moss, E.H., 1952. Grassland of the Peace River Region, Western Canada. *Canadian Journal of Botany* Vol. 30: 98-124.
- Moss, E.H., 1953. Marsh and bog vegetation in Northwestern Alberta. *Canadian Journal of Botany* 31:448-470.
- Moss, E. H. 1955. The vegetation of Alberta. *Botanical Review*. 21(9): 493-567.
- Moss, E.H. 1983. *Flora of Alberta*, Second Edition, Revised by J.G. Packer. University of Toronto Press, Toronto.
- Mueller, D., R. Bowman, and W. McGinnies. 1985. Effects of tillage and manure on emergence and establishment of Russian Wildrye in a Saltgrass meadow. *Journal of Range Management* 38:497-500.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2001. Version 1.6 . Arlington, Virginia, USA: NatureServe. Available: <http://www.natureserve.org/explorer>.

- Pearson Timberline Forestry Consultants 1993. Wainwright dunes ecological reserve vegetation change/disturbance assessment. Prepared for Alberta Environmental Protection, Resource Information Division, Edmonton, Alberta.
- Parish, R., R. Coupe, and D. Lloyd (eds.). 1996. Plants of southern interior British Columbia. Lone Pine Publishing, Vancouver, BC.
- Pojar, J. 1982. Boreal and Subalpine Grasslands of Northern BC. In Nicholson, McLean & Baker, ed., Grassland Ecology and Classification Symposium Proceedings June 1982. BC Ministry of Forests. Victoria, British Columbia:249-261.
- Polster, D.F., and L.E. Watson. 1979. Vegetation classification and evaluation – Elk Island National Park. Techman Ltd. For Parks Canada. 134p.
- Redmann, R.E. & A.G. Schwarz. 1986. Dry grassland plant communities in Wood Buffalo National Park, Alberta. The Canadian Field Naturalist Vol. 100: 526-532
- Remphrey, W. R.; Steeves, T. A.; Neal, B. R. 1983. The morphology and growth of *Arctostaphylos uva-ursi* (bearberry): an architectural analysis. Canadian Journal of Botany. 61: 2430-2450.
- Renewable Resources Consulting Ltd.. 1975. Dillberry Lake - resource inventory and analysis. Prepared for Alberta Provincial Parks, Department of Lands & Forests, Edmonton, Alberta.
- Rintoul, J. 1984. Silver Valley: Biophysical Resource Analysis. Alberta Forestry, Lands and Wildlife, Edmonton, Alberta.
- Schneider, R.E., D. Faber-Langendoen, R.C.Crawford, A.S.Weakley. 1997. The Status of Biodiversity in the Great Plains: Great Plains Vegetation Classification. The Nature Conservancy, Great Plains Program. Minneapolis, MN. 75 pp.
- Schwarz, A. and R. Wein. 1997. Threatened dry grasslands in the continental Boreal forests of Wood Buffalo National Park. Canadian Journal of Botany 75:1363-1370.
- Severson, Kieth E.; Garrett, E. Chester. 1974. Growth characteristics of bearberry in the Black Hills. Res. Note. RM-254. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 3 p.
- Smith, W.W., and N. Kondla. 1972. An ecological survey of Dry Island Buffalo Jump Provincial Park. Provincial Parks Planning, Department of Lands and Forests, Government of Alberta (Alberta Environment), Edmonton, Alberta.

- Stacey, E.C. 1927. Ecological Studies on the Native Grasses of the Grande Prairie - Beaverlodge District. A thesis submitted to the Committee on Graduate Studies of the University of Alberta, Edmonton, Alberta.
- Steeves, T.A., M.W. Steeves, and A.R. Olson. 1991. Flower development in *Amelanchier alnifolia* (Maloideae). *Canadian Journal of Botany* 69:844-857.
- Strong, W.L. 1999. Botanical, Vegetation and Ecological Resource Survey of the proposed Coal Valley II Mine Extension and Subregional Ecosession Classification. Luscar Ltd - Coal Valley Mine, Edson Alberta.
- Thilenius, J. F., G. R. Brown, and A. L. Medina. 1995. Vegetation on semi-arid rangelands, Cheyenne River Basin, Wyoming. USDA Forest Service. General Technical Report RM-GTR-263. Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 60 pp.
- Thompson, W. 1994. Relationship of the national wetlands inventory to an ecological site classification of wetlands in northeast Montana. Masters Thesis. University of Montana, Missoula. 60 pp.
- Thorpe, J. and R. Godwin. 1993. Vegetation Survey of the Manito Sand Hills. Applied Plant Ecology Section, Saskatchewan Research Council, Publication No. E2550-1-E-93.
- Timoney, K., and A. Robinson. 1998. Floristic, rare plant, and vegetation survey of the blackfoot provincial recreation area uplands (1997-1998). Treeline Ecological Research Report prepared for; Operations, Natural Resources Service Parkland Region, Alberta Environmental Protection, Rimbey, Alberta.
- Ungar, I. A. 1974c. Halophyte communities of Park County, Colorado. *Bulletin of the Torrey Botanical Club* 101:145-152.
- United States Department of Agriculture. reprinted 1988. Range plant handbook. Dover ed. Mineola, N.Y.
- USFS [U.S. Forest Service]. 1992. Draft habitat types of the Little Missouri National Grasslands. Medora and McKenzie ranger districts, Custer National Forest. Dickinson, ND.
- Vujnovic, K. and J. Bentz. 2001. Preliminary classification of native wheat grass (*Agropyron* spp.) community types in Alberta. Alberta Environment Report for the Alberta Natural Heritage Information Center, Prepared by Geowest Environmental Consultants Ltd., Edmonton, Alberta.

- Vujnovic, K., and R.W. Wein. 1997. An inventory of remnant prairie grasslands within the central parkland natural sub-region of Alberta. Report prepared for Park Services, Alberta Environmental Protection, Edmonton, Alberta.
- Wallis, C. 1977. Red Deer River Resource Evaluation - 1977. Alberta Recreation, Parks and Wildlife, Edmonton, Alberta.
- Wallis, C. 1990. Reconnaissance survey of saline wetlands and springs in the Grassland-Parkland region of Eastern Alberta. World Wildlife Fund, Prairie for Tomorrow, Edmonton, Alberta, Canada.
- Weerstra, B.G. and A.C. Holcroft Weerstra. 1998. Preliminary classification of plains rough fescue (*Festuca hallii*) community types within the central parkland subregion of Alberta. Alberta Natural Heritage Information Center, Alberta Environment, Edmonton, Alberta.
- Western Rangeland Consultants Inc. 1999. Mitigation Measures and Further Recommendations for Rare Native Grasslands on the Cochrane Property and Three other adjacent to Highway 43, near Grande Prairie. Alberta Transportation and Utilities, Edmonton, Alberta.
- Wheeler, G.W. 1976. Some grassland and shrubland communities in the parklands of Central Alberta. A thesis submitted to the faculty of graduate studies. Department of Plant Science, U of Alberta
- Wilkinson, K. 1981. Remnant and Early Settlement Prairies and Solonchic Soils in the Peace River District. Master's Thesis, Dept. of Biology, University of Calgary.
- Willoughby, M. 2000. Guide to range plant community types and carrying capacity for the dry and central mixedwood subregions of Alberta : third approximation. Alberta Environment, Forest Management Division; Alberta Agriculture Food and Rural Development, Public Lands Division, Edmonton, Alberta.
- Willoughby, M., and D. Downing. 1995. Deciduous plant communities and carrying capacity for the boreal ecoprovince of Alberta. Alberta Environmental Protection, Edmonton, Alberta.
- Wroe., R.A. 1971. Synecology of a *Festuca Scabrella* Torr. Grassland. Masters Thesis, Department of Plant Science, University of Alberta, Edmonton, Alberta.