


MACROFUNGI OF LA BUTTE CREEK, FIDLER-GREYWILLOW AND COLIN-CORNWALL WILDLAND PROVINCIAL PARKS



PREPARED FOR:
ALBERTA COMMUNITY DEVELOPMENT
PARKS AND PROTECTED AREAS DIVISION

PREPARED BY:
WILLIAM RICHARDS AND DIANE MURRAY

JANUARY 2003

Executive Summary

During 2001 and 2002 a project to collect and identify the macrofungi of La Butte Creek, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks, in northeastern Alberta was undertaken. This project was part of a greater biophysical investigation carried out by Parks and Protected Areas' staff, and volunteers from a variety of scientific backgrounds.

The fungi found belong to three major groups; the Ascomycota (cup fungi), Basidiomycota (mushrooms, rusts, smuts, etc.) and Myxomycota (slime molds).

One hundred of the 156 species documented through this study have been previously recorded as occurring in Alberta. Valid Alberta records have yet to be found for the remaining 56 species. Also two species discovered (*Gloeophyllum abietinum* and *Myriosclerotinia caricic-ampullaceae*) are reportedly rare in North America.

This report may be cited as:

Richards, W. M. and D. M. Murray. 2002. Macrofungi of La Butte Creek, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks, Alberta. Prepared for Alberta Community Development, Parks and Protected Areas Division, Edmonton, Alberta.

Table of Contents

| | |
|--|----|
| Introduction..... | 1 |
| Methods | 1 |
| Results..... | 3 |
| Discussion..... | 4 |
| Summary..... | 4 |
| Recommendations..... | 5 |
| Literature Cited..... | 6 |
| General References..... | 8 |
| List of Figures | iv |
| Appendix 1. Taxonomic listing of macrofungi from La Butte Creek, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks with note of previous Alberta records..... | 9 |
| Appendix 2. Annotated species list for La Butte Creek Wildland Provincial Park..... | 15 |
| Appendix 3. Annotated species list for Fidler-Greywillow Wildland Provincial Park..... | 19 |
| Appendix 4. Annotated species list for Colin-Cornwall Lakes Wildland Provincial Park..... | 21 |
| Appendix 5. Partial list of literature accounts of fungi occurrences in Alberta | 24 |
| Appendix 6. Photographs of macrofungi from La Butte, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks..... | 28 |

List of Figures

| | |
|--|---|
| Map 1. Location of La Butte Creek, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks..... | 2 |
|--|---|

Introduction

During July 2001 and July 2002 fieldwork was undertaken by staff of Parks and Protected Areas Division of Alberta Community Development, and volunteers from a variety of scientific backgrounds. The purpose of the fieldwork was to collect baseline biophysical information in three recently established protected areas in northeastern Alberta. As part of this study, the collection and identification of the macrofungi of La Butte Creek, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks (see Map 1) was conducted. This paper represents a snapshot of the mycological flora of these sites, and by no means comprises a definitive listing of fungi for these areas.

These parks are primarily within the Kazan Upland Sub-region of the Canadian Shield Natural Region in northeastern Alberta. La Butte Creek Wildland Provincial Park contains diverse areas of wetland complexes and Precambrian shield outcrops. La Butte Creek itself is a large stream draining from the Kazan Upland. A small portion of the park is bordering on the Slave River in the Peace River Lowland Sub-region.

Fidler-Greywillow Wildland Provincial Park contains one of the most diverse sections of the north shore of Lake Athabasca in Alberta. It has extensive sandy beaches, beach ridges and bedrock outcrops. This area contains the best exposure of Athabasca Sandstone outcrops in the region and has plant species not found south of Lake Athabasca. The park also incorporates a number of islands, including Bustard Island, the largest island in the province.

Colin-Cornwall Lakes Wildland Provincial Park contains diverse areas of wetland complexes and Precambrian shield outcrops. The park is primarily within the Kazan Upland Sub-region.


Methods

Observations and collections were limited to the macrofungi. Macrofungi form “fruiting bodies” that are visible to the naked eye and are generally greater than 1 cm in diameter (Redhead 1997). These fruiting bodies are the mushrooms, toadstools and puff-balls that are commonly perceived by the public as “mushrooms”. Both terrestrial and arboreal fungi were collected. Fungi were searched for at the microhabitat level (Redhead and Berch 1997), where searches included such habitats as: woody plants (living trees and shrubs); shed bark; leaf and needle litter; coarse woody debris (dead trees, standing or fallen); beaches and shorelines of lakes, wetlands and creeks; bogs; soil; burn sites and other areas of organic accumulation.

Flying squirrel and other rodents include truffle and secotioid fungi in their diet (Zabel and Walters 1997 & Currah *et al* 2002). For this reason subterranean searches for truffle and false truffles were undertaken at all locations where evidence of rodent digging was noted. This search required scraping and sifting through a large quantity of duff using a three prong garden fork. Searches were restricted to the humus layer and covered an area approximately 40 cm square, which centered on the rodent’s original excavation.

As specimens were collected they were placed in numbered and labeled waxed paper bags. Notes on habitat, species name or family (if known); and specimen and spore colour were included. Where spore colour was not readily discernible but required for identification, an

attempt to acquire a spore print was made by placing the specimen on white paper before depositing it in the waxed bag.



Map 1. Locations of Fidler-Greywillow, La Butte Creek and Collin-Cornwall Lakes Wildlands

Scientific names are based on current nomenclature. Different references were used to identify the different groups of fungi as follows: Brandrud *et al.* (1990, 1992, 1994, 1998) was used for *Cortinarius*; Overholts (1967) and Gilbertson and Ryvarden (1986-1987) for the polypore groups; Martin and Alexopolos (1969) for the *Myxomycetes*; Smith *et al.* (1979) and Kauffman (1918) for the *Agarics*; and Moser (1983) for *Polyporales*, *Boletales*, *Agaricales*, and *Russulales*. In addition several important works were used as general identification guides, these include: Arora (1986), Philips (1991), Pomerleau (1980), Smith *et al.* (1981), Walters *et al.* (1981, 1986, 1991, 1995). Names noted with *cf. (conferre)*, compare with the species noted but because of the presence (or lack) of a feature during the keying process, a definitive naming could not be made at this time. Where possible, tentative identifications were made in the field, but all specimens were submitted for verification or identification (Schulz 2002).

Results

A complete list of the species found at each park is presented in the appendices. Table 1 contains a summary of the number of species and genera encountered for each park.

Table 1. Numbers of genera represented and species of macrofungi that were collected in three parks.

| Park | Number of Genera | Number of Species |
|----------------------|------------------|-------------------|
| La Butte Creek | 55 | 85 |
| Fidler-Greywillow | 32 | 64 |
| Colin-Cornwall Lakes | 42 | 61 |
| Total | 129 | 156 |

The fungi found belong to three major groups; the Ascomycota (cup fungi), Basidiomycota (mushrooms, rusts, smuts, etc.) and Myxomycota (slime molds). See Appendix 1 for a complete listing of species within these groups. The Ascomycetes are represented by five families with a total of nine species. The Basidiomycetes are represented by 25 families and contain a total of 143 species. The slime moles are represented by three families, each with one species.

Various sources were checked for previous occurrences of fungi in Alberta. One hundred of the 156 species documented by this study have been noted by various authors as occurring in Alberta. These authors include: Abbott 1987, 1989, and 1990; Abbott and Currah 1989; Baranyay 1968; Bradbury 1998; Calgary's Natural Areas 1980; Danielson 1984; Gilbertson 1981; Gilbertson and Ryvarden 1986-1987; Kernaghan and Currah 1998; Larsen and Denison 1978; Lawrence and Hiratsuka 1972; Paul and Etheridge 1958; Redhead 1988 and 1989; Robinson-Jeffrey and Loman 1963; Schalkwijk 1975 & 1989; Schalkwijk-Barendsen 1991 & 1991a; Thomas *et al.* 1960; and Traquair 1980. These authors were the first to report a particular species as occurring in Alberta. Valid Alberta records have yet to be found for the remaining 56 species.

Few species were abundant in any of the parks or in any habitat. *Collybia dryophila* (Bull.:Fr) Kum. within the Family Tricholomataceae was represented at each park, occurring a total of 15 times, and were the most common fungi at La Butte Creek with 10 occurrences. There was a total of 28 species represented within the Tricholomataceae and 31 within the Family

Polyporaceae. The Family Cortinariaceae had 36 species recorded. There were 90 species encountered only once during this study.

No truffles (in the Family Tuberaceae) or false truffles (in the Family Rhizopogonaceae) were discovered in the process of searching rodent dig sites. However, at several locations small gelatinous globules resembling false truffles were discovered. These globules were later determined to be egg sacks of an undetermined spider species.

Discussion

No Alberta records have been found for 56 (35.8%) of the species collected in this study. This number of new records for the province is suspected to be high, as no systematic search of herbaria was made, nor have all written reports been reviewed.

Only the microfungi of Alberta are fully recorded in checklist format (Sigler and Flis 1998). Checklists for the macrofungi have not yet been completed.

Virtually nothing is known of the relative status of any fungi species in Alberta, however two species found in the present study are considered rare. *Gloeophyllum abietinum* (Fr.) Karst., is considered “a very rare species in North America” and is known from Arizona, (Gilbertson and Ryvarden 1986-1987). Murrill (1908) describes the synonymous *G. abietinellum* Murr. as being known only from the Rocky Mountains. There was but one specimen collected during this study; it was obtained from Fidler-Greywillow from dead black spruce.

Schalkwijk-Barendsen (1991) reports *Myriosclerotinia caricic-ampullaceae* (Nyberg.) N.F. Buchw as a “rare, subarctic species”. This species has been collected twice (1976 and 1977) at the same location near Winterburn, Alberta. Both prior collections were from sedge as was the present collection from Colin-Cornwall Lakes. As the Schalkwijk-Barendsen collections were believed to be the first report of *M. caricic-ampullaceae* in northwestern North America, these collections were deposited at both the Herbarium (ALTA), University of Alberta, Edmonton, and at the National Mycological Herbarium (DAOM) Ottawa, Ontario (Schalkwijk-Barendsen 1991).

Summary

The total number of species collected in the three parks was minimal as was the number of specimens within each species. There were two main reasons for this:

- Weather conditions leading up to the field surveys had been extremely dry, not just for weeks but for the last several seasons.
- The sample period was restricted in all cases to the dry, midsummer season, which is typically slow in the fruiting of fleshy fungi. Thus, most specimens were obtained from woody substrates or along wetland margins.

The sample period for each location was short and not repeated. According to Redhead and Berch (1997), “all possible sites and microhabitats within these sites must be examined repeatedly as seasons progress”. Repeated visits to these remote sites were not, however, possible.

Of the 156 species of fungi documented, 35.8 % (56 species) were not previously reported for Alberta. The number of apparently new species is high, and probably related to the following factors:

- No systemic review of herbaria was done for fungi;
- Fungi are not often included in biophysical studies, so are under-reported.
- A through review of the literature pertaining to fungi in Alberta was not completed.

In Alberta, the macrofungi species have not been ranked for “rarity”, however:

- Two species collected during this study are considered rare: *Gloeophyllum abietinum*, (Gilbertson and Ryvarden 1986-1987) and *Myriosclerotinia caricic-ampullaceae*, (Schalkwijk-Barendsen 1991).
- Other species may also be rare, but that is impossible to substantiate at this time due to lack of information.

Fungi are one of the most important groups of organisms on the planet. They are responsible for most of the recycling of organic material back to the soil in a form that can be reused by other organisms. Fungi are vitally important for the growth of trees and most other plants, through mycorrhizal association. Currah and Van Dyk (1986) have, in one study, documented mycorrhizal association with 179 species of perennial plants native to Alberta.

Recommendations

There is a lack of knowledge of the macrofungi in Alberta. More time and research need to be devoted to understanding this group of organisms in the province.

These needs can be achieved by the following two steps:

1. Compile all known records from the existing literature for macrofungi within Alberta (Appendix 5 contains a partial list).
2. Compile all herbarium records for collections of macrofungi in Alberta.

Once these have been accomplished, provincial-wide surveys should be instituted. At a minimum, fungi investigations should be a part of all future biophysical assessments within the province. This will provide a starting point on which to base status assessments of the various species.

Literature Cited

- Abbott, S. P. 1987. The First Foray. The Stinkhorn, 1 (1): 12. Edmonton.
- Abbott, S. P. 1989. Key to Boleti of Alberta. University of Alberta, Edmonton. 12pp.
Mimeograph.
- Abbott, S. P. 1990. Shaw Lake. The Stinkhorn, 4 (1): 38-41. Edmonton.
- Abbott, S.P. and R.S. Currah. 1989. The larger cup fungi and other Ascomycetes of Alberta, an annotated checklist. Devonian Botanic Garden, University of Alberta, Edmonton. 96pp.
- Baranyay, J.A. 1968. Fungi collected during forest diseases surveys in northern Alberta and the district of Mackenzie, Northwest Territories. For Branch Dept. Public no. 1238, 25p.
- Bradbury, S.M., Danielson, R.M., and Visser, S. 1998. Ectomycorrhizas of regenerating stands of lodgepole pine (*Pinus contorta*). Can. J. Bot. 76: 218-227.
- Brandrud, T. E. , Linström, H., Merlund, H., Melot, J., and Muskos, S. 1990, 1992, 1994, 1998. Cortinarius Flora Photographica. Vol. 1-4. Cortinarius HB, Mattfors, Sweden (*Cortinarius*).
- Calgary's Natural Areas. 1980, A Report Prepared by the Natural Areas Committee of the Calgary Field Naturalists Society, December 1980. Harold W. Pinel, Editor. (on line at <http://weaselhead.org/learn/slimesmoldsfungi.htm>; viewed <last date February 10, 2003>).
- Currah, R.S., and Van Dyk, M. 1986. A survey of some perennial vascular plant species native to Alberta for the occurrence of mycorrhizal fungi. Can. Field-Nat. 100: 330-342.
- Currah RS; Smreciu EA; Lehesvirta T; Niemi M; Larsen KW. 2000. Fungi in the winter diets of northern flying squirrels and red squirrels in the boreal mixedwood forest of northeastern Alberta. Canadian Journal of Botany-Revue Canadienne de Botanique. 78(12):1514-1520.
- Danielson, R.M. 1984. Ectomycorrhizal associations in jackpine stands in northeastern Alberta, Can. J. Bot. 62:932-939.
- Gilbertson, R.L. 1981. North American wood-rotting fungi that cause brown rots. Mycotaxon 12:372—416.
- Gilbertson, R.L. and L. Ryvarden. 1986-1987. North American polypores. Fungiflora Vols. 1—2. Oslo, Norway. (*Polypores*).
- Hiratsuka, Y. 1987. Forest tree diseases of the Prairie Provinces. Env. Can. North. For. Res. Cent. Inf. Rep. No. NOR-X-286
- Kauffman, C. H. 1918. The Agaricaceae of Michigan. Wynkoop Hallenback Crawford Co., State Printers, Lansing, Michigan, USA (*Agarics*).
- Kernaghan, G. and Currah, R. S. 1998. Ectomycorrhizal fungi at tree line in the Canadian Rockies. Mycotaxon 69: 39-80.
- Lawrence J. J. and Hiratsuka. 1972. Forest fungi collection in Banff National Park, Alberta. Information Report NOR-X-22, Northern Forest Research Centre.
- Larsen, H. J. and W. C. Denison. 1978. A checklist of operculate cup-fungi (Pezizales) of North America west of the Great Plains. Mycotaxon 7:68-90.
- Martin, G. W. and Alexopolos, C. J. 1969. The Myxomycetes. University of Iowa Press, Iowa City, USA (*Myxomycetes*).
- Moser, M. (1983). *Keys to Agarics and Boleti (Polyporales, Boletales, Agaricales, Russulales)* [English translation by Plant, S., ed. Kibby, G.].
- Natural Resources Canada. 1996. Aspen trunk rot. Pages 52-53 in Common tree diseases of British Columbia. Canadian Forest Service, Pacific Forestry Centre, Victoria, BC. Edited by E.A. Allen, D.J. Morrison, and G. Wallis. 178 p. (on-line at

- http://www.pfc.cfs.nrcan.gc.ca/diseases/CTD/Group/Heart/heart14_e.html; viewed <last date February 26, 2003>).
- Overholts, L. O. 1967. The Polyporaceae of the United States, Alaska, and Canada. The University of Michigan Press, Michigan, USA (*Polypores*).
- Paul, G. and Etheridge, D.E. 1958. Decay of aspen (*Populus tremuloides* Michx.) and balsam poplar (*P. balsamifera* L.) in the Lesser Slave Lake region of Alberta. Alberta Dep. Land For., For. Serv. Div., Edmonton, Alberta, and Can. Dep. Agric., For. Biol. Div., V+Calgary, Alberta, Mimeograph.
- Redhead, S.A. 1988. Notes on the genus Xeromphalina (Agaricales, Xerulaceae) in Canada: biogeography, nomenclature, taxonomy. Can. J. Bot. 66:479-507.
- Redhead, S.A. 1989. A biogeographical overview of the Canadian mushroom flora. Can. J. Bot. 67:3003—62.
- Redhead, S.A. 1997. Macrofungi of British Columbia: requirements for inventory. Res. Br., B.C. Min. For., and Wildl. Br. B.C. Min. Envirn., Lands and Parks. Victoria, B.C. Work Pap. 28/1997. 119 pgs.
- Redhead, S.A., and S. Berch. 1997. Standardized Inventory Methodologies for Components of British Columbia's Biodiversity: MACROFUNGI (including the phyla Ascomycota and Basidiomycota). Version 1.1. Resources inventory committee approved standard. British Columbia.
- Robinson-Jeffrey, R. C., and A. A. Loman. 1963. Fungi Isolated in Culture from Red Heartwood Stain and Advance decay of Lodgepole Pine in Alberta. Can. J. Bot. 41:1371-1375.
- Schalkwijk, H.M.E. 1975. Mushrooms of the Edmonton Area - edible and poisonous. Printed and published in Edmonton 34pp.
- Schalkwijk, H.M.E. 1989. Leni Schalkwijk's checklist of Alberta Fungi. The Stinkhorn, Edmonton 3 (1): 1-66.
- Schalkwijk-Barendsen, H. M. E. 1991. Mushrooms of Western Canada. Lone Pine Publishing, Edmonton, Alberta, Canada.
- Schalkwijk-Barendsen, H.M.E. 1991a. Checklist to accompany the book – Mushrooms of Western Canada, published by author.
- Schulz, M. 2002. Personal Commutations. Mycologist, Department of Biological Science. University of Alberta. Edmonton Alberta.
- Sigler, L. and A.L. Flis. 1998. University of Alberta microfungus collection and herbarium (UAMH) Devonian Botanic Garden, Edmonton, AB, Canada.
- Smith, A. H., Smith, H. V., and Weber, N.S. 1979. How to Know the Gilled Mushrooms. Wm. C. Brown Company Publishers, Dubuque, Iowa, USA (*Agarics*).
- Thomas, G.P., D. E. Etherridge and G. Paul. 1960. Fungi and decay in aspen and balsam poplar in the boreal forest region, Alberta. Can J. Bot. 38: 459-466.
- Traquair, J.A. 1980. The first Mary Elliot Mycological foray. Alberta Naturalist. 10: 4-6.
- Zabel, C. J. and J. R. Waters. 1997. Food preferences of captive northern flying squirrels from the Lassen National Forest in northeastern California. Northwest Science, 71:103–107.
- Zeglen, S. 1997 . Tree wounding and partial-cut harvesting: a literature review for British Columbia (Pest management report ; no. 14) Ministry of Forests, Pacific Forestry Centre, Canadian Forest Service.

General References

- Arora, D. 1986. *Mushrooms Demystified*. Ten Speed Press, Berkley, California, USA (*General*).
Philips, R. 1991. *Mushrooms of North America*. Little, Brown and Company, Toronto, Ontario, Canada (*General*).
Pomerleau, R. 1980. *Flore des Champignons au Quebec*. Les Editions la Presse, Ltee, Ottawa, Canada (*General*).
Smith, A. H., Smith, H. V., and Weber, N.S. 1981. *How to Know the Non-gilled Mushrooms*. Wm. C. Brown Company Publishers, Dubuque, Iowa, USA (*General non-Agarics*).
Walters, V. L., Breitenbach, J., and Kränzlin, F. 1981. 1986, 1991. 1995. *Fungi of Switzerland*. Vol. 1-4. Verlag Mykologia, Luzern, Switzerland (*General*).

Appendix 1. Taxonomic listing of macrofungi recorded for this study with reference to known Alberta records

| Division | Subdivision | Class | Order | Family | Species | Known Alberta Record |
|----------|-----------------|----------------|--------------|-----------------|---|-----------------------------|
| Eumycota | | | | | | |
| | Ascomycotina | | | | | |
| | | Discomycetes | | | | |
| | | | Leotiales | | | |
| | | | | Leotiaceae | | |
| | | | | | <i>Bisporella citrina</i> (Batch ex Fr.) Korf & Carpenter | Abbott and Currah (1989) |
| | | | | | <i>Neobulgaria pura</i> (Fr.) Petrak | |
| | | | | Sclerotiniaceae | | |
| | | | | | <i>Myriosclerotinia caricic-ampullaceae</i> (Nyberg.) N.F. Buchw. | Schalkwijk (1989) |
| | | | Pezizales | | | |
| | | | | Humariaceae | | |
| | | | | | <i>Scutellinia scutellata</i> (L. ex St. Amans) Lambotte | Traquair (1980) |
| | | | | Morchellaceae | | |
| | | | | | <i>Morchella elata</i> Fr. | Schalkwijk (1975) |
| | | | | | <i>Gyromitra esculenta</i> (Pers.) Fr. | Larsen and Denison (1978) |
| | | | | Pezizaceae | | |
| | | | | | <i>Melastiza chateri</i> (W. G. Smith) Boud. | Abbott and Currah (1989) |
| | | | | | <i>Peziza badia</i> Pers. et Merát | Abbott and Currah (1989) |
| | | | | | <i>Peziza praetervisa</i> Bres. | Larsen and Denison (1978) |
| | Basidiomycotina | | | | | |
| | | Gasteromycetes | | | | |
| | | | Lycoperdales | | | |
| | | | | Lycoperdaceae | | |
| | | | | | <i>Lycoperdon perlatum</i> Pers.: Pers. | Traquair (1980) |
| | | | | Astraeaceae | | |
| | | | | | <i>Astraeus hygrometricus</i> (Pers.) Morgan | Danielson (1984) |
| | | Hymenomycetes | | | | |
| | | | Agaricales | | | |
| | | | | Agaricaceae | | |
| | | | | | <i>Agaricus bernardii</i> (Quél.) Sacc. | |
| | | | | | <i>Agaricus silvicola</i> Vitt. Sac. | Schalkwijk-Barendsen (1991) |
| | | | | Bolbitiaceae | | |
| | | | | | <i>Agrocybe praecox</i> (Persoon: Fries) Fayod | Schalkwijk (1989) |
| | | | | | <i>Agrocybe putaminum</i> (Fr.) Sing. | |
| | | | | | | |
| | | | | | | |

| Division | Subdivision | Class | Order | Family | Species | Previous Alberta Record |
|----------|-------------|-------|-------|----------------|---|-----------------------------|
| | | | | Boletaceae | | |
| | | | | | <i>Leccinum insigne</i> Smith | Abbott (1989) |
| | | | | | <i>Leccinum scabrum</i> (Fr.) S. F. Gray | Abbott (1989) |
| | | | | Coprinaceae | | |
| | | | | | <i>Coprinus hemerobius</i> Fr. | |
| | | | | | <i>Coprinus lagopoides</i> Karst. | Schalkwijk (1989) |
| | | | | | <i>Coprinus micaceus</i> (Bull.:Fr.) Fr. | Thomas et al (1960) |
| | | | | | <i>Coprinus truncorum</i> (Scop.) Fr. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Psathyrella cf. gracilis</i> (Fr.) Quél. | |
| | | | | | <i>Psathyrella cf. phegophila</i> Romagn. | |
| | | | | | <i>Psathyrella cf. velutina</i> (Fr.) Singer | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Psathyrella subnuda</i> (Karst.) A. H. Smith | Schalkwijk-Barendsen (1991) |
| | | | | Cortinariaceae | | |
| | | | | | <i>Cortinarius armillatus</i> (Fr.) Fr. | Danielson (1984) |
| | | | | | <i>Cortinarius brunneus</i> var. <i>grandicolor</i> (Fr.:Fr.) Lindstr. & Melot ¹ | Kernaghan and Currah (1998) |
| | | | | | <i>Cortinarius cf. casimiri</i> (Velen) Huijsman | |
| | | | | | <i>Cortinarius cf. huronensis</i> Ammirati & Smith | |
| | | | | | <i>Cortinarius cf. jubarinus</i> France:DM | |
| | | | | | <i>Cortinarius cf. privignus</i> Fr. | |
| | | | | | <i>Cortinarius cf. raphanoides</i> (Fr.) Fr. | |
| | | | | | <i>Cortinarius cf. vespertinus</i> (Fr.:Fr.) Fr. | |
| | | | | | <i>Cortinarius citrinifolius</i> Smith | |
| | | | | | <i>Cortinarius helobius</i> Romagn. | |
| | | | | | <i>Cortinarius mucosus</i> (Bull. Ex Fr.) Fr. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Cortinarius obtusus</i> (Fr.:Fr.) Fr. | Bradbury et al (1998) |
| | | | | | <i>Cortinarius trivialis</i> Lange | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Crepidotus ellipsoideus</i> Hes. et Smith | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Crepidotus mollis</i> (Fr.) Stde. | Traquair (1980) |
| | | | | | <i>Dermocybe cinnamomeobadia</i> (R. Hry.) Mos. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Dermocybe uliginosa</i> (Berk.) Mos. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Galerina cinctula</i> Orton | |
| | | | | | <i>Galerina hypnorum</i> (Schrank ex Fr.) Kuehn. | |
| | | | | | <i>Galerina mniophila</i> (Lash.) Kuhn. | |
| | | | | | <i>Galerina paludosa</i> (Fr.) Kuh. | Redhead (1989) |
| | | | | | <i>Gymnopilus cf. humicola</i> Harding ex Singer | |
| | | | | | <i>Hebeloma cf. longicaudum</i> (Fr.) S. S. Lge | |
| | | | | | <i>Hebeloma cf. velatum</i> Peck. | |
| | | | | | <i>Inocybe cf. calamistrata</i> (Fr.) Gill. | |

| Division | Subdivision | Class | Order | Family | Species | Previous Alberta Record |
|----------|-------------|-------|-------|------------------|---|-----------------------------|
| | | | | | <i>Inocybe cf. fastigata</i> (Schaeff.) Quél | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Inocybe cf. frigidula</i> Favre | |
| | | | | | <i>Inocybe cf. gymnocarpa</i> Kuehn | |
| | | | | | <i>Inocybe cf. petiginosa</i> (Fr.: Fr.) Gillet. | Schalkwijk (1989) |
| | | | | | <i>Inocybe lacera</i> (Fr.) Kummer | |
| | | | | | <i>Inocybe mixtilis</i> Britz. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Inocybe</i> sp. infected with <i>Cladosporium</i> sp | |
| | | | | | <i>Tubaria furfuracea</i> (Pers. ex Fr.) Gillet | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Tubaria hiemalis</i> Romagn. Ex Bon. | |
| | | | | | <i>Tubaria pellucida</i> (Bull. Ex Fr.) Gill. | |
| | | | | | <i>Tubaria romagnesiana</i> Arnolds | |
| | | | | Entolomataceae | | |
| | | | | | <i>Rhodocybe caelata</i> (Fr.) Mre. | |
| | | | | Hygrophoraceae | | |
| | | | | | <i>Hygrophorus</i> sp. | |
| | | | | Paxillaceae | | |
| | | | | | <i>Hygrophoropsis aurantiaca</i> (Wulf. ex Fr.) Maire | Schalkwijk-Barendsen (1991) |
| | | | | Pleurotaceae | | |
| | | | | | <i>Pleurotus ostreatus</i> (Jacq. ex Fr.) Singer | Schalkwijk-Barendsen (1991) |
| | | | | Pluteaceae | | |
| | | | | | <i>Pluteus cervinus</i> (Schaeff.) Kumm. | Traquair (1980) |
| | | | | | <i>Pluteus pestatus</i> (Fr.) Gillet | Schalkwijk (1989) |
| | | | | Russulaceae | | |
| | | | | | <i>Lactarius cf. obscuratus</i> (Lasch) Fr. | |
| | | | | | <i>Lactarius fragilis</i> (Burmington) Hesler & Smith | |
| | | | | | <i>Russula cf. maculata</i> Quél. et Roz. | |
| | | | | | <i>Russula emetica</i> (Schaeff. Ex Fr.) S. F. Gray | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Russula fragilis</i> (Pers. ex. Fr.) Fr. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Russula vesca</i> Fr. | |
| | | | | Strophariaceae | | |
| | | | | | <i>Kuehneromyces lignicola</i> (Peck.) Redhead | Schalkwijk (1989) |
| | | | | | <i>Pholiota spumosa</i> (Fr.) Sing. | Abbott (1990) |
| | | | | Tricholomataceae | | |
| | | | | | <i>Armillaria mellea</i> (Vahl ex Fr.) Karsten | Paul and Etheridge (1958) |
| | | | | | <i>Clitocybe truncicola</i> (Pk.) Sacc. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Collybia confusa</i> Orton | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Collybia dryophila</i> (Bull.:Fr.) Kum. | Abbott (1990) |
| | | | | | <i>Collybia marasmioides</i> (Britz) Bresinsky & Romagn | |

| Division | Subdivision | Class | Order | Family | Species | Previous Alberta Record |
|----------|-----------------|------------------|-------|--------|--|---------------------------------|
| | | | | | <i>Fayodia pseudoclusilis</i> (Joss. & Konr.) Sing. | |
| | | | | | <i>Hemimycena gracilis</i> (Quél.) Sing. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Hygrocybe cantharellus</i> (Schw.) Mull. | Schalkwijk (1989) |
| | | | | | <i>Hygrotrama cf. hymenocephalum</i> (Sm. & Hesl.) Sing | |
| | | | | | <i>Laccaria laccata</i> (Scop.: Fr.) Berk. & Br | Traquair (1980) |
| | | | | | <i>Laccaria proxima</i> (Boud.) Pat. | Danielson (1984) |
| | | | | | <i>Lentinus lepideus</i> (Fr. ex Fr.) Fr. | |
| | | | | | <i>Lentinus sulcatus</i> Berkeley | Calgary's Natural Areas (1980) |
| | | | | | <i>Leptista (=Clitocybe) sp.</i> | |
| | | | | | <i>Marasmius pallidocephalus</i> Gillam | Traquair (1980) |
| | | | | | <i>Marasmius scorodonius</i> (Fr.) Fr. | |
| | | | | | <i>Marasmius strictipes</i> (Pk.) Singer | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Mycena cf. citrinomarginata</i> Gillet | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Mycena cf. galopus</i> (Pers.:Fr.) Kumm. | |
| | | | | | <i>Mycena cf. stipata</i> M. Geest & Schwobel | |
| | | | | | <i>Mycena filopes</i> (Bull.:Fr.) Kummer | |
| | | | | | <i>Mycena leucogala</i> (Cooke) Sacc. | |
| | | | | | <i>Mycena stannea</i> (Fr.) Quélét | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Omphalina ericetorum</i> (Fr.) M. Lange | Traquair (1980) |
| | | | | | <i>Omphalina pyxidata</i> (Bull. Ex Fr.) Quél. | |
| | | | | | <i>Oudemansiella longipes</i> (St. Ann.) Moser | |
| | | | | | <i>Phyllotopsis nidulans</i> (Fr.) Singer | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Xeromphalina fraxinophila</i> Smith | Redhead (1988) |
| | Aphyllophorales | | | | | |
| | | Corticiaceae | | | | |
| | | | | | <i>Peniophora</i> sp. | Abbott (1987) |
| | | | | | <i>Plicatuopsis crispa</i> (Fr.) Reid. | Bradbury (1968) |
| | | Ganodermataceae | | | | |
| | | | | | <i>Gandoderma appplanatum</i> (Pers.) Pat. | Bradbury (1968) |
| | | Hydnaceae | | | | |
| | | | | | <i>Sarcodon scabrosus</i> (Fr.) Karst | Abbott (1990) |
| | | Hymenochaetaceae | | | | |
| | | | | | <i>Coltricia perrenis</i> (Fr.) Murr. | Danielson (1984) |
| | | | | | <i>Phellinus chrysoloma</i> (Fr.) Donk. | Gilbertson and Ryvarden (1987) |
| | | | | | <i>Phellinus ignarius</i> (L.:Fr.) Quél. ² | Thomas et al (1960) |
| | | | | | <i>Phellinus pini</i> (Thore.:Fr.) A. Ames ³ | Robinson-Jeffrey & Loman (1963) |
| | | | | | <i>Phellinus tremulae</i> (Bond.) Bond. & Boriss. ⁴ | Paul and Etheridge (1958) |

| Division | Subdivision | Class | Order | Family | Species | Previous Alberta Record |
|----------|-------------|-------|-------|------------------|--|---------------------------------|
| | | | | Polyporaceae | | |
| | | | | | <i>Coriolopsis gallica</i> (Fr.) Ryv. | Gilbertson and Ryvarden (1986) |
| | | | | | <i>Coriolopsis</i> sp. | |
| | | | | | <i>Daedaleopsis confragosa</i> (Bolt.:Fr.) Schroet. | Schalkwijk (1989) |
| | | | | | <i>Fomes fomentarius</i> (L.:Fr.) Kickx | Thomas et al (1960) |
| | | | | | <i>Fomitopsis cajanderi</i> (Karst.) Kotl. et Pouz. ⁵ | Robinson-Jeffrey & Loman (1963) |
| | | | | | <i>Fomitopsis pinicola</i> (Swartz.:Fr.) Karst. ⁶ | Robinson-Jeffrey & Loman (1963) |
| | | | | | <i>Fomitopsis rosea</i> (Alb. et Schw.:Fr.) Karst. | Gilbertson (1981) |
| | | | | | <i>Gloeophyllum abietinum</i> (Fr.) Karst. | |
| | | | | | <i>Gloeophyllum protractum</i> (Fr.) Imaz. | Gilbertson and Ryvarden (1986) |
| | | | | | <i>Gloeophyllum sepiarium</i> (Fr.) Karst. ⁷ | Robinson-Jeffrey & Loman (1963) |
| | | | | | <i>Gloeophyllum trabium</i> (Fr.) Murr. | Traquair (1980) |
| | | | | | <i>Gleoporus taxicola</i> (Pers.:Fr.) Gilbn & Ryv. | Gilbertson and Ryvarden (1986) |
| | | | | | <i>Inonotus obliquus</i> (Pers.:Fr.) Pilát | Gilbertson and Ryvarden (1986) |
| | | | | | <i>Irpex lacteus</i> (Fr.:Fr.) Fr. | Schalkwijk (1989) |
| | | | | | <i>Piptoporus betulinus</i> (Bull.:Fr.) Karst. ⁸ | Bradbury (1968) |
| | | | | | <i>Polyporus alveolaris</i> (DC.:Fr.) Bond & Sing. ⁹ | Abbott (1987) |
| | | | | | <i>Polyporus brumalis</i> Pers.:Fr. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Polyporus elegans</i> Bull.:Fr. | Traquair (1980) |
| | | | | | <i>Polyporus melanopus</i> Fr. | Gilbertson and Ryvarden (1987) |
| | | | | | <i>Polyporus squamosus</i> Huds.: Fr. | Gilbertson and Ryvarden (1987) |
| | | | | | <i>Polyporus varius</i> Fr. | Lawrence and Hiratsuka (1972) |
| | | | | | <i>Polyporus varius</i> Persoon:Fries forma nummularis | Schalkwijk (1989) |
| | | | | | <i>Pycnoporus cinnabarinus</i> (Jacq.:Fr.) Karst. | Schalkwijk-Barendsen (1991) |
| | | | | | <i>Skeletocutis amorpha</i> (Fr.) Kokl. & Pouz. | Gilbertson and Ryvarden (1987) |
| | | | | | <i>Trametes cervina</i> (Schw.) Bres. | |
| | | | | | <i>Trametes hirsuta</i> (Wulf.:Fr.) Pil. | Calgary's Natural Areas (1980) |
| | | | | | <i>Trametes pubescens</i> (Schum.:Fr.) Pilát ¹⁰ | Bradbury (1968) |
| | | | | | <i>Trametes suaveolens</i> L.:Fr. | Calgary's Natural Areas (1980) |
| | | | | | <i>Trametes versicolor</i> (L.: Fr.) Pilát | Gilbertson and Ryvarden (1987) |
| | | | | | <i>Trichaptum abietinum</i> (Dicks.:Fr.) Ryv. ¹¹ | Bradbury (1968) |
| | | | | | <i>Trichaptum biforme</i> (Fr. in Kl.) Ryv | Abbott (1990) |
| | | | | Schizophyllaceae | | |
| | | | | | <i>Schizophyllum commune</i> Fr. | Schalkwijk-Barendsen (1991) |

| Division | Subdivision | Class | Order | Family | Species | Previous Alberta Record |
|--------------|-------------|-------------|--------------|-----------------|--|---------------------------|
| | | | | Stereaceae | | |
| | | | | | <i>Chondrostereum purpureum</i> Fr. ex Fr. ¹² | Paul and Etheridge (1958) |
| | | | | | <i>Stereum hirsutum</i> Fr. | Bradbury (1968) |
| | | | | | <i>Stereum striatum</i> (Fr.:Fr.) Fr. | |
| | | | | Thelephoraceae | | |
| | | | | | <i>Thelephora terrestris</i> Fr. | Danielson (1984) |
| | | | | Dacrymycetales | | |
| | | | | | Dacrymycetaceae | |
| | | | | | <i>Dacrymyces deliquescens</i> (Mérat) Duby | Bradbury (1968) |
| | | | | | <i>Dacrymyces palmatus</i> | |
| | | | Tremellales | | | |
| | | | | Tremellaceae | | |
| | | | | | <i>Exidia glandulosa</i> Fr. | Traquair (1980) |
| | | | | | <i>Exidia nucleata</i> (Schw.) Burt. | |
| | | | | | <i>Tremella foliacea</i> (Pers. & S. F. Gray) Pers. | |
| Myxomycotina | | | | | | |
| | | Myxomycetes | | | | |
| | | | Liceales | | | |
| | | | | Reticulariaceae | | |
| | | | | | <i>Lycogala epidendrum</i> (L.) Fries | Bradbury (1968) |
| | | | Physarales | | | |
| | | | | Physaraceae | | |
| | | | | | <i>Leocarpus fragilis</i> (Dicks) Rost. | |
| | | | Stemonitales | | | |
| | | | | Stemonitidaceae | | |
| | | | | | <i>Stemonitis splendens</i> Rost. | |

¹ Previously recorded as species only.

² Recorded as *Fomes igniarius* (L. ex Fr) Kickx, and recognized as a synonym by Zeglen (1997)

³ Recorded as *Fomes pini* (Thore: Fr.) Fr., and recognized as a synonym by Hiratsuka (1987)

⁴ Recorded as *Fomes igniarius* var. *populinus* (Neuman) Camp., and recognized as a synonym by Natural Resources Canada (1996)

⁵ Recorded as *Fomes cajanderi* Karst., and recognized as a synonym by Gilbertson and Ryvarden (1986-1987)

⁶ Recorded as *Fomes pinicola* (Sw.:Fr.) Cooke, and recognized as a synonym by Hiratsuka (1987)

⁷ Recorded as *Lenzites saepiaria* (Wulfen:Fr.) Fr., and recognized as a synonym by Hiratsuka (1987)

⁸ Recorded as *Polyporus betulinus* (Bull.:Fr.) Fr., and recognized as a synonym by Hiratsuka (1987)

⁹ Recorded as *Favolus alveolaris*, and recognized as a synonym by Gilbertson and Ryvarden (1986-1987)

¹⁰ Recorded as *Polyporus pubescens* (Schum.:Fr.) Pilat, and recognized as a synonym by Gilbertson and Ryvarden (1986-1987)

¹¹ Recorded as *Polyporus abietinus* (Dickson:Fr.) Donk, and recognized as a synonym by Gilbertson and Ryvarden (1986-1987)

¹² Recorded as *Stereum purpureum* (Pers.) Fr., and recognized as a synonym by Hiratsuka (1987)

Appendix 2. Annotated species list for La Butte Creek Wildland

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|---|--|-----------|-----------|------------|--------------|-----------------|
| (? <i>Inocybe</i> sp.) infected with <i>Cladosporium</i> sp | on moss with black spruce | 12V | 485975.90 | 6581957.42 | July 15 2001 | LBC107a |
| <i>Chondrostereum purpureum</i> | on dead aspen poplar bark | 12V | 491127.72 | 6580205.28 | July 8 2001 | LBC032 |
| <i>Clitocybe truncicola</i> | in moss on rotted log | 12V | 477088.94 | 6584366.43 | July 14 2001 | LBC070 |
| <i>Collybia cf. dryophila</i> | on moss in spruce bog | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC097 |
| <i>Collybia cf. dryophila</i> | on moss in spruce bog | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC098 |
| <i>Collybia dryophila</i> | on ground in moss in willow forest | 12V | 474464.27 | 6587728.02 | July 6 2001 | LBC011 |
| <i>Collybia dryophila</i> | on moss in white spruce forest | 12V | 474126.65 | 6585095.84 | July 7 2001 | LBC029 |
| <i>Collybia dryophila</i> | on aspen poplar leaves | 12V | 491127.72 | 6580205.28 | July 8 2001 | LBC034 |
| <i>Collybia dryophila</i> | on moss in aspen forest | 12V | 491127.72 | 6580205.28 | July 8 2001 | LBC035 |
| <i>Collybia dryophila</i> | in moss of white spruce /aspen forest | 12V | 491127.72 | 6580205.28 | July 8 2001 | LBC036 |
| <i>Collybia dryophila</i> | in moss in black spruce bog | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC044 |
| <i>Collybia dryophila</i> | with white spruce and balsam poplar | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC088a |
| <i>Collybia dryophila</i> | in white spruce stand | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC091 |
| <i>Collybia marasmoides</i> | on ground in moss in willow forest | 12V | 474464.27 | 6587728.02 | July 6 2001 | LBC010 |
| <i>Coprinus hemerobius</i> | on mud of beaverdam | 12V | 489206.86 | 6580689.96 | July 7 2001 | LBC016b |
| <i>Coprinus micaceus</i> | on equisetum sp. | 12V | 489206.86 | 6580689.96 | July 7 2001 | LBC016 |
| <i>Coprinus micaceus</i> | on moss leaves under willow | 12V | 474853.33 | 658 41.35 | July 6 2001 | LBC016a |
| <i>Coriolopsis gallica</i> | on fallen aspen with bark remaining | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC082 |
| <i>Coriolopsis</i> sp. | on fallen aspen with bark remaining | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC120 |
| <i>Cortinarius cf. huronensis</i> | [with moss] | 12V | 478813.83 | 6579354.52 | July 15 2001 | LBC106 |
| <i>Cortinarius cf. huronensis</i> | on moss with black spruce | 12V | 485975.90 | 6581957.42 | July 15 2001 | LBC107 |
| <i>Cortinarius cf. privignus</i> | in duff of white spruce/aspen poplar stand | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC067 |
| <i>Cortinarius cf. raphanoides</i> | on moss in black spruce bog | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC094 |
| <i>Cortinarius cf. vespertinus</i> | in moss in black spruce bog | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC043 |
| <i>Cortinarius citrinifolius</i> | [?] | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC093 |
| <i>Cortinarius mucosus</i> | on game trail in aspen forest, with grasses, bearberry & vetch | 12V | 474126.65 | 6585095.84 | July 7 2001 | LBC023 |
| <i>Cortinarius trivialis</i> | in moss in alder forest | 12V | 474126.65 | 6585095.84 | July 7 2001 | LBC024 |

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|--------------------------------------|--|-----------|-----------|------------|--------------|-----------------|
| <i>Crepidotus ellipsoideus</i> | on dead alder | 12V | 486095.89 | 6582093.18 | July 10 2001 | LBC050 |
| <i>Crepidotus ellipsoideus</i> | on alder/old, dry | 12V | 477499.07 | 6584378.30 | July 10 2001 | LBC059 |
| <i>Crepidotus ellipsoideus</i> | on wood | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC092 |
| <i>Crepidotus mollis</i> | on dead wood | 12V | 477783.17 | 6584510.43 | July 14 2001 | LBC063 |
| <i>Dacrymyces palmatus</i> | on birch and alder/specimen missing | 12V | 486095.89 | 6582093.18 | July 10 2001 | LBC048 |
| <i>Dacrymyces deliquescens</i> | on rotten balsam poplar | 12V | 474126.65 | 6585095.84 | July 7 2001 | LBC030 |
| <i>Daedaleopsis confragosa</i> | on older fire killed log in black spruce bog | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC040 |
| <i>Daedaleopsis confragosa</i> | on white birch twig | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC076 |
| <i>Exidia glandulosa</i> | on alder | 12V | 477088.94 | 6584366.43 | July 14 2001 | LBC074 |
| <i>Exidia nucleata</i> | on dead alder | 12V | 486095.89 | 6582093.18 | July 10 2001 | LBC122 |
| <i>Fayodia pseudoclusilis</i> | on partly decomposed willow leaves | 12V | 474464.27 | 6587728.02 | July 6 2001 | LBC012 |
| <i>Fomes fomentarius</i> | on white birch | 12V | 474859.30 | 6587445.19 | July 6 2001 | LBC003 |
| <i>Fomes fomentarius</i> | on birch | 12V | 474919.40 | 6587555.91 | July 6 2001 | LBC008 |
| <i>Fomes fomentarius</i> | on alder | 12V | 477499.07 | 6584378.30 | July 10 2001 | LBC119 |
| <i>Fomitopsis cajanderi</i> | on rotting balsam poplar log | 12V | 474919.40 | 6587555.91 | July 6 2001 | LBC006a |
| <i>Fomitopsis pinicola</i> | on dead spruce log | 12V | 481025.28 | 6582607.45 | July 9 2001 | LBC039 |
| <i>Fomitopsis pinicola</i> | on spruce stump | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC078 |
| <i>Gandoderma applanatum</i> | on wood | 12V | 474729.26 | 6587026.10 | July 6 2001 | LBC004 |
| <i>Gleophyllum sepiarium</i> | on dead wood | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC053 |
| <i>Gleophyllum sepiarium</i> | on white spruce | 12V | 499258.40 | 6579539.65 | July 15 2001 | LBC108 |
| <i>Gleophyllum trabium</i> | on fire killed pine | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC046 |
| <i>Gymnopilus cf. humicola</i> | on wood | 12V | 477088.94 | 6584366.43 | July 14 2001 | LBC071 |
| <i>Hemimycena gracilis</i> | on moss in white spruce forest | 12V | 489206.86 | 6580689.96 | July 8 2001 | LBC026 |
| <i>Hygrocybe cantharellus</i> | in sphagnum | 12V | 493979.68 | 6577631.01 | July 15 2001 | LBC111 |
| <i>Hygrophoropsis aurantiaca</i> | on rotten log in swamp with calla and alder | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC086 |
| <i>Hygrotrama cf. hymenocephalum</i> | on sphagnum | 12V | 493979.68 | 6577631.01 | July 15 2001 | LBC109 |
| <i>Inocybe cf. calamistrata</i> | on willow | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC095 |
| <i>Inocybe cf. fastigata</i> | on sphagnum in black spruce bog | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC058 |
| <i>Inocybe cf. frigidula</i> | on ground in mixed white spruce/aspen stand | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC096 |
| <i>Inocybe cf. gymnocarpa</i> | in saline meadow with willow | 12V | 478813.83 | 6579354.52 | July 15 2001 | LBC116 |
| <i>Inocybe cf. petiginosa</i> | in moss in black spruce swamp | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC045 |

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|------------------------------------|---|-----------|-----------|------------|--------------|-----------------|
| <i>Inocybe mixtilis</i> | in moss with black spruce | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC055 |
| <i>Irpex lacteus</i> | on bark of dead alder | 12V | 491127.72 | 6580205.28 | July 8 2001 | LBC033 |
| <i>Irpex lacteus</i> | on wood | 12V | 477499.07 | 6584378.30 | July 15 2001 | LBC102 |
| <i>Leccinum insigne</i> | on edge of aspen stand and black spruce bog | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC037 |
| <i>Leccinum scabrum</i> | on bare ground on shoreline of La Butte Creek | 12V | 477783.17 | 6584510.43 | July 14 2001 | LBC062 |
| <i>Lentinus lepideus</i> | on dead/dry large driftwood logs | 12V | 474016.20 | 6585200.54 | July 7 2001 | LBC051 |
| <i>Lentinus lepideus</i> | on fire killed black spruce, in area with Labrador tea & wild red raspberry | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC081 |
| <i>Lentinus sulcatus</i> | on dead/dry wood | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC123 |
| <i>Leocarpus fragilis</i> | on lichen in moss | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC069 |
| <i>Lycogala epidendrum</i> | on wood | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC068 |
| <i>Marasmius pallidocephalus</i> | in moss in white spruce/white birch forest | 12V | 475063.66 | 6586400.20 | July 7 2001 | LBC020 |
| <i>Marasmius pallidocephalus</i> | on bits of spruce cone & needles | 12V | 474126.65 | 6585095.84 | July 7 2001 | LBC028 |
| <i>Marasmius scorodonius</i> | in sphagnum | 12V | 493979.68 | 6577631.01 | July 15 2001 | LBC114 |
| <i>Marasmius strictipes</i> | on wood in white spruce forest-willow area | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC060 |
| <i>Melastiza chateri</i> | in moss on mud of beaverdam | 12V | 489206.86 | 6580689.96 | July 8 2001 | LBC038 |
| <i>Mycena cf. citrinomarginata</i> | in moss in white spruce/white birch forest | 12V | 475063.66 | 6586400.20 | July 7 2001 | LBC021 |
| <i>Mycena cf. stipata</i> | [?] | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC057 |
| <i>Omphalina ericetorum</i> | on sphagnum in spruce bog | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC047 |
| <i>Omphalina ericetorum</i> | on moss in black spruce with sphagnum and rushes | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC052 |
| <i>Omphalina ericetorum</i> | in sphagnum | 12V | 493979.68 | 6577631.01 | July 15 2001 | LBC117 |
| <i>Oudemansiella longipes</i> | on wood in young aspen stand | 12V | 493979.68 | 6577631.01 | July 15 2001 | LBC118 |
| <i>Oudemansiella longipes</i> | in moss with alder | 12V | 478813.83 | 6579354.52 | July 15 2001 | LBC105 |
| <i>Oudemansiella longipes</i> | in moss with alder | 12V | 478813.83 | 6579354.52 | July 15 2001 | LBC105a |
| <i>Peniophora sp.</i> | on old spruce | 12V | 499258.40 | 6579539.65 | July 15 2001 | LBC115 |
| <i>Phellinus pini</i> | on old wood | 12V | 477499.07 | 6584378.30 | July 15 2001 | LBC101 |
| <i>Phellinus tremulae</i> | one meter from ground on dead balsam poplar | 12V | 474919.40 | 6587555.91 | July 7 2001 | LBC075 |
| <i>Phellinus tremulae</i> | on aspen poplar | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC077 |
| <i>Phyllotopsis nidulans</i> | on dead spruce branch | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC066 |
| <i>Piptoporus betulinus</i> | on dead birch | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC079 |

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|--|--|-----------|-----------|------------|--------------|-----------------|
| <i>Piptoporus betulinus</i> | on dead birch | 12V | 474853.33 | 6585541.35 | July 7 2001 | LBC080 |
| <i>Pleurotus ostreatus</i> | on balsam poplar | 12V | 474729.26 | 6587026.10 | July 6 2001 | LBC002 |
| <i>Plicaturopsis crispa</i> | on birch | 12V | 474919.40 | 6587555.91 | July 6 2001 | LBC009 |
| <i>Plicaturopsis crispa</i> | on alder | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC083 |
| <i>Plicaturopsis crispa</i> | on alder/ old, dry | 12V | 477499.07 | 6584378.30 | July 10 2001 | LBC121 |
| <i>Pluteus cervinus</i> | on rotting balsam poplar log | 12V | 474919.40 | 6587555.91 | July 6 2001 | LBC006 |
| <i>Pluteus pestatus</i> | growing out the end of a rotted birch log | 12V | 489206.86 | 6580689.96 | July 8 2001 | LBC025 |
| <i>Polyporus alveolaris</i> | on aspen | 12V | 477783.17 | 6584510.43 | July 14 2001 | LBC061a |
| <i>Polyporus elegans</i> | on alder | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC090 |
| <i>Polyporus squamosus</i> | on rotted stump | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC061 |
| <i>Polyporus varius</i> | on birch and rotting balsam poplar | 12V | 474919.40 | 6587555.91 | July 6 2001 | LBC007 |
| <i>Polyporus varius</i> | on balsam poplar | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC054 |
| <i>Polyporus varius forma nummularis</i> | on dead wood | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC085 |
| <i>Psathyrella cf. gracilis</i> | on leaf litter amongst alder at creek edge | 12V | 482191.49 | 6582371.79 | July 15 2001 | LBC104 |
| <i>Psathyrella cf. velutina</i> | on ground in alder flat | 12V | 475063.66 | 6586400.20 | July 7 2001 | LBC018 |
| <i>Psathyrella cf. velutina</i> | on ground in alder flat | 12V | 475063.66 | 6586400.20 | July 7 2001 | LBC019 |
| <i>Psathyrella subnuda</i> | on moss in white spruce stand | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC084 |
| <i>Psathyrella subnuda</i> | on ground in open dry meadow with aspen and willow | 12V | 478813.83 | 6579354.52 | July 15 2001 | LBC113 |
| <i>Pycnoporus cinnabarinus</i> | on dead alder | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC041 |
| <i>Rhodocybe caelata</i> | [?] | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC087 |
| <i>Rhodocybe caelata Cf.</i> | in black spruce bog | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC100 |
| <i>Russula cf. fragilis</i> | on alder bark in balsam poplar forest | 12V | 492236.38 | 6580251.83 | July 12 2001 | LBC099 |
| <i>Russula emetica</i> | on sphagnum in wet birch/alder swamp | 12V | 493979.68 | 6577631.01 | July 15 2001 | LBC110 |
| <i>Russula fragilis</i> | in alder swamp | 12V | 474126.65 | 6585095.84 | July 7 2001 | LBC022 |
| <i>Russula fragilis</i> | in white spruce stand | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC089 |
| <i>Russula fragilis</i> | on mossy log | 12V | 477499.07 | 6584378.30 | July 15 2001 | LBC103 |
| <i>Russula vesca</i> | on the shoreline of recently drawn down La Butte Creek | 12V | 477783.17 | 6584510.43 | July 14 2001 | LBC064 |
| <i>Sarcodon scabrosus</i> | under aspen /specimen moldy | 12V | 477499.07 | 6584378.30 | July 14 2001 | LBC073 |
| <i>Schizophyllum commune</i> | on dead wood | 12V | 485975.90 | 6581957.42 | July 10 2001 | LBC042 |
| <i>Scutellinia scutellata</i> | on mud of beaverdam | 12V | 489206.86 | 6580689.96 | July 8 2001 | LBC031 |
| <i>Stereum hirsutum</i> | rotted birch log | 12V | 474859.30 | 6587445.19 | July 6 2001 | LBC005 |
| <i>Trametes pubescens</i> | on wood | 12V | 486112.51 | 6581970.06 | July 10 2001 | LBC056 |
| <i>Tremella foliacea</i> | on dead alder | 12V | 486095.89 | 6582093.18 | July 10 2001 | LBC049 |
| <i>Tremella foliacea</i> | on alder | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC083a |
| <i>Trichaptum biforme</i> | on birch | 12V | 474919.40 | 6587555.91 | July 6 2001 | LBC013 |
| <i>Tubaria furfuracea</i> | on dead wood with moss in white spruce forest | 12V | 475891.84 | 6580230.43 | July 11 2001 | LBC088 |
| <i>Tubaria pellucida</i> | [?] | 12V | 475063.66 | 6586400.20 | July 7 2001 | LBC017 |

Appendix 3. Annotated species list for Fidler-Greywillow Wildland

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|--|---|-----------|-----------|------------|--------------|-----------------|
| <i>Agaricus bernardii</i> | on sand near balsam poplar | 12V | 494503 | 6513031 | July 24 2001 | FG070 |
| <i>Armillaria mellea</i> | on ground | 12V | 547141 | 6565461 | July 20 2001 | FG037 |
| <i>Astraeus hygrometricus</i> | on sand at upper beach level | 12V | 532885.51 | 6552112.57 | July 18 2001 | FG014 |
| <i>Astraeus hygrometricus</i> | in sand 2-4 cm deep | 12V | 547141 | 6565461 | July 20 2001 | FG035 |
| <i>Astraeus hygrometricus</i> | on sand of upper beach | 12V | 521564.95 | 6532104.53 | July 21 2001 | FG041 |
| <i>Bisporella citrina</i> | on rotting stick | 12V | 535656.32 | 6554192.38 | July 19 2001 | FG024 |
| <i>Collybia dryophila</i> | on moss in spruce forest | 12V | 535131.81 | 6553858.37 | July 19 2001 | FG019 |
| <i>Collybia dryophila</i> | on feather/feather moss | 12V | 535483.64 | 6554269.03 | July 20 2001 | FG031 |
| <i>Collybia dryophila</i> | on moss | 12V | 535483.64 | 6554269.03 | July 20 2001 | FG033 |
| <i>Coltricia perrenis</i> | on sand with reindeer lichen | 12V | 536660 | 6554586 | July 19 2001 | FG015 |
| <i>Coltricia perrenis</i> | on humus on sand | 12V | 547141 | 6565461 | July 20 2001 | FG040 |
| <i>Coltricia perrenis</i> | on sand along path, with lichen and blueberry | 12V | 536643 | 6555716 | July 22 2001 | FG049 |
| <i>Coltricia perrenis</i> | on sand | 12V | 514758 | 6517473 | July 24 2001 | FG061 |
| <i>Coriolopsis gallica</i> | on fire killed pine | 12V | 547141 | 6565461 | July 20 2001 | FG036 |
| <i>Cortinarius armillatus</i> | on ground in aspen/balsam poplar stand | 12V | 509615 | 6520911 | July 24 2001 | FG069 |
| <i>Cortinarius brunneus var. grandicolor</i> | on moss with black spruce | 12V | 535131.81 | 6553858.37 | July 19 2001 | FG020 |
| <i>Cortinarius brunneus var. grandicolor</i> | on sand at lake edge | 12V | 529525 | 6553460 | July 23 2001 | FG053 |
| <i>Cortinarius cf. casimiri</i> | on sand with rushes at spray line of lake | 12V | 536660 | 6554586 | July 19 2001 | FG017 |
| <i>Cortinarius cf. casimiri</i> | on sand with rushes at spray line of lake | 12V | 535131.81 | 6553858.37 | July 19 2001 | FG018 |
| <i>Cortinarius cf. jubarinus</i> | on sphagnum | 12V | 535483.64 | 6554269.03 | July 20 2001 | FG032 |
| <i>Cortinarius helobius</i> | on sand at lake edge | 12V | 529525 | 6553460 | July 23 2001 | FG053 |
| <i>Daedaleopsis confragosa</i> | on balsam poplar | 12V | 529525 | 6553460 | July 23 2001 | FG056 |
| <i>Dermocybe cinnamomeobadia</i> | in sphagnum | 12V | 509615 | 6520911 | July 24 2001 | FG063 |
| <i>Dermocybe uliginosa</i> | in moss on open bog | 12V | 535213.77 | 6553895.56 | July 18 2001 | FG010 |
| <i>Fomes fomentarius</i> | on birch | 12V | 532787.66 | 6552523.92 | July 17 2001 | FG008 |
| <i>Fomes fomentarius</i> | on birch | 12V | 533249 | 6552781 | July 26 2001 | FG071 |
| <i>Fomitopsis pinicola</i> | on black spruce and jackpine | 12V | 535131.81 | 6553858.37 | July 18 2001 | FG012 |
| <i>Fomitopsis pinicola</i> | on dead black spruce | 12V | 534057 | 6553874 | July 22 2001 | FG051 |
| <i>Fomitopsis pinicola</i> | on balsam poplar | 12V | 509615 | 6520911 | July 24 2001 | FG065 |
| <i>Galerina hypnorum</i> | on moss in black spruce stand | 12V | 532688.21 | 6552404.78 | July 17 2001 | FG002 |
| <i>Galerina paludosa</i> | on moss with black spruce | 12V | 535131.81 | 6553858.37 | July 19 2001 | FG022 |
| <i>Galerina paludosa</i> | on leaf litter in black spruce/aspen | 12V | 535092.25 | 6553951.79 | July 19 2001 | FG026 |
| <i>Galerina paludosa</i> | in sphagnum | 12V | 535483.64 | 6554269.03 | July 20 2001 | FG030 |
| <i>Gandoderma applanatum</i> | on wood | 12V | 533249 | 6552781 | July 26 2001 | FG072 |
| <i>Gandoderma applanatum</i> | on wood | 12V | 533249 | 6552781 | July 26 2001 | FG073 |
| <i>Gloeophyllum abietinum</i> | on dead black spruce | 12V | 537833 | 6555131 | July 22 2001 | FG046 |
| <i>Gymnopilus humicola</i> | on birch | 12V | 535131.81 | 6553858.37 | July 19 2001 | FG021 |

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|----------------------------------|---|-----------|-----------|------------|--------------|-----------------|
| <i>Hebeloma cf. velatum</i> | edge of pine and creek | 12V | 535656.32 | 6554192.38 | July 19 2001 | FG027 |
| <i>Inocybe cf. frigidula</i> | on sand | 12V | 534970.30 | 6553653.18 | July 20 2001 | FG028 |
| <i>Laccaria laccata</i> | on leaf litter in black spruce/aspen | 12V | 535483.64 | 6554269.03 | July 20 2001 | FG029 |
| <i>Laccaria proxima</i> | on sand | 12V | 537833 | 6555131 | July 22 2001 | FG047 |
| <i>Lactarius cf. obscuratus</i> | in leaf litter | 12V | 547141 | 6565461 | July 20 2001 | FG038 |
| <i>Lactarius fragilis</i> | in shade of upturned stump, on moss in sand | 12V | 534057 | 6553874 | July 22 2001 | FG050 |
| <i>Leccinum insigne</i> | with aspen poplar, occasional birch and balsam poplar | 12V | 509615 | 6520911 | July 24 2001 | FG064 |
| <i>Leccinum scabrum</i> | in moss under aspen/birch/white spruce | 12V | 529525 | 6553460 | July 23 2001 | FG054 |
| <i>Leptista (=Clitocybe) sp.</i> | on moss | 12V | 535213.77 | 6553895.56 | July 18 2001 | FG013 |
| <i>Mycena stannea</i> | on moss | 12V | 547141 | 6565461 | July 20 2001 | FG039 |
| <i>Omphalina ericetorum</i> | in spruce bog with liverworts, sphagnum & other mosses | 12V | 532885.51 | 6552112.57 | July 17 2001 | FG001 |
| <i>Omphalina ericetorum</i> | on moss with black spruce (with FG022 - <i>Galerina paludosa</i> Fr.) | 12V | 535131.81 | 6553858.37 | July 19 2001 | FG074 |
| <i>Omphalina pyxidata</i> | on moss in black spruce stand | 12V | 532688.21 | 6552404.78 | July 17 2001 | FG006 |
| <i>Peziza badia</i> | growing on sand with moss & liverworts | 12V | 536660 | 6554586 | July 19 2001 | FG016 |
| <i>Phellinus tremulae</i> | on balsam poplar | 12V | 509615 | 6520911 | July 24 2001 | FG066 |
| <i>Piptoporus betulinus</i> | on birch | 12V | 509615 | 6520911 | July 24 2001 | FG068 |
| <i>Pluteus cervinus</i> | on birch log | 12V | 532688.21 | 6552404.78 | July 17 2001 | FG004 |
| <i>Pluteus cervinus</i> | on alder | 12V | 533081.88 | 6552783.92 | July 17 2001 | FG005 |
| <i>Polyporus brumalis</i> | on willow | 12V | 535656.32 | 6554192.38 | July 19 2001 | FG023 |
| <i>Polyporus elegans</i> | on birch or alder | 12V | 535131.81 | 6553858.37 | July 18 2001 | FG009 |
| <i>Polyporus elegans</i> | on wood | 12V | 535131.81 | 6553858.37 | July 18 2001 | FG011 |
| <i>Polyporus elegans</i> | on alder | 12V | 529525 | 6553460 | July 23 2001 | FG055 |
| <i>Polyporus</i> sp. | on sand | 12V | 535483.64 | 6554269.03 | July 20 2001 | FG034 |
| <i>Russula cf. maculata</i> | on ground | 12V | 535656.32 | 6554192.38 | July 19 2001 | FG025 |
| <i>Russula emetica</i> | on sphagnum in black spruce bog | 12V | 537185 | 6555035 | July 22 2001 | FG048 |
| <i>Thelephora terrestris</i> | on moss | 12V | 532885.51 | 6552112.57 | July 17 2001 | FG003 |
| <i>Trichaptum biforme</i> | on birch | 12V | 532787.66 | 6552523.92 | July 17 2001 | FG007 |
| <i>Trichaptum biforme</i> | on balsam poplar | 12V | 509615 | 6520911 | July 24 2001 | FG067 |

Appendix 4. Annotated species list for Colin-Cornwall Lakes Wildland

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|---------------------------------|---|-----------|-----------|------------|---------------|-----------------|
| <i>Agrocybe praecox</i> | on sand under burnt pine | 12V | 546429 | 6605262 | July 8, 2002 | CC074 |
| <i>Agrocybe putaminum</i> | on soil in open pine forest | 12V | 546429 | 6605262 | July 8, 2002 | CC073 |
| <i>Chondrostereum purpureum</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC029 |
| <i>Collybia confusa</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 7, 2002 | CC085 |
| <i>Collybia dryophila</i> | with feather moss | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC009 |
| <i>Collybia dryophila</i> | on rotted wood | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC044 |
| <i>Coprinus lagopoides</i> | on burnt wood | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC036 |
| <i>Coriolopsis gallica</i> | on fire killed pine | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC056 |
| <i>Coriolopsis gallica</i> | on dead conifer | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC061 |
| <i>Coriolopsis gallica</i> | on dead pine | 12V | 551063 | 6608253 | July 8, 2002 | CC078 |
| <i>Cortinarius obtusus</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 7, 2002 | CC084 |
| <i>Cortinarius obtusus</i> | with moss under black spruce | 12V | 546605.00 | 6599986.00 | July 9, 2002 | CC099 |
| <i>Fomes fomentarius</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC013 |
| <i>Fomes fomentarius</i> | on black spruce | 12V | 541646.42 | 6602125.3 | July 9, 2002 | CC103 |
| <i>Fomitopsis cajanderi</i> | on fire killed pine | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC054 |
| <i>Fomitopsis cajanderi</i> | on dead wood | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC059 |
| <i>Fomitopsis pinicola</i> | on dead conifer | 12V | 541646.42 | 6602125.3 | July 6, 2002 | CC003 |
| <i>Fomitopsis pinicola</i> | on dead conifer | 12V | 541646.42 | 6602125.3 | July 6, 2002 | CC005 |
| <i>Fomitopsis pinicola</i> | on dead white spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC010 |
| <i>Fomitopsis pinicola</i> | on birch | 12V | 550734.03 | 6601468.7 | July 9, 2002 | CC100 |
| <i>Fomitopsis rosea</i> | on rotted log | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC030 |
| <i>Fomitopsis rosea</i> | on cut pine stump | 12V | 546429 | 6605262 | July 8, 2002 | CC075 |
| <i>Galerina cinctula</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 8, 2002 | CC081 |
| <i>Galerina mniophila</i> | with moss under black spruce | 12V | 546605.00 | 6599986.00 | July 9, 2002 | CC097 |
| <i>Gandoderma applanatum</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC035 |
| <i>Gleophyllum protractum</i> | on fire killed pine | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC057 |
| <i>Gleophyllum sepiarium</i> | on fire killed pine | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC058 |
| <i>Gleophyllum sepiarium</i> | on fire killed conifer | 12V | 540362.00 | 6600468.00 | July 7, 2002 | CC066 |
| <i>Gleophyllum sepiarium</i> | on dead conifer | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC008 |
| <i>Gleophyllum trabium</i> | on fire killed black spruce | 12V | 543755.00 | 6602888.00 | July 8, 2002 | CC072 |
| <i>Gleoporus taxicola</i> | on unpeeled pine logs of cabin | 12V | 546610 | 6600000.6 | July 10, 2002 | CC105 |
| <i>Gyromitra esculenta</i> | on wood on ground | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC032 |
| <i>Gyromitra esculenta</i> | on burnt soil | 12V | 540362.00 | 6600468.00 | July 7, 2002 | CC064 |
| <i>Gyromitra esculenta</i> | on buried wood in burnt pine forest | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC108 |
| <i>Hebeloma cf. longicaudum</i> | with moss under black spruce | 12V | 546605.00 | 6599986.00 | July 9, 2002 | CC096 |
| <i>Hygrophorus sp.</i> | with moss under black spruce (specimen missing) | 12V | 551063 | 6608253 | July 8, 2002 | CC086 |
| <i>Inocybe lacera</i> | on ground in burnt pine forest | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC014 |
| <i>Inonotus obliquus</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC017 |
| <i>Irpex lacteus</i> | on fire killed birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC023 |
| <i>Irpex lacteus</i> | on dead wood | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC037 |
| <i>Kuehneromyces lignicola</i> | on burnt ground with moss | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC062 |
| <i>Kuehneromyces lignicola</i> | on sand under burnt pine | 12V | 546429 | 6605262 | July 8, 2002 | CC111 |
| <i>Lycogala epidendrum</i> | on burnt black spruce | 12V | 543755.00 | 6602888.00 | July 8, 2002 | CC070a |

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|---|---|--------|-----------|------------|---------------|--------------|
| <i>Marasmius scorodonius</i> | on burnt soil with moss | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC041 |
| <i>Morchella elata</i> | on burnt ground, under white spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC027 |
| <i>Mycena cf. galopus</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 8, 2002 | CC080 |
| <i>Mycena filipes</i> | with moss under black spruce | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC052 |
| <i>Mycena leucogala</i> | on burnt soil | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC042 |
| <i>Myriosclerotinia caricic-ampullaceae</i> | on Carex sp. | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC053 |
| <i>Neobulgaria pura</i> | on burnt black spruce | 12V | 543755.00 | 6602888.00 | July 8, 2002 | CC070b |
| <i>Omphalina ericetorum</i> | with moss under black spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC046 |
| <i>Omphalina ericetorum</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 8, 2002 | CC082 |
| <i>Peziza praetervisa</i> | on burnt soil | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC011 |
| <i>Phellinus chrysoluma</i> | on fire killed pine | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC055 |
| <i>Phellinus chrysoluma</i> | on dead white spruce | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC063 |
| <i>Phellinus chrysoluma</i> | on unpeeled pine logs of cabin | 12V | 546610 | 6600000.6 | July 10, 2002 | CC104 |
| <i>Phellinus ignarius</i> | on dead wood | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC048 |
| <i>Phellinus ignarius</i> | on birch | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC049 |
| <i>Phellinus ignarius</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC033 |
| <i>Phellinus ignarius</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC034 |
| <i>Phellinus ignarius</i> | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC087 |
| <i>Phellinus tremulae</i> | on aspen popular | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC015 |
| <i>Pholiota spumosa</i> | on soil | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC021 |
| <i>Pholiota spumosa</i> | on soil | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC022 |
| <i>Pholiota spumosa</i> | on wood in burnt ground, under white spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC026 |
| <i>Pholiota spumosa</i> | on wood in burnt ground, under white spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC028 |
| <i>Pholiota spumosa</i> | on burnt soil | 12V | 540362.00 | 6600468.00 | July 7, 2002 | CC065 |
| <i>Pholiota spumosa</i> | on burnt soil with moss | 12V | 543755.00 | 6602888.00 | July 8, 2002 | CC071 |
| <i>Pholiota spumosa</i> | on buried wood | 12V | 551063 | 6608253 | July 8, 2002 | CC077 |
| <i>Piptoporus betulinus</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC020 |
| <i>Polyporus elegans</i> | on dead wood | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC047 |
| <i>Polyporus elegans</i> | on birch | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC050 |
| <i>Polyporus elegans</i> | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC088 |
| <i>Polyporus melanopus</i> | on rotted log | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC019 |
| <i>Polyporus varius</i> | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC093 |
| <i>Polyporus varius</i> | on birch | 12V | 546610 | 6600000.6 | July 10, 2002 | CC106 |
| <i>Psathyrella cf. phegophila</i> | on burnt ground, under white spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC025 |
| <i>Pycnoporus cinnabarinus</i> | on birch | 12V | 540362.00 | 6600468.00 | July 7, 2002 | CC068 |
| <i>Sarcodon scabrosus</i> | on ground | 12V | 541646.42 | 6602125.3 | July 6, 2002 | CC006 |
| <i>Skeletocutis amorphia</i> | on alder | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC018 |
| <i>Skeletocutis amorphia</i> | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC092 |
| <i>Skeletocutis amorphia</i> | on burnt alder | 12V | 541646.42 | 6602125.3 | July 9, 2002 | CC101 |
| Specimen Missing | burnt soil | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC038 |
| Specimen Unidentifiable | on dead spruce | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC031 |
| <i>Stemonitis splendens</i> | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC091 |
| <i>Stereum hirsutum</i> | on birch | 12V | 540362.00 | 6600468.00 | July 7, 2002 | CC067 |

| Species | Habitat | NAD 83 | Easting | Northing | Date | Specimen No. |
|----------------------------------|------------------------------|-----------|-----------|------------|---------------|-----------------|
| <i>Stereum striatum</i> | on birch | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC060 |
| <i>Thelephora terrestris</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 8, 2002 | CC083 |
| <i>Trametes cervina</i> | log in cabin wall | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC016 |
| <i>Trametes hirsuta</i> | on dead black spruce | 12V | 546605.00 | 6599986.00 | July 9, 2002 | CC095 |
| <i>Trametes hirsuta</i> | on fire killed pine | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC110 |
| <i>Trametes suaveolens</i> | on birch | 12V | 541646.42 | 6602125.3 | July 9, 2002 | CC102 |
| <i>Trametes versicolor</i> | on dead pine | 12V | 546429 | 6605262 | July 8, 2002 | CC076 |
| <i>Trichaptum abietinum</i> | on white spruce | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC090 |
| <i>Trichaptum biforme</i> | on fire killed birch | 12V | 541646.42 | 6602125.3 | July 6, 2002 | CC001 |
| <i>Trichaptum biforme</i> | on birch | 12V | 541646.42 | 6602125.3 | July 6, 2002 | CC002 |
| <i>Trichaptum biforme</i> | on dead wood | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC007 |
| <i>Trichaptum biforme</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC012 |
| <i>Trichaptum biforme</i> | on fire killed birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC024 |
| <i>Trichaptum biforme</i> | on birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC045 |
| <i>Trichaptum biforme</i> | on fire killed birch | 12V | 540362.00 | 6600468.00 | July 7, 2002 | CC069 |
| <i>Trichaptum biforme</i> | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC094 |
| <i>Tubaria hiemalis</i> | balsam poplar on the ground | 12V | 541646.42 | 6602125.3 | July 6, 2002 | CC004 |
| <i>Tubaria hiemalis</i> | on wood | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC051 |
| <i>Tubaria romagnesiana</i> | on rotted birch | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC039 |
| <i>Tubaria romagnesiana</i> | on burnt soil | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC040 |
| Unidentifiable | on birch | 12V | 529639.00 | 6600841.00 | July 9, 2002 | CC089 |
| Unidentifiable yellow slime | on sphagnum | 12V | 541366.00 | 6599947.00 | July 10, 2002 | CC109 |
| <i>Xeromphalina fraxinophila</i> | on burnt soil with moss | 12V | 541742.77 | 6602023.6 | July 6, 2002 | CC043 |
| <i>Xeromphalina fraxinophila</i> | with moss under black spruce | 12V | 551063 | 6608253 | July 8, 2002 | CC079 |
| <i>Xeromphalina fraxinophila</i> | with moss under black spruce | 12V | 546605.00 | 6599986.00 | July 9, 2002 | CC098 |
| <i>Xeromphalina fraxinophila</i> | with moss under black spruce | 12V | 541366.00 | 6599947.00 | July 7, 2002 | CC107 |

Appendix 5. Partial list of literature accounts of fungi occurrences in Alberta.

- Abbott, L. L. 1990. Ministik Lake Foray. The Stinkhorn, 4 (1):17. Edmonton.
- Abbott, L. L. 1991. Ministik Lake Foray 1990. The Stinkhorn, 5 (1):11. Edmonton.
- Abbott, L. L. 1991. Shaw Lake Foray 1990. The Stinkhorn, 5 (1):20-21. Edmonton.
- Abbott, S. P. 1987. A Return Foray. The Stinkhorn, 1 (1):12. Edmonton.
- Abbott, S. P. 1987. EMS Display at the Provincial Museum. The Stinkhorn, 4 (1):29-31. Edmonton.
- Abbott, S. P. 1987. EMS Mushroom Display. The Stinkhorn, 1 (1):19-20. Edmonton.
- Abbott, S. P. 1987. Foray at the Botanic Garden. The Stinkhorn, 1 (1):14-15. Edmonton.
- Abbott, S. P. 1987. The First Foray. The Stinkhorn, 1 (1):13. Edmonton.
- Abbott, S. P. 1989. Key to Boleti of Alberta. University of Alberta, Edmonton. 12pp. Mimeograph.
- Abbott, S. P. 1990. *Amanita muscaria* Poisonings. The Stinkhorn, 4 (1): 21. Edmonton.
- Abbott, S. P. 1990. *Entoloma Grande* Poisonings. The Stinkhorn, 4 (1): 20. Edmonton.
- Abbott, S. P. 1990. Pet Poisonings -- A Narrative.... The Stinkhorn, 5 (1): 22. Edmonton.
- Abbott, S. P. 1990. Shaw Lake. The Stinkhorn, 4 (1): 38-42. Edmonton.
- Abbott, S. P. 1990. Stinkhorns. The Stinkhorn, 4 (1): 32. Edmonton.
- Abbott, S. P. and L. L. Abbott. 1991. Alberta *Verpa* Report. The Stinkhorn, 5 (1): 13-18. Edmonton.
- Abbott, S. P. and R. S. Currah. 1989. The larger cup fungi and other Ascomycetes of Alberta, an annotated checklist. Devonian Botanic Garden, University of Alberta, Edmonton. 96 pp.
- Abbott, S.P. & R.S. Currah. 1997. The Helvellaceae: Systematic revision and occurrence in northern and northwestern North America. Mycotaxon 62:1-125.
- Abbott, S.P. 1992. Systematic studies of the Helvellaceae in northern and northwestern North America. University of Alberta. (Canadian Theses Service microfiche TH 77275 National Library)
- Abbott, S.P. and R.S. Currah. 1997. The Helvellaceae: systematic revision and occurrence in northern and northwestern North America. Mycotaxon 62: 1-125.
- Abbott, S.P., and Currah. R.S. 1988. The genus *Helvella* in Alberta. Mycotaxon 33: 229-250.
- Addy, H.D., Hambleton, S. and Currah, R.S. 2000. Distribution and molecular characterization of the root endophyte *Phialocephala fortinii* along an environmental gradient in the boreal forest of Alberta. Mycol. Res. 104: 1213-1221.
- Anonymous 1991. Stinkhorn Notes. The Stinkhorn, 5 (1):40. Edmonton
- Baranyay, J. A. 1967. Notes on Hypoxylon canker of aspen in Alberta. For. Chron. 43:372-280.
- Baranyay, J.A. 1968. Fungi collected during forest diseases surveys in northern Alberta and the district of Mackenzie, Northwest Territories. For Branch Dept. Public no. 1238, 25p.
- Boyetchko, S.M, and J. P. Tewari. 1993. Occurrence of vesicular-arbuscular mycorrhizal fungi in Alberta, Canada. Zeitschrift Fur Naturforschung C-A Journal of Biosciences. 48(11-12):923-929.
- Bradbury, SM. 1998. Ectomycorrhizas of lodgepole pine (*Pinus contorta*) seedlings originating from seed in southwestern Alberta cut blocks. Can. J. Bot.- Rev. Can. Bot. 76: 213-217.
- Brodie, H. J. 1966. A new species of *Cyathus* from the Canadian Rockies. Canadian J. Bot. 44:1235-1237.
- Brodie, H. J. 1968. The Nidulariaceae of Canada. Canadian Field-Naturalist 82: 2-14

- Brodie, H.J. 1970. A previously unnamed species of Cyathus from the Cypress Hills, Alberta. Canadian Journal of Botany. 48: 749-750
- Carmichael, J. W., W. B. Kendrick, I. L. Connerse and L. Sigler. 1980. Genera of Hyphomycetes. The Univ. of Alberta Press, Edmonton, Canada. 386p.
- Cormack, M.W. 1948. Winter crown rot or snow mold of alfalfa, clovers, and grasses in Alberta. Can. J. Res. (Sec. C) 26:71—85.
- Currah RS; Smreciu EA; Lehesvirta T; Niemi M; Larsen KW. 2000. Fungi in the winter diets of northern flying squirrels and red squirrels in the boreal mixedwood forest of northeastern Alberta. Canadian Journal of Botany-Revue Canadienne de Botanique. 78(12):1514-1520.
- Currah, R. 1986. A new species of Ascodesmis from Alberta. Mycologia 78: 198-201.
- Danielson R.M. 1984. Ectomycorrhizal associations in jack pine stands in northeastern Alberta. Canadian Journal of Botany 62: 932-939.
- Danielson, R. M. (no date). Key to Mushrooms and their allies found in Alberta. University of Calgary. Mimeo. 66p.
- Danielson, R.M. 1979. Hypogeous ascomycetes in Alberta Canada, with two new North American records. Mycotaxon, 9:445-450
- Danielson, R.M., and S. Visser. 1989. Host response to inoculation and behaviour of introduced and indigenous ectomycorrhizal fungi of jack pine grown on oil-sands tailings. Canadian Journal of Forest Research vol. 19: 1412-1421
- Danielson, R.M. 1984. Ectomycorrhizal associations in jackpine stands in northeastern Alberta, Can. J. Bot. 62:932-939.
- Gilbertson, R. L. and L. Ryvarden. 1986. North American Polypores Vol. 1. Fungiflora, Oslo, Norway. 433p.
- Gilbertson, R. L. and L. Ryvarden. 1987. North American Polypores Vol. II. Fungiflora, Oslo, Norway. 885p.
- Gilbertson, R.L. 1981. North American wood-rotting fungi that cause brown rots. Mycotaxon 12:372—416.
- Ginns, J. 1989. Descriptions and notes for some unusual North American corticioid fungi (Aphyllophorales, Corticiaceae). Memoirs N.Y. Botanical Garden 49: 129-137.
- Hambleton, S., Huhtinen, and S. Currah, R.S. 1999. *Hymenoscyphus ericae*: a new record from western Canada. Mycol. Res. 103:1391-1397.
- Hiratsuka, Y., A.A. Loman. 1984. Decay of aspen and balsam poplar in Alberta. Can. For. Res, Cent. North. For. Res. Cent. Edmonton Alberta. Inf. Rep. NOR-X-262.
- Hiratsuka, Y., Gibbard, D.A., Bakowsky, O., and Maier, G.B. 1990. Classification and measurement of aspen decay and stain in Alberta. For. Can. Northwest Reg., North. For. Cent., Edmonton, Alberta. Inf. Rep. NOR-X-314. 29p.
- Hiratsuka, Y., P. J. Maruyama. 1985. Western gall rusts. Can. For. Res, Cent. North. For. Res. Cent. Edmonton Alberta. Pest Leafl. NFRC PL 27-85.
- Hutchison, L.J. 1999. Wood-inhabiting microfungi isolated from *Populus tremuloides* from Alberta and northeastern British Columbia. Can. J. Bot. 77: 898-905.
- Hutchison, L.J. 1999. Wood-inhabiting microfungi isolated from *Populus tremuloides* from Alberta and northeastern Brotidh Columbia. Can. J. Bot. 77: 898-905.
- Kernaghan, G. and Currah, R.S. 1998. Ectomycorrhizal fungi at tree line in the Canadian Rockies. Mycotaxon 69: 39-80.
- Kernaghan, G., Currah, R.S. and Bayer, R.J. 1997. Russulaceous ectomycorrhizae of *Abies lasiocarpa* and *Picea engelmannii*. Canadian Journal of Botany 75: 1843-1850.

- Larsen, H. J. and W. C. Denison. 1978. A checklist of operculate cup-fungi (Pezizales) of North America west of the Great Plains. *Mycotaxon* 7:68-90.
- Lawrence J.J. and Hiratsuka. 1972. Forest fungi collection in Banff National Park, Alberta. Information Report NOR-X-22, Northern Forest Research Centre.
- Lawrence J.J. and Hiratsuka. 1972. Forest fungi collection in Jasper National Park, Alberta. Information Report NOR-X-21, Northern Forest Research Centre.
- Lawrence J.J. and Hiratsuka. 1972. Forest fungi collection in Waterton Lakes National Park, Alberta. Information Report NOR-X-20, Northern Forest Research Centre.
- Loman, A. A., G. Paul. 1963. Decay of lodgepole pine in two foothills sections of the boreal forest in Alberta. *For. Chron.* 39:422-435
- Lowe J.L. & R.L. Gilbertson. 1961, Synopsis of Polyporaceae of the Western United States and Canada. *Mycoloia* vol. 53.
- Lumley, T.C., S.P. Abbott & R.C. Currah. 2000. Microscopic ascomycetes isolated from rotting wood in the boreal forest. *Mycotaxon* 74:395-414.
- Marcia Monreal, S.M. Berch, and Mary Berbee. 1999. Molecular diversity of ericoid mycorrhizal fungi. *Can. J. Bot.* 77: 1580-1594.
- Martin K.J. & R. L. Gilbertson 1977. Synopsis of wood-rotting fungi on spruce in North America. *Mycotaxon* 6: 43-77.
- Mugala, M.S., P.V. Blenis, Y. Hiratsuka, and K.I. Mallet. 1989. Infection of lodgepole pine and white spruce by Alberta isolates of *Armillaria*. *Canad. J. For. Res.* 19: 685-689.
- Parmelee, J. A. 1971. The genus *Gymnosporangium* in western Canada. *Can. J. Bot.* 49:903-926
- Parmelee, J.A. 1984. Microfungi parasitic on vascular plants in Waterton Lakes National Park, Alberta, and environs. *Agriculture Canada Tech. Bull.* 1984-11E, Ottawa. 32 pp.
- Paul, G. and Etheridge, D.E. 1958. Decay of aspen (*Populus tremuloides* Michx.) and balsam poplar (*P. balsamifera* L.) in the Lesser Slave Lake region of Alberta. Alberta Dep. Land For., For. Serv. Div., Edmonton, Alberta, and Can. Dep. Agric., For. Biol. Div., V+Calgary, Alberta, Mimeograph.
- Redhead, S.A. 1984. Arrhenia and Rimbachia, expanded generic concepts, and a reevaluation of *Leptoglossum* with emphasis on muscicolous North American taxa. *Canad. J. Bot.* 62: 865-892.
- Redhead, S.A. 1989. A biogeographical overview of the Canadian mushroom flora. *Canad. J. Bot.* 67: 3003-3062.
- Robinson-Jeffrey, C., A. A. Loman. 1963. Fungi Isolated in culture from red heartwood stain and advanced decay of lodgepole pine in Alberta. *Can. J. Bot.* 41:1371-1375.
- Schalkwijk, H.M.E. 1975. Mushrooms of the Edmonton Area- edible and poisonous. Printed and published in Edmonton 34pp.
- Schalkwijk, H.M.E. 1989. Leni Schalkwijk's checklist of Alberta Fungi. The Stinkhorn, Edmonton 3 (1): 1-66.
- Schalkwijk-Barendsen, H.M.E. 1991. Checklist of fungi to accompany the book *Mushrooms of Western Canada*. Schalkwijk-Barendsen, Edmonton, Alta. published by author.
- Schalkwijk-Barendsen, H.M.E. 1991. *Mushrooms of western Canada*. Lone Pine Publishing, Edmonton. 414 pp.
- Schalkwyk, H.M.E. 1987. Notes from members. The Stinkhorn. 1 (1): 23. Edmonton
- Schalkwyk, H.M.E. 1989. Leni Schalkwyk's checklist of Alberta Fungi. The Stinkhorn. 3 (1): 1-66. Edmonton
- Schulz, M. 2002. Common Wood Decay Fungi And Polypores of the Edmonton Area. Mycologist, Department of Biological Science. University of Alberta. Edmonton Alberta. Mimeograph
- Sigler, L. and A.L. Flis. 1998. University of Alberta microfungus collection and herbarium (UAMH) Devonian Botanic Garden, Edmonton, AB, Canada.

- Stoyke, G., and Currah, R.S. 1991. Endophytic fungi from the mycorrhizae of alpine ericoid plants. Can. J. Bot. 69: 347–352.
- Thomas, G. P. , D. E. Etheridge, G. Paul. 1960. Fungi and decay in aspen and balsam poplar in the boreal forest region, Alberta. Can J. Bot.38:549-466.
- Thormann MN; Currah RS; Bayley SE. 2002. The relative ability of fungi from Sphagnum fuscum to decompose selected carbon substrates. CANADIAN JOURNAL OF MICROBIOLOGY. 48(3):204-211.
- Traquair, J.A. 1980. The first Mary Elliot Mycological foray. Alberta Naturalist. 10: 4-6
- Tsuneda, A., Thormann, M.N., and R.S. Currah. 2000. *Scleroconidioma* , a new genus of dematiaceous Hyphomycetes. Can. J. Bot. 78:1294-1298.
- Zalasky, H. 1964. Nomenclature and description of *Diplodia tumefaciens* (Shear) Zalasky (=Macrophoma tumefaciens Shear apud Hubert). Can. J. Bot. 42:1049-1055.
- Ziller, W. G. 1974. The Tree Rusts of Western Canada. Canadian Forestry Service Pub. No. 1329. Pacific Forest Research Centre, Victoria, B. C. 272p.

Appendix 6. Photographs of macrofungi from La Butte Creek, Fidler-Greywillow and Colin-Cornwall Lakes Wildland Provincial Parks



LB061 *Polyporus squamosus*, La Butte Creek Wildland Provincial Park.
Photographer: Drajs Vujnovic



FG034 *Polyporus* sp.? , [Specimen parasitized and with hymenium eaten off (?)] Fidler-Greywillow Wildland Provincial Park.

Photographer: Drajs Vujnovic



FG053 *Cortinarius brunneus* var. *grandicolor*, on sand at lake edge, Fidler-Greywillow Wildland Provincial Park.

Photographer: Lorna Allen



FG024 *Scutellinia scutellata*, on mud of beaver dam, Fidler-Greywillow Wildland Provincial Park.

Photographer: Lorna Allen



FG016 *Peziza badia*, Fidler-Greywillow Wildland Provincial Park.

Photographer: Lorna Allen



FG001 *Omphalina ericetorum*, Fidler-Greywillow Wildland Provincial Park.
Photographer: Lorna Allen



LB082 *Coriolopsis gallica*, La Butte Creek Wildland Provincial Park.
Photographer: Lorna Allen



LB074 *Exidea glandosa*, La Butte Creek Wildland Provincial Park.
Photographer: Lorna Allen



Laccaria sp., La Butte Creek Wildland Provincial Park.
Photographer: Lorna Allen



Fomitopsis pinicola On black spruce, Colin-Cornwall Lakes
Wildland Provincial Park
Photographer: Drajs Vujnovic



Gyromitra esculenta, Colin-Cornwall Lakes Wildland Provincial Park
Photographer: Drajs Vujnovic